

Classifying cubic s -regular graphs of orders $22p$ and $22p^2$

A. A. Talebi and N. Mehdipoor

Communicated by Z. Marciniak

ABSTRACT. A graph is s -regular if its automorphism group acts regularly on the set of s -arcs. In this study, we classify the connected cubic s -regular graphs of orders $22p$ and $22p^2$ for each $s \geq 1$, and each prime p .

1. Introduction

In this study, all graphs considered are assumed to be undirected, finite, simple, and connected, unless stated otherwise. For a graph X , $V(X)$, $E(X)$, $Arc(X)$, and $Aut(X)$ denote its vertex set, edge set, arc set, and full automorphism group, respectively. Let G be a subgroup of $Aut(X)$. For $u, v \in V(X)$, uv denotes the edge incident to u and v in X , and $N_X(u)$ denotes the neighborhood of u in X , that is, the set of vertices adjacent to u in X .

A graph \tilde{X} is called a covering of a graph X with projection $p : \tilde{X} \rightarrow X$ if there is a surjection $p : V(\tilde{X}) \rightarrow V(X)$ such that $p|_{N_{\tilde{X}}(\tilde{v})} : N_{\tilde{X}}(\tilde{v}) \rightarrow N_X(v)$ is a bijection for any vertex $v \in V(X)$ and $\tilde{v} \in p^{-1}(v)$.

A permutation group G on a set Ω is said to be semiregular if the stabilizer G_v of v in G is trivial for each $v \in \Omega$, and is regular if G is transitive, and semiregular.

Let K be a subgroup of $Aut(X)$ such that K is intransitive on $V(X)$. The quotient graph X/K induced by K is defined as the graph such that

2000 MSC: 05C25, 20b25.

Key words and phrases: s -regular graphs, s -arc-transitive graphs, symmetric graphs, regular covering.

the set Ω of K -orbits in $V(X)$ is the vertex set of X/K and $B, C \in \Omega$ are adjacent if and only if there exist a $u \in B$ and $v \in C$ such that $\{u, v\} \in E(X)$.

A covering \tilde{X} of X with a projection p is said to be regular (or N -covering) if there is a semiregular subgroup N of the automorphism group $Aut(\tilde{X})$ such that graph X is isomorphic to the quotient graph \tilde{X}/N , say by h , and the cubic map $\tilde{X} \rightarrow \tilde{X}/N$ is the composition ph of p and h (in this paper, all functions are composed from left to right). If N is a cyclic or an elementary Abelian, then, \tilde{X} is called a cyclic or an elementary Abelian covering of X , and if \tilde{X} is connected, N becomes the covering transformation group.

An s -arc in a graph X is an ordered $(s+1)$ -tuple (v_0, v_1, \dots, v_s) of vertices of X such that v_{i-1} is adjacent to v_i for $1 \leq i \leq s$, and $v_{i-1} \neq v_{i+1}$ for $1 \leq i < s$; in other words, a directed walk of length s that never includes a backtracking. For a graph X and a subgroup G of $Aut(X)$, X is said to be G -vertex-transitive, G -edge-transitive, or G - s -arc-transitive if G is transitive on the sets of vertices, edges, or s -arcs of X , respectively, and G - s -regular if G acts regularly on the set of s -arcs of X . A graph X is said to be vertex-transitive, edge-transitive, s -arc-transitive, or s -regular if X is $Aut(X)$ -vertex-transitive, $Aut(X)$ -edge-transitive, $Aut(X)$ - s -arc-transitive, or $Aut(X)$ - s -regular, respectively. In particular, 1-arc-transitive means arc-transitive, or symmetric.

Covering techniques have long been known as a powerful tool in topology, and graph theory. Regular covering of a graph is currently an active topic in algebraic graph theory. Tutte [17, 18] showed that every finite cubic symmetric graph is s -regular for some $s \geq 1$, and this s is at most five. It follows that every cubic symmetric graph has an order of the form $2mp$ for a positive integer m and a prime number p . In order to know all cubic symmetric graphs, we need to classify the cubic s -regular graphs of order $2mp$ for a fixed positive integer m and each prime p . Conder and Dobcsányi [4, 5] classified the cubic s -regular graphs up to order 2048 with the help of the ‘‘Low index normal subgroups’’ routine in MAGMA system [2]. Cheng and Oxley [3] classified the cubic s -regular graphs of order $2p$. Recently, by using the covering technique, cubic s -regular graphs with order $2p^2$, $2p^3$, $4p$, $4p^2$, $6p$, $6p^2$, $8p$, $8p^2$, $10p$, $10p^2$, $12p$, $12p^2$, $14p$ and $16p$ were classified in [1, 7 – 12, 15, 16].

In this paper, by employing the covering technique, and group-theoretical construction, we investigate connected cubic s -regular graphs of orders $22p$ and $22p^2$ for each $s \geq 1$, and each prime p .

2. Preliminaries

We start by introducing two propositions for later applications in this paper.

Proposition 2.1. [14, *Theorem 9*] Let X be a connected symmetric graph of prime valency and G a s -regular subgroup of $\text{Aut}(X)$ for some $s \geq 1$. If a normal subgroup N of G has more than two orbits, then it is semiregular and G/N is an s -regular subgroup of $\text{Aut}(X_N)$, where X_N is the quotient graph of X corresponding to the orbits of N . Furthermore, X is a N -regular covering of X_N .

Proposition 2.2. [18] If X is an s -arc regular cubic graph, then $s \leq 5$.

Remark. If X is a regular graph with valency k on n vertices and $s \geq 1$, then there exactly $nk(k-1)^{s-1}$ s -arcs. It follows that if X is s -arc transitive then $|\text{Aut}(X)|$ must be divisible by $nk(k-1)^{s-1}$, and if X is s -regular, then $|\text{Aut}(X)| = nk(k-1)^{s-1}$. In particular, a cubic arc-transitive graph X is s -regular if and only if $|\text{Aut}(X)| = (3n)2^{s-1}$.

3. Cubic s -regular graphs of orders $22p$ and $22p^2$

In this section, we investigate the connected cubic s -regular graphs of orders $22p$ and $22p^2$, where p is a prime. We have the following lemma, by [4, 5].

Lemma 3.1. Let p be a prime. Let X be a connected cubic symmetric graph. If X has order $22p$, and $p \leq 89$, then X is isomorphic to one of the 2-regular graph $F242$ with order 242, the 3-regular graphs $F110$, $F506A$ with orders 110, 506 respectively, or the 4-regular graph $F506B$ with order 506.

Lemma 3.2. Let p be a prime. Then, there is no cubic symmetric graph of order $22p$ for $p > 89$.

Proof. Suppose that X is a connected cubic symmetric graph of the order $22p$, where $p > 89$ is a prime. Set $A := \text{Aut}(X)$. By proposition 2.2, and [18], X is at most 5-regular. Then, $|A| = 2^s \cdot 3 \cdot 11 \cdot p$, where $1 \leq s \leq 5$. Then we deduce that solvable. Because if not, then by the classification of finite simple groups, its composition factors would have to be one of the following non-abelian simple groups

$$\begin{aligned} M_{11}, M_{12}, PSL(2, 11), PSL(2, 19), PSL(2, 23), \\ PSL(2, 32), \text{ or } PSU(5, 2). \end{aligned} \tag{3.1}$$

Since $p > 89$, this contradicts the order of A . Therefore, A is solvable, and hence, elementary Abelian. Let N is a minimal normal subgroup of A . Then, N is an elementary Abelian. Hence, N is 2, 3, 11, or p -group. Then, N has more than two orbits on $V(X)$, and hence it is semiregular, by proposition 2.1. Thus, $|N| \mid 22p$. Therefore $|N| = 2, 11, \text{ or } p$. In each case, we get a contradiction.

case I): $|N| = p$

If $|N| = p$, then the quotient graph X_N of X relative to N is an A/N -symmetric graph of the order 22 , by Proposition 2.1. It is impossible by [4]. Suppose that $Q := O_p(A)$ be the maximal normal p -subgroup of A . Therefore, $O_p(A) = 1$.

case II): $|N| = 2$

If $|N| = 2$, then Proposition 2.1, implies that the quotient graph X_N corresponding to orbits of N has odd number of vertices and valency 3, which is impossible.

case III): $|N| = 11$

Now, we consider the quotient graph $X_N = X/N$ of X relative to N , where A/N is a s -regular of $Aut(X_N)$. Let K/N be a minimal normal subgroup of A/N . By the same argument as above K/N is solvable, and elementary Abelian. Then, we must have $|K/N| = 2, \text{ or } p$. Consequently $|K| = 22, \text{ or } 11p$. If $|K| = 22$, we consider the quotient graph $X_K = X/K$ of X relative to K , where A/K is a s -regular of $Aut(X_K)$. By Proposition 2.1, X_K is an A/K -symmetric graph of the order p . Then, with the same reasoning as case II, we get a contradiction. Now, suppose that $|K| = 11p$. Since $p > 89$, K has a normal subgroup of order p , which is characteristic in K and hence is normal in A , contradicting to $O_p(A) = 1$. □

Theorem 3.3. Let p be a prime. Let X be a connected cubic symmetric graph. Let p be a prime. Let X be a connected cubic symmetric graph. If X has order $22p$ then, X is isomorphic to one of the 2-regular graph $F242$ with order 242, the 3-regular graphs $F110, F506A$ with orders 110, 506 respectively, or the 4-regular graph $F506B$ with order 506.

Proof. By Lemmas 3.1 and 3.2, the proof is complete. □

Theorem 3.4. Let p be a prime. Then, there is no cubic symmetric graph of order $22p^2$.

Proof. For $p \leq 7$, by [3], there is no connected cubic symmetric graph of the order $22p^2$. Thus we may assume that $p \geq 11$. Suppose that X is a connected cubic symmetric graph of the order $22p^2$, where $p > 7$ is

a prime. Set $A := \text{Aut}(X)$. Then, $|A| = 2^s \cdot 3 \cdot 11 \cdot p^2$, where $1 \leq s \leq 5$. First, suppose that A is nonsolvable. Then, A is a product of isomorphic non-abelian simple groups. By the classification of finite simple groups, its composition factors would have to be a non-abelian simple $\{2, 3, 11\}$ -group, or $\{2, 3, 11, p\}$ -group. Let q be a prime. Then, by [13, pp. 12-14], and [6], a non-abelian simple $\{2, q, p\}$ -group is one of the groups

$$A_5, A_6, PSL(2, 7), PSL(2, 8), PSL(2, 17), PSL(3, 3), \\ PSU(3, 3), \text{ or } PSU(4, 2). \quad (3.2)$$

But, this is contradiction to the order of A . Then, composition factors is a $\{2, 3, 11, p\}$ -group. By the classification of finite simple groups, its composition factors would have to be one of the following non-abelian simple groups listed in (3.1). However, this contradicts the order of A . Therefore, A is solvable, and hence, elementary Abelian. Let N is a minimal normal subgroup of A . Then, N is an elementary Abelian. Hence, N is 2, 3, 11, or p -group. Then, N has more than two orbits on $V(X)$, and hence it is semiregular, by proposition 2.1. Thus, $|N| \mid 22p^2$. Therefore $|N| = 2, 11, p$, or p^2 . In each case, we get a contradiction.

case I): $|N| = p^2$

If $|N| = p^2$, then the quotient graph X_N of X relative to N is an A/N -symmetric graph of the order 22, by Proposition 2.1. It is impossible by [4]. Suppose that $Q := O_p(A)$ be the maximal normal p -subgroup of A .

case II): $|N| = p$

Now, we consider the quotient graph $X_N = X/N$ of X relative to N , where A/N is a s -regular of $\text{Aut}(X_N)$. Let K/N be a minimal normal subgroup of A/N . By the same argument as above K/N is solvable, and elementary Abelian. Then, we must have $|K/N| = 2, 11$, or p . Now, by considering the quotient graph X_K with the same reasoning as lemma 3.2, a contradiction can be obtained.

case III): $|N| = 11$

Now, we consider the quotient graph $X_N = X/N$ of X relative to N , where A/N is a s -regular of $\text{Aut}(X_N)$. Let K/N be a minimal normal subgroup of A/N . By the same argument as above K/N is solvable, and elementary Abelian. Then, we must have $|K/N| = 2, p$, or p^2 . Consequently $|K| = 22, 11p$, or $11p^2$. Then, with the same reasoning as case III of lemma 3.2, we arrive at a contradiction.

case IV): $|N| = 2$

In this case by the argument as in the case II of Lemma 3.2 a similar contradiction is obtained. \square

References

- [1] M. Alaeiyan and M. K. Hosseinipoor *A classification of the cubic s -regular graphs of orders $12p$ and $12p^2$* , Acta Universitatis Apulensis (2011), 153–158.
- [2] W. Bosma and J. Cannon, *Handbook of Magma Function*, Sydney University Press, Sydney, 1994.
- [3] Y. Cheng and J. Oxley, *On weakly symmetric graphs of order twice a prime*, J. Combin. Theory Ser. B 42 (1987), 196–211.
- [4] M. D. E. Conder, *Trivalent (cubic) symmetric graphs on up to 2048 vertices*, J (2006). <http://www.math.auckland.ac.nz/conder/symm-cubic2048list.txt>.
- [5] M. D. E. Conder and P. Dobcsányi, *Trivalent symmetric graphs on up to 768 vertices*, J. Combin. Math. Combin. Comput. 40 (2002) 41–63.
- [6] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, and R. A. Wilson, *Atlas of Finite Groups*, Clarendon Press, Oxford, 1985.
- [7] Y. Q. Feng and J. H. Kwak, *Classifying cubic symmetric graphs of order $10p$ or $10p^2$* , Sci. China Ser. A 49 (2006), 300–319.
- [8] Y. Q. Feng and J. H. Kwak, *Cubic symmetric graphs of order a small number times a prime or a prime square* J. Combin. Theory Ser. B 97 (2007), 627–646.
- [9] Y. Q. Feng and J. H. Kwak, *Cubic symmetric graphs of order twice an odd prime-power*, J. Aust. Math. Soc. 81 (2006), 153–164.
- [10] Y. Q. Feng and J. H. Kwak, *One-regular cubic graphs of order a small number times a prime or a prime square*, J. Aust. Math. Soc. 76 (2004), 345–356.
- [11] Y. Q. Feng, J. H. Kwak and K. Wang, *Classifying cubic symmetric graphs of order $8p$ or $8p^2$* , European J. Combin. 26 (2005), 1033–1052.
- [12] Y. Q. Feng, J. H. Kwak and M. Y. Xu, *Cubic s -regular graphs of order $2p^3$* , J. Graph Theory 52 (2006), 341–352.
- [13] D. Gorenstein, *Finite Simple Groups*, Plenum Press, New York, 1982.
- [14] P. Lorimer, *Vertex-transitive graphs: Symmetric graphs of prime valency*, J. Graph Theory 8 (1984), 55–68.
- [15] J.M. Oh, *A classification of cubic s -regular graphs of order $14p$* , Discrete Math. 309 (2009), 2721–2726.
- [16] J.M. Oh, *A classification of cubic s -regular graphs of order $16p$* , Discrete Math. 309 (2009), 3150–3155.
- [17] W. T. Tutte, *A family of cubical graphs*, Proc. Cambridge Philos. Soc. 43 (1947), 459–474.
- [18] W. T. Tutte, *On the symmetry of cubic graphs*, Canad. J. Math. 11 (1959), 621–624.

CONTACT INFORMATION

A. A. Talebi,
N. Mehdipoor Department of Mathematics, University of
Mazandaran, Babolsar, Iran
E-Mail: a.talebi@umz.ac.ir,
nargesmehdipoor@yahoo.com

Received by the editors: 07.03.2012
and in final form 14.08.2013.