

H19 AND MICRORNA-675 AS POTENTIAL BIOMARKERS FOR PROSTATE CANCER

M.M. MOHAMMED¹, M.M. RAMZY^{1*}, B.A. KAMEL¹,
G.T. MOHAMED², S.H.I. HAFIZ¹, W.F. SEDIK¹

¹ Department of Biochemistry, Faculty of Medicine, Minia University,
Egypt

² Department of Clinical Pathology, Minia Oncology Center, Egypt

E-mail: mourad61@111@gmail.com, maggiemaher24@gmail.com,
magymaher@mu.edu.eg, bothinahmed777@gmail.com,
tahagigi7@gmail.com, salwahamdy090@gmail.com,
waelsdk@yahoo.com, wael.hassanain@mu.edu.eg

Corresponding author: Ramzy M.M., e-mail: maggiemaher24@gmail.com

Prostate cancer (PCa) is considered the most common malignancy in men globally. Although prostate specific antigen (PSA) is still the only universally utilized biomarker related to prostate cancer, however PSA is not an ideal biomarker and its level can be altered by many factors. The aim of this study is to study serum H19 and miR-675 expression in prostate cancer patients and their potential diagnostic & prognostic significance targeting to find a non-invasive biomarker. This study comprised 25 PCa patients and 25 healthy controls. The expression of serum H19 and miR-675 was detected by real-time polymerase chain reaction while N-Cadherin as an adhesion marker was measured by ELISA (enzyme-linked immunosorbent assay). The expression levels of serum H19 and miR-675 were increased in prostate cancer patients compared with controls. Also serum level of N-cadherin was higher in prostate cancer patients. Upregulation of H19 and miR-675 correlated with PCa metastasis and higher Gleason score. By using the ROC curve, the sensitivity of H19 and miR-675 was higher than PSA for the diagnosis of PCa. The results suggest that H19 and miR-675 expression could be considered potential noninvasive diagnostic and prognostic biomarkers for PCa.

Key words: Prostate cancer; H19; miRNA-675; N Cadherin; Adhesion; Biomarker.

H19 ТА мікроРНК-675 ЯК ПОТЕНЦІЙНІ
БІОМАРКЕРИ РАКУ ПРОСТАТИ

© ІНСТИТУТ КЛІТИННОЇ БІОЛОГІЇ ТА ГЕНЕТИЧНОЇ
ІНЖЕНЕРІЇ НАН УКРАЇНИ, 2026

Рак простати (РП) вважається найпоширенішим злоякісним утворенням серед чоловіків по всьому світу. Хоча простат-специфічний антиген (ПСА) досі є єдиним універсально використовуваним біомаркером, пов'язаним із раком простати, він не є ідеальним, і багато факторів можуть змінювати його рівень. Мета цього дослідження полягала у вивченні експресії H19 та miR-675 у сироватці пацієнтів із раком простати та їхньої потенційної і прогностичної значущості для виявлення неінвазивного біомаркера. Учасниками цього дослідження стали 25 пацієнтів із раком простати та 25 здорових осіб для контролю. Експресію H19 та miR-675 у сироватці виявляли за допомогою ПЛР в реальному часі, а N-кадгерин як маркер адгезії вимірювали за допомогою ІФА (імуноферментного аналізу). Рівні експресії H19 та miR-675 в сироватці були вищими в пацієнтів із раком простати порівняно з контрольною групою. Крім того, рівень N-кадгерину в сироватці був вищим у пацієнтів із раком простати. Підвищена експресія H19 та miR-675 корелювала з метастазами раку простати й вищою оцінкою за шкалою Глісона. Крива ROC продемонструвала, що для діагностики раку простати чутливість H19 та miR-675 була вищою, ніж ПСА. Згідно з отриманими результатами експресію H19 та miR-675 можна вважати потенційним неінвазивним діагностичним та прогностичним біомаркером раку простати.

Ключові слова: рак простати, H19, мікроРНК-675, N-кадгерин, адгезія, біомаркер.

REFERENCES

- Al Shareef, Z., Al-Shahrabi, R., Saheb Sharif-Askari, F., AlDhmanie, A., Alshamsi, Y., Zarooni, A.A., and Bendardaf, R., Incidence and risk factors of prostate cancer among the Northern and Eastern parts of the United Arab Emirates population, *The Prostate*, 2024, vol. 84, no. 2, pp. 185–192. <https://doi.org/10.1002/pros.24637>
- Alhamzo, J.S.J., and Abdulazeez, A.M., Prostate cancer: MRI image detection based on deep learning – a review, *Indones. J. Computer Sci.*, 2024, vol. 13, no. 3. <https://doi.org/10.33022/ijcs.v13i3.4045>
- Azzalini, E., and Bonin, S., Molecular diagnostics of prostate cancer: impact of molecular tests, *Asian. J. Androl.*, 2024, vol. 26, no.6, pp. 562–566. <https://doi.org/10.4103/aja202411>

- Bacci, L., Aiello, A., Ripoli, C., Loria, R., Pugliese, D., Pierconti, F., Rotili, D., Strigari, L., Pinto, F., Bassi, P.F., et al., H19-dependent transcriptional regulation of $\beta 3$ and $\beta 4$ integrins upon estrogen and hypoxia favors metastatic potential in prostate cancer, *Int. J. Mol. Sci.*, 2019, vol. 20, no. 16, pp. 4012. <https://doi.org/10.3390/ijms20164012>
- Bozgeyik, E., and Bozgeyik, I., Non-coding RNA variations in oral cancers: a comprehensive review, *Gene*, 2023, vol. 851, pp. 147012. <https://doi.org/10.1016/j.gene.2022.147012>
- Cai, X., and Cullen, B.R., The imprinted H19 noncoding RNA is a primary microRNA precursor, *RNA*, 2007, vol. 13, no. 3, pp. 313–316. <https://doi.org/10.1261/rna.351707>
- Chen, X., Li, Y., Zuo, C., Zhang, K., Lei, X., Wang, J., Yang, Y., Zhang, J., Ma, K., Wang, S., et al., Long non-coding RNA H19 regulates glioma cell growth and metastasis via miR-200a-mediated CDK6 and ZEB1 expression, *Front. Oncol.*, 2021, vol. 11, pp. 757650. <https://doi.org/10.3389/fonc.2021.757650>
- Conti, D.V., Darst, B.F., Moss, L.C., Saunders, E.J., Sheng, X., Chou, A., Strom, S.S., et al., Trans-ancestry genome-wide association meta-analysis of prostate cancer identifies new susceptibility loci and informs genetic risk prediction, *Nat. Genet.*, 2021, vol. 53, no. 1, pp. 65–75. <https://doi.org/10.1038/s41588-020-00748-0>
- Darmadi, D., Chugaeva, U.Y., Saleh, R.O., Hjaz, A., Saleem, H.M., Ghildiyal, P., Alwaily, E.R., Alawadi, A., Alnajjar, M.J., and Ihsan, A., Critical roles of long noncoding RNA H19 in cancer, *Cell Biochem. Func.*, 2024, vol. 42, no. 3, pp. e4018. <https://doi.org/10.1002/cbf.4018>
- Debnath, P., Huiem, R.S., Dutta, P., and Palchaudhuri, S., Epithelial-mesenchymal transition and its transcription factors, *Biosci. Rep.*, 2022, vol. 42, no. 1, pp. BSR20211754. <https://doi.org/10.1042/BSR20211754>
- Dey, B.K., Pfeifer, K., and Dutta, A., The H19 long noncoding RNA gives rise to microRNAs miR-675-3p and miR-675-5p to promote skeletal muscle differentiation and regeneration, *Genes Dev.*, 2014, vol. 28, no. 5, pp. 491–501. <https://doi.org/10.1101/gad.234419.113>
- Drivalos, A., Chrisofos, M., Efsthathiou, E., Kapranou, A., Kollaitis, G., Koutlis, G., Antoniou, N., Karanastasis, D., Dimopoulos, M.A., and Bamias, A., Expression of $\alpha 5$ -integrin, $\alpha 7$ -integrin, E-cadherin, and N-cadherin in localized prostate cancer, *Urologic Oncology: Seminars and Original Investigations*, 2016, vol. 34, no. 4, pp. 165.e11–165.e18. <https://doi.org/10.1016/j.urolonc.2015.11.016>
- Fares, J., Fares, M.Y., Khachfe, H.H., Salhab, H.A., and Fares, Y., Molecular principles of metastasis: a hallmark of cancer revisited, *Signal Transduction and Targeted Therapy*, 2020, vol. 5, no. 1, pp. 28. <https://doi.org/10.1038/s41392-020-0134-x>
- Gao, L.M., Xu, S.F., Zheng, Y., Wang, P., Zhang, L., Shi, S.S., Wu, T., Li, Y., Zhao, J., Tian, Q., et al., Long non-coding RNA H19 is responsible for the progression of lung adenocarcinoma by mediating methylation-dependent repression of CDH1 promoter, *J. Cell. Mol. Med.*, 2019, vol. 23, no. 9, pp. 6411–6428. <https://doi.org/10.1111/jcmm.14533>
- Graça, I., Pereira-Silva, E., Henrique, R., Packham, G., Crabb, S.J., and Jernimo, C., Epigenetic modulators as therapeutic targets in prostate cancer, *Clin. Epigenet.*, 2016, vol. 8, no. 1, pp. 98. <https://doi.org/10.1186/s13148-016-0264-8>
- Herman, A.B., Tsitsipatis, D., and Gorospe, M., Integrated lncRNA function upon genomic and epigenomic regulation, *Mol. Cell*, 2022, vol. 82, no. 12, pp. 2252–2266. <https://doi.org/10.1016/j.molcel.2022.05.027>
- John, J., Adam, A., Kaestner, L., Spies, P., Mutambirwa, S., Lazarus, J., et al., The South African prostate cancer screening guidelines, *South Afric. Med. J.*, 2024, vol. 114, no. 5, pp. 15–18. https://hdl.handle.net/10520/ejc-m_samj_v114_n5_a7
- Karamali, N., Daraei, A., Rostamlou, A., Mahdavi, R., Akbari Jonoush, Z., Ghadiri, N., Mahmoudi, Z., Mardi, A., Javidan, M., Sohrabi, S.S., and Baradaran, B., Decoding contextual crosstalk: revealing distinct interactions between non-coding RNAs and unfolded protein response in breast cancer, *Cancer Cell Int.*, 2024, vol. 24, no. 1, pp. 104. <https://doi.org/10.1186/s12935-024-03296-3>
- Kratzer, T.B., Mazzitelli, N., Star, J., Dahut, W.L., Jemal, A., and Siegel, R.L., Prostate cancer statistics, *CA: A Cancer J. Clinicians*, 2025. <https://doi.org/10.3322/caac.70028>
- Lecerf, C., Le Bourhis, X., and Adriaenssens, E., The long non-coding RNA H19: an active player with multiple facets to sustain the hallmarks of cancer, *Cell. Mol. Life Sci.*, 2019, vol. 76, pp. 4673–4687. <https://doi.org/10.1007/s00018-019-03240-z>
- Lee, K.M., Nelson, T.J., Bryant, A., Teerlink, C.C., Gulati, R., Pagadala, M.S., and Lynch, J.A., Genetic risk and likelihood of prostate cancer detection on first biopsy by ancestry, *JNCI*, 2024, vol. 116, no. 5, pp. 753–757. <https://doi.org/10.1093/jnci/djae002>
- Li, J., Zhang, G., Liu, C.G., Xiang, X., Le, M.T., Sethi, G., Wang, L., Goh, B.C., and Ma, Z., The potential role of exosomal circRNAs in the tumor microenvironment: insights into cancer diagnosis and therapy, *Theranostics*, 2022, vol. 12, no. 1, pp. 87–104. <https://doi.org/10.7150/thno.64096>

- Liao, J., Chen, B., Zhu, Z., Du, C., Gao, S., Zhao, G., Zhao, P., Wang, Y., Wang, A., Schwartz, Z., et al., Long noncoding RNA (lncRNA) H19: an essential developmental regulator with expanding roles in cancer, stem cell differentiation, and metabolic diseases, *Genes Dis.*, 2023, vol. 10, no. 4, pp. 1351–1366. <https://doi.org/10.1016/j.gendis.2023.02.008>
- Liao, S., Yu, C., Liu, H., Zhang, C., Li, Y., and Zhong, X., Long non-coding RNA H19 promotes the proliferation and invasion of lung cancer cells and regulates the expression of E-cadherin, N-cadherin, and vimentin, *Oncotargets and Therapy*, 2019, vol. 12, pp. 4099–4107. <https://doi.org/10.2147/OTT.S185156>
- Liu, C., Chen, Z., Fang, J., Xu, A., Zhang, W., and Wang, Z., H19-derived miR-675 contributes to bladder cancer cell proliferation by regulating p53 activation, *Tumor Biol.*, 2016, vol. 37, pp. 263–270. <https://doi.org/10.1007/s13277-015-3779-2>
- Mahal, B.A., Aizer, A.A., Efsthathiou, J.A., and Nguyen, P.L., Association of very low prostate-specific antigen levels with increased cancer-specific death in men with high-grade prostate cancer, *Cancer*, 2016, vol. 122, no. 1, pp. 78–83. <https://doi.org/10.1002/cncr.29691>
- McNeal, J.E., Redwine, E.A., Freiha, F.S., and Stamey, T.A., Zonal distribution of prostatic adenocarcinoma: correlation with histologic pattern and direction of spread, *Amer. J. Surgical Pathol.*, 1988, vol. 12, no. 12, pp. 897–906
- Montanari, M., Rossetti, S., Cavaliere, C., D’Aniello, C., Malzone, M.G., Vanacore, D., Franco, R.D., Mantia, E.L., Iovane, G., Piscitelli, R., et al., Epithelial–mesenchymal transition in prostate cancer: an overview, *Oncotarget*, 2017, vol. 8, no. 21, pp. 35376–35389. <https://doi.org/10.18632/oncotarget.15686>
- Moreira-Silva, F., Henrique, R., and Jernimo, C., From therapy resistance to targeted therapies in prostate cancer, *Frontiers in Oncology*, 2022, vol. 12, pp. 877379. <https://doi.org/10.3389/fonc.2022.877379>
- Nalla, A.K., Estes, N., Patel, J., and Rao, J.S., N-cadherin mediates angiogenesis by regulating monocyte chemoattractant protein-1 expression via PI3K/Akt signaling in prostate cancer cells, *Exper. Cell Res.*, 2011, vol. 317, no. 17, pp. 2512–2521. <https://doi.org/10.1016/j.yexcr.2011.07.024>
- Nokkeaw, A., Thamjamrassri, P., Chantaravisoot, N., Tangkijvanich, P., and Ariyachet, C., Long non-coding RNA H19 promotes proliferation in hepatocellular carcinoma cells via H19/miR-107/CDK6 axis, *Oncol. Res.*, 2023, vol. 31, no. 6, pp. 989–1005. <https://doi.org/10.32604/or.2023.030395>
- Peperstraete, E., Lecerf, C., Collette, J., Vennin, C., Raby, L., Völkel, P., Angrand, P.O., Winter, M., Bertucci, F., Finetti, P., et al., Enhancement of breast cancer cell aggressiveness by lncRNA H19 and its miR-675 derivative: insight into shared and different actions, *Cancers*, 2020, vol. 12, no. 7, pp. 1730. <https://doi.org/10.3390/cancers12071730>
- Pope, C., Piekos, S.C., Chen, L., Mishra, S., and Zhong, X.B., The role of H19, a long non-coding RNA, in mouse liver postnatal maturation, *PLoS One*, 2017, vol. 12, no. 11, p. e0187557. <https://doi.org/10.1371/journal.pone.0187557>
- Rivera-Izquierdo, M., Perez de Rojas, J., Martínez-Ruiz, V., Pérez-Gymez, B., Sánchez, M.J., Khan, K.S., and Jiménez-Moleyn, J.J., Obesity as a risk factor for prostate cancer mortality: a systematic review and dose-response meta-analysis of 280,199 patients, *Cancers*, 2021, vol. 13, no. 16, pp. 4169. <https://doi.org/10.3390/cancers13164169>
- Roche, J., The epithelial-to-mesenchymal transition in cancer, *Cancers*, 2018, vol. 10, no. 2, pp. 52. <https://doi.org/10.3390/cancers10020052>
- Shaker, O.G., Safa, A., Khairy, A., and Abozeid, N.F., Serum long non-coding RNA H19/micro RNA-675-5p axis as a probable diagnostic biomarker in inflammatory bowel disease, *Mol. Biol. Rep.*, 2023, vol. 50, no. 11, pp. 9029–9036. <https://doi.org/10.1007/s11033-023-08777-8>
- Singh, N., Ramnarine, V.R., Song, J.H., Pandey, R., Padi, S.K., Nouri, M., Olive, V., Kobelev, M., Okumura, K., McCarthy, D., et al., The long noncoding RNA H19 regulates tumor plasticity in neuroendocrine prostate cancer, *Nat. Commun.*, 2021, vol. 12, p. 7349. <https://doi.org/10.1038/s41467-021-26901-9>
- Steenbuck, I.D., Cosenza-Contreras, M., Froehlich, K., Mayer, B., Kurowski, K., Werner, T., Reinold, M., Fahrner, M., Hause, F., Seredynska, A., et al., Serum proteome profiling identifies N-cadherin and C-Met as early marker candidates of therapeutic response to neoadjuvant chemotherapy in breast cancer, *bioRxiv*, 2024. <https://doi.org/10.1101/2024.05.24.595719>
- Tao, S., Wang, J., Li, F., Shi, B., Ren, Q., Zhuang, Y., and Qian, X., Extracellular vesicles released by hypoxia-induced tumor-associated fibroblasts impart chemoresistance to breast cancer cells via long noncoding RNA H19 delivery, *FASEB J.*, 2024, vol. 38, no. 2, p. e23165. <https://doi.org/10.1096/fj.202300203R>
- Thakur, K.K., Kumar, A., Banik, K., Verma, E., Khattoon, E., Harsha, C., and Kunnumakkara, A.B., Long noncoding RNAs in triple-negative breast cancer: a new frontier in the regulation of tumorigenesis, *J. Cell. Physiol.*, 2021, vol. 236, no. 12, pp. 7938–7965. <https://doi.org/10.1002/jcp.30463>
- Tiwari, N., Gheldof, A., Tatari, M., and Christofori, G.,

- EMT as the ultimate survival mechanism of cancer cells, *Semin. Cancer Biol.*, 2012, vol. 22, no. 3, pp. 194–207. <https://doi.org/10.1016/j.semcancer.2012.02.013>
- VanGuilder, H.D., Vrana, K.E., and Freeman, W.M., Twenty-five years of quantitative PCR for gene expression analysis, *Biotechniques*, 2008, vol. 44, no. 5, pp. 619–626. <https://doi.org/10.2144/000112776>
- Vennin, C., Spruyt, N., Dahmani, F., Julien, S., Bertucci, F., Finetti, P., Thierry, C., Bourette, P.R., Bourhis, X.L., and Adriaenssens, E., H19 non-coding RNA-derived miR-675 enhances tumorigenesis and metastasis of breast cancer cells by downregulating c-Cbl and Cbl-b, *Oncotarget.*, 2015, vol. 6, no. 30, pp. 29209–29223. <https://doi.org/10.18632/oncotarget.4976>
- Vergara, D., Simeone, P., Franck, J., Trerotola, M., Giudetti, A., Capobianco, L., Tinelli, A., Bellomo, C., Fournier, I., Gaballo, A., et al., Translating epithelial mesenchymal transition markers into the clinic: novel insights from proteomics, *EuPA Open Proteomics*, 2016, vol. 10, pp. 31–41. <https://doi.org/10.1016/j.euprot.2016.01.003>
- Wang, G., Zhao, D., Spring, D.J., and DePinho, R.A., Genetics and biology of prostate cancer, *Genes and Development*, 2018, vol. 32, no. 17–18, pp. 1105–1140. <https://doi.org/10.1101/gad.315739.118>
- Yan, J., Zhang, Y., She, Q., Li, X., Peng, L., Wang, X., Liu, S., Shen, X., Zhang, W., Dong, Y., et al., Long noncoding RNA H19/miR-675 axis promotes gastric cancer via FADD/Caspase 8/Caspase 3 signaling pathway, *Cell. Physiol. Biochem.*, 2017, vol. 42, no. 6, pp. 2364–2376. <https://doi.org/10.1159/000480028>
- Zhang, W., Zhou, K., Zhang, X., Wu, C., Deng, D., and Yao, Z., Roles of the H19/microRNA-675 axis in the proliferation and epithelial-mesenchymal transition of human cutaneous squamous cell carcinoma cells, *Oncology Reports*, 2021, vol. 45, no. 4, p. 39. <https://doi.org/10.3892/or.2021.7990>
- Zhu, M., Chen, Q., Liu, X., Sun, Q., Zhao, X., Deng, R., Wang, Y., Huang, J., Xu, M., Yan, J., and Yu, J., lncRNA H19/miR-675 axis represses prostate cancer metastasis by targeting TGFBI, *FEBS J.*, 2014, vol. 281, no. 16, pp. 3766–3775. <https://doi.org/10.1111/febs.12902>

Received May 24, 2025

Received July 23, 2025

Accepted March 18, 2026