

J. O. Hamzat¹ (Univ. Lagos, Nigeria),

A. A. El-Ashwah (Faculty of Science Damietta Univ., New Damietta, Egypt)

SOME PROPERTIES OF A GENERALIZED MULTIPLIER TRANSFORM ON ANALYTIC p -VALENT FUNCTIONS

ДЕЯКІ ВЛАСТИВОСТІ УЗАГАЛЬНЕНОГО МУЛЬТИПЛІКАТИВНОГО ПЕРЕТВОРЕННЯ НА АНАЛІТИЧНИХ p -ВАЛЕНТНИХ ФУНКЦІЯХ

For a function

$$f(z) = z^p + \sum_{k=1}^{\infty} a_{k+p} z^{k+p}$$

where $p \in \mathbb{N}$, the authors investigate some properties of a more general multiplier transform on analytic p -valent functions in an open unit disk. The applications of the obtained results to fractional calculus are pointed out, while several other corollaries follow as simple consequences.

Для функції

$$f(z) = z^p + \sum_{k=1}^{\infty} a_{k+p} z^{k+p},$$

де $p \in \mathbb{N}$, досліджено деякі властивості більш загального мультиплікативного перетворення на аналітичних p -валентних функціях у відкритому одиничному колі. Розглянуто застосування отриманих результатів до дробового числення, а деякі інші результати отримано як прості наслідки.

1. Introduction and preliminaries. Let Γ denote the class of analytic functions $f(z)$, having the series representation

$$f(z) = z + \sum_{k=2}^{\infty} a_k z^k \quad (1)$$

and normalized by $f'(0) - 1 = 0 = f(0)$ in the open unit disk $U = \{z \in \mathbb{C} : |z| < 1\}$.

Also let Γ_p denote the class of analytic p -valent functions $f(z)$ having the form

$$f(z) = z^p + \sum_{k=1}^{\infty} a_{k+p} z^{k+p}, \quad p \in \mathbb{N}. \quad (2)$$

A function $U = u(x, y)$ is said to be harmonic if it is a real-valued function having continuous partial derivatives of order one and two, and satisfying

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0.$$

However, a continuous complex-valued function $f(z) = u(x, y) + iv(x, y)$ is said to be harmonic in a complex domain D say, if both the real and imaginary parts $u(x, y)$ and $v(x, y)$, respectively, are harmonic in D . The geometric function theory is mostly interested in the survey of properties of analytic functions (see [2, 5]). Given any simply connected region $R \subset D$, we can say that

¹ Corresponding author, e-mail: jhamzat@unilag.edu.ng.

$$f(z) = h(z) + \overline{g(z)},$$

where h and g are analytic in the connected region R . Conventionally, we refer to h and g as the analytic and co-analytic parts, respectively. Thus a necessary and sufficient condition for function f to be locally univalent and orientation preserving is that

$$|h'(z)| > |g'(z)| \in R,$$

see [1, 4] among others. Let \mathcal{H} denote the family of p -valent harmonic function in U . Then h and g can be expressed as

$$h(z) = z^p + \sum_{k=1}^{\infty} a_{k+p} z^{k+p}, \quad g(z) = \sum_{k=0}^{\infty} b_{k+p} z^{k+p}$$

for $p \in \mathbb{N}$ and, in particular, $0 \leq |b_p| < 1$.

Therefore, we write that

$$f(z) = z^p + \sum_{k=1}^{\infty} a_{k+p} z^{k+p} + \overline{\sum_{k=0}^{\infty} b_{k+p} z^{k+p}}. \tag{3}$$

As a special case, if the co-analytic part of f is identically zero (i.e., $g = 0$), then the family of orientation preserving, normalized harmonic univalent functions reduces to the usual class of normalized analytic functions. For function $f(z)$ of the form (1), Swamy [10], in 2012 introduced and studied a multiplier differential operator $I_{\alpha,\beta}^n f(z)$ given by

$$I_{\alpha,\beta}^n f(z) = z + \sum_{k=2}^{\infty} \left(\frac{\alpha + k\beta}{\alpha + \beta} \right)^n a_k z^k,$$

see also [8].

Furthermore, we define for function $f(z)$ of the form (2), a linear differential operator $L_{\alpha,\beta,\gamma}^{n,p} f(z)$ such that

$$\begin{aligned} L_{\alpha,\beta,\gamma}^{1,p} f(z) &= \frac{\alpha f(z) + \beta z f'(z) + \gamma z(z f'(z))'}{\alpha + \beta p + \gamma p^2}, \\ L_{\alpha,\beta,\gamma}^{2,p} f(z) &= L_{\alpha,\beta,\gamma}^{1,p} f(z) \left(L_{\alpha,\beta,\gamma}^{1,p} f(z) \right), \\ L_{\alpha,\beta,\gamma}^{3,p} f(z) &= L_{\alpha,\beta,\gamma}^{1,p} f(z) \left(L_{\alpha,\beta,\gamma}^{2,p} f(z) \right), \\ &\dots\dots\dots \\ L_{\alpha,\beta,\gamma}^{n,p} f(z) &= L_{\alpha,\beta,\gamma}^{1,p} f(z) \left(L_{\alpha,\beta,\gamma}^{n-1,p} f(z) \right), \end{aligned} \tag{4}$$

where $p \in \mathbb{N}$, $n, \alpha, \beta \geq 0$ and α is real such that $\alpha + \beta + \gamma > 0$. It follows from (4) that

$$L_{\alpha,\beta,\gamma}^{n,p} f(z) = z^p + \sum_{k=1}^{\infty} \left(\frac{\alpha + \beta(k+p) + \gamma(k+p)^2}{\alpha + \beta p + \gamma p^2} \right)^n a_{k+p} z^{k+p}. \tag{5}$$

Remark. Suppose that the function $f(z)$ has the form (2), it is easily verified from (5) that

$$L_{\alpha,0,0}^{0,p}f(z) = f(z) \in \Gamma_p \quad \text{and} \quad L_{\alpha,0,0}^{0,1}f(z) = f(z) \in \Gamma_p.$$

It is obvious that the operator $L_{\alpha,\beta,\gamma}^{n,p}f(z)$ generalizes many existing operators of this kind which were introduced and studied by different authors. For instance,

- (i) $L_{\alpha,\beta,\gamma}^{n,1}f(z) = I_{\alpha,\beta,\gamma}^n f(z)$ studied by Makinde et al. [8];
- (ii) $L_{\alpha,\beta,0}^{n,1}f(z) = I_{\alpha,\beta}^n f(z)$ studied by Swamy [10];
- (iii) $L_{\alpha,1,\gamma}^{n,1}f(z) = I_{\alpha}^n f(z)$, $\alpha > -1$ studied by Cho and Srivastava [3];
- (iv) $L_{1,\beta,0}^{n,1}f(z) = N_{\beta}^n f(z)$ studied by Swamy [10].

With reference to (5), we can write that

$$L_{\alpha,\beta,\gamma}^{n,p}f(z) = H_{\alpha,\beta,\gamma}^{n,p}f(z) + \overline{G_{\alpha,\beta,\gamma}^{n,p}f(z)}. \quad (6)$$

Now using (6), we give the following definition.

Definition 1. Let $f(z)$ be of the form (3), then $f(z) \in \mathcal{H}_{\mu}^{n,p}(\alpha, \beta, \gamma)$ if it satisfies the condition that

$$\Re \left\{ \frac{z \left(H_{\alpha,\beta,\gamma}^{n,p}f(z) \right)' - \overline{z \left(G_{\alpha,\beta,\gamma}^{n,p}f(z) \right)'}}{H_{\alpha,\beta,\gamma}^{n,p}f(z) + \overline{G_{\alpha,\beta,\gamma}^{n,p}f(z)}} \right\} \geq \mu \quad (7)$$

for $p \in \mathbb{N}$, $0 \leq \mu < p$, $n, \beta, \gamma \geq 0$ and α is real such that $\alpha + \beta + \gamma > 0$.

In addition, suppose that

$$\mathcal{V}_{\mathcal{H}}^{n,p}(\alpha, \beta, \gamma, \mu) = \mathcal{V}_{\mathcal{H}}^{n,p} \cap \mathcal{H}_{\mu}^{n,p}(\alpha, \beta, \gamma), \quad (8)$$

where $\mathcal{V}_{\mathcal{H}}^{n,p}$ is the harmonic functions with varying arguments consists of functions f of the form (3) in $\mathcal{H}_{\mu}^{n,p}$ for which there exists a real number σ such that

$$\psi_{k+p} + k\sigma \equiv \pi \pmod{2\pi}, \quad \tau_{k+p} + k\sigma \equiv 0 \pmod{2\pi}, \quad k \geq 1, \quad (9)$$

where

$$\psi_{k+p} = \arg(a_{k+p}) \quad \text{and} \quad \tau_{k+p} = \arg(b_{k+p}).$$

At this juncture, we shall obtain a sufficient coefficient condition for function f of the form (3) to be in the aforementioned class $\mathcal{H}_{\mu}^{n,p}(\alpha, \beta, \gamma)$. It is noted that this coefficient condition is also necessary for functions belonging to the class $\mathcal{V}_{\mathcal{H}}^{n,p}(\alpha, \beta, \gamma)$.

2. Necessary and sufficient coefficient for the class $\mathcal{H}_{\mu}^{n,p}(\alpha, \beta, \gamma)$.

Theorem 2.1. Let $f(z)$ be of the form (3). Then, for $p \in \mathbb{N}$, $0 \leq \mu < p$, $\alpha > 0$, $\beta, \gamma \geq 0$, $\alpha + \beta + \gamma > 0$, and $|b_p| < \frac{p-\mu}{p+\mu}$, $f \in \mathcal{H}_{\mu}^{n,p}(\alpha, \beta, \gamma)$ if

$$\sum_{k=1}^{\infty} \left[\frac{k+p-\mu}{p-\mu} |a_{k+p}| + \frac{k+p+\mu}{p-\mu} |b_{k+p}| \right] \times \left(\frac{\alpha + \beta(k+p) + \gamma(k+p)^2}{\alpha + \beta p + \gamma p^2} \right)^n \leq 1 - \frac{p+\mu}{p-\mu} |b_p|. \quad (10)$$

Proof. We begin the prove by showing that the condition (7) is satisfied if the inequality (10) holds true for the coefficient of f defined in (3). Using (6) and (7), we have

$$\omega(z) = \left\{ \frac{z \left(H_{\alpha, \beta, \gamma}^{n, p} f(z) \right)' - \overline{z \left(G_{\alpha, \beta, \gamma}^{n, p} f(z) \right)'}}{p \left(H_{\alpha, \beta, \gamma}^{n, p} f(z) + \overline{G_{\alpha, \beta, \gamma}^{n, p} f(z)} \right)} \right\} = \frac{M(z)}{N(z)}, \quad (11)$$

where

$$M(z) = \frac{z}{p} \left(H_{\alpha, \beta, \gamma}^{n, p} f(z) \right)' - \frac{\overline{z}}{p} \left(G_{\alpha, \beta, \gamma}^{n, p} f(z) \right)'$$

and

$$N(z) = H_{\alpha, \beta, \gamma}^{n, p} f(z) + \overline{G_{\alpha, \beta, \gamma}^{n, p} f(z)}.$$

Here, we recall that $\Re(\omega) > \frac{\mu}{p}$ if and only if

$$|p - \mu + p\omega| \geq |p + \mu - p\omega|.$$

Then, from (11), it suffices to show that

$$|M(z) + (p - \mu)N(z)| \geq |M(z) - (p + \mu)N(z)|$$

and

$$|M(z) + (p - \mu)N(z)| - |M(z) - (p + \mu)N(z)| \geq 0. \quad (12)$$

Having substituted for the values of $M(z)$ and $N(z)$ in (12), we obtain

$$\begin{aligned} & |M(z) + (p - \mu)N(z)| - |M(z) - (p + \mu)N(z)| \geq \\ & \geq 2(p - \mu)|z|^p - \sum_{k=1}^{\infty} 2(k + p) \left(\frac{\alpha + \beta(k + p) + \gamma(k + p)^2}{\alpha + \beta p + \gamma p^2} \right)^n |a_{k+p}| |z|^{k+p} - \\ & - \sum_{k=0}^{\infty} 2(k + p) \left(\frac{\alpha + \beta(k + p) + \gamma(k + p)^2}{\alpha + \beta p + \gamma p^2} \right)^n |b_{k+p}| |z|^{k+p} + \\ & + 2\mu \sum_{k=1}^{\infty} \left(\frac{\alpha + \beta(k + p) + \gamma(k + p)^2}{\alpha + \beta p + \gamma p^2} \right)^n |a_{k+p}| |z|^{k+p} - \\ & - 2\mu \sum_{k=0}^{\infty} \left(\frac{\alpha + \beta(k + p) + \gamma(k + p)^2}{\alpha + \beta p + \gamma p^2} \right)^n |b_{k+p}| |z|^{k+p}, \end{aligned}$$

that is,

$$\begin{aligned} & |M(z) + (p - \mu)N(z)| - |M(z) - (p + \mu)N(z)| \geq \\ & \geq 2(p - \mu)|z|^p - \sum_{k=1}^{\infty} 2(k + p - \mu)Y^n |a_{k+p}| |z|^{k+p} - \sum_{k=0}^{\infty} 2(k + p + \mu)Y^n |b_{k+p}| |z|^{k+p} \geq \end{aligned}$$

$$\begin{aligned} &\geq 2(p - \mu)|z|^p \left\{ 1 - \sum_{k=1}^{\infty} \frac{k + p - \mu}{p - \mu} Y^n |a_{k+p}| - \sum_{k=0}^{\infty} \frac{k + p + \mu}{p - \mu} Y^n |b_{k+p}| \right\} \geq \\ &\geq 2(p - \mu)|z|^p \left\{ 1 - \frac{p + \mu}{p - \mu} |b_p| - \sum_{k=1}^{\infty} \left[\frac{k + p - \mu}{p - \mu} |a_{k+p}| - \frac{k + p + \mu}{p - \mu} |b_{k+p}| \right] Y^n \right\} \geq 0 \end{aligned}$$

by virtue of inequality (10) where $Y = \frac{\alpha + \beta(k + p) + \gamma(k + p)^2}{\alpha + \beta p + \gamma p^2}$.

This shows that $f \in \mathcal{H}_\mu^{n,p}(\alpha, \beta, \gamma)$.

Theorem 2.1 is proved.

Corollary 2.1. Let $f(z) \in \mathcal{H}_\mu^{n,1}(\alpha, \beta, \gamma)$. Then, for $0 \leq \mu < 1$, $\alpha > 0$, $\beta, \gamma \geq 0$ and $\alpha + \beta + \gamma > 0$,

$$\sum_{k=2}^{\infty} \left[\frac{k - \mu}{1 - \mu} |a_k| + \frac{k + \mu}{1 - \mu} |b_k| \right] \left(\frac{\alpha + \beta k + \gamma k^2}{\alpha + \beta + \gamma} \right)^n \leq 1 - \frac{1 + \mu}{1 - \mu} |b_1|. \tag{13}$$

Corollary 2.2. Let $f(z) \in \mathcal{H}_0^{n,1}(\alpha, \beta, \gamma)$. Then, for $\alpha > 0$, $\beta, \gamma \geq 0$ and $\alpha + \beta + \gamma > 0$,

$$\sum_{k=2}^{\infty} [k|a_k| + k|b_k|] \left(\frac{\alpha + \beta k + \gamma k^2}{\alpha + \beta + \gamma} \right)^n \leq 1 - |b_1|.$$

Corollary 2.3. Let $f(z) \in \mathcal{H}_0^{0,1}(\alpha, \beta, \gamma)$. Then, for $\alpha > 0$, $\beta, \gamma \geq 0$ and $\alpha + \beta + \gamma > 0$,

$$\sum_{k=2}^{\infty} [k|a_k| + k|b_k|] \leq 1 - |b_1|. \tag{14}$$

Next we obtain both the necessary and sufficient condition for function f of the form (3) given the condition (8).

Theorem 2.2. $f \in \mathcal{V}_{\mathcal{H}}^{n,p}(\alpha, \beta, \gamma, \mu)$ if and only if

$$\sum_{k=1}^{\infty} \left[\frac{k + p - \mu}{p - \mu} |a_{k+p}| + \frac{k + p - 2\mu}{p - \mu} |b_{k+p}| \right] \left(\frac{\alpha + \beta(k + p) + \gamma(k + p)^2}{\alpha + \beta p + \gamma p^2} \right)^n \leq 1 - \frac{p + \mu}{p - \mu} |b_p| \tag{15}$$

for $p \in \mathbb{N}$, $0 \leq \mu < p$, $\alpha > 0$, $\beta, \gamma \geq 0$ and $\alpha + \beta + \gamma > 0$.

Proof. Since $\mathcal{V}_{\mathcal{H}}^{n,p}(\alpha, \beta, \gamma, \mu) \subset \mathcal{H}_\mu^{n,p}(\alpha, \beta, \gamma)$. Then the necessary condition part of the theorem shall be established. Suppose that $f \in \mathcal{V}_{\mathcal{H}}^{n,p}(\alpha, \beta, \gamma, \mu)$, then appealing to (6) and (7), we have that

$$\Re \left\{ \left[\frac{z \left(H_{\alpha, \beta, \gamma}^{n,p} f(z) \right)' - \overline{z \left(G_{\alpha, \beta, \gamma}^{n,p} f(z) \right)'}}{H_{\alpha, \beta, \gamma}^{n,p} f(z) + \overline{G_{\alpha, \beta, \gamma}^{n,p} f(z)}} \right] - \mu \right\} \geq 0.$$

Equivalently, we can write that

$$\Re \left\{ \frac{pz^p + \sum_{k=1}^{\infty} (k + p) Y^n a_{k+p} z^{k+p} - \sum_{k=0}^{\infty} (k + p) Y^n b_{k+p} \overline{z^{k+p}}}{z^p + \sum_{k=1}^{\infty} Y^n a_{k+p} z^{k+p} + \sum_{k=0}^{\infty} Y^n b_{k+p} \overline{z^{k+p}}} - \mu \right\} \geq$$

$$\begin{aligned} &\geq \left\{ \frac{[(p - \mu) - (p + \mu)|b_p|]}{(1 + |b_p|) + \sum_{k=1}^{\infty} Y^n |a_{k+p}| |z^k| + \left| \frac{\bar{z}}{z} \right|^p \sum_{k=1}^{\infty} Y^n |b_{k+p}| |\bar{z}^k|} \right\} - \\ &- \left\{ \frac{-\sum_{k=1}^{\infty} (k + p - \mu) Y^n |a_{k+p}| |z^k|}{(1 + |b_p|) + \sum_{k=1}^{\infty} Y^n |a_{k+p}| |z^k| + \left| \frac{\bar{z}}{z} \right|^p \sum_{k=1}^{\infty} Y^n |b_{k+p}| |\bar{z}^k|} \right\} - \\ &- \left\{ \frac{-\left| \frac{\bar{z}}{z} \right|^p \sum_{k=1}^{\infty} (k + p + \mu) Y^n |b_{k+p}| |\bar{z}^k|}{(1 + |b_p|) + \sum_{k=1}^{\infty} Y^n |a_{k+p}| |z^k| + \left| \frac{\bar{z}}{z} \right|^p \sum_{k=1}^{\infty} Y^n |b_{k+p}| |\bar{z}^k|} \right\} \geq 0 \end{aligned}$$

and Y is as earlier defined.

The above condition must hold for all the values of z such that $|z| = r < 1$. With σ as in (9), we obtain

$$\begin{aligned} &\frac{[(p - \mu) - (p + \mu)|b_p|] - \sum_{k=1}^{\infty} [(k + p - \mu)|a_{k+p}| - (k + p + \mu)|b_{k+p}|]}{(1 + |b_p|) + \sum_{k=1}^{\infty} [|a_{k+p}| + |b_{k+p}|] \left(\frac{\alpha + \beta(k + p) + \gamma(k + p)^2}{\alpha + \beta p + \gamma p^2} \right)^n r^k} \times \\ &\times \frac{\left(\frac{\alpha + \beta(k + p) + \gamma(k + p)^2}{\alpha + \beta p + \gamma p^2} \right)^n r^k}{(1 + |b_p|) + \sum_{k=1}^{\infty} [|a_{k+p}| + |b_{k+p}|] \left(\frac{\alpha + \beta(k + p) + \gamma(k + p)^2}{\alpha + \beta p + \gamma p^2} \right)^n r^k} \geq 0. \end{aligned} \tag{16}$$

Suppose that (15) does not hold, then the numerator in (16) is negative for r sufficiently close to 1. Thus there exists point $z_0 = r_0$, $0 < r_0 < 1$ for which the quotient in (16) is negative and this negates our assumption that $f \in \mathcal{V}_{\mathcal{H}}^{n,p}(\alpha, \beta, \gamma, \mu)$. Therefore, we can conclude that it is necessary as well as sufficient that (15) holds true whenever $f \in \mathcal{V}_{\mathcal{H}}^{n,p}(\alpha, \beta, \gamma, \mu)$ and this ends the proof of Theorem 2.2.

Next we obtain both the growth and distortion results.

Theorem 2.3. *Let $f \in \mathcal{H}_{\mu}^{n,p}(\alpha, \beta, \gamma)$. Then*

$$|f(z)| \geq (1 - |b_p|)r^p - \left[\frac{p - \mu}{1 + p - \mu} - \frac{p + \mu}{1 + p - \mu} |b_p| \right] \left(\frac{\alpha + \beta p + \gamma p^2}{\alpha + \beta(1 + p) + \gamma(1 + p)^2} \right)^n r^{p+1}$$

or

$$|f(z)| \leq (1 + |b_p|)r^p + \left[\frac{p - \mu}{1 + p - \mu} - \frac{p + \mu}{1 + p - \mu} |b_p| \right] \left(\frac{\alpha + \beta p + \gamma p^2}{\alpha + \beta(1 + p) + \gamma(1 + p)^2} \right)^n r^{p+1}.$$

Proof. From (3), we have

$$f(z) = z^p + \sum_{k=1}^{\infty} a_{k+p} z^{k+p} + \overline{\sum_{k=0}^{\infty} b_{k+p} z^{k+p}}.$$

Now

$$\begin{aligned}
 |f(z)| &\leq (1 + |b_p|)r^p + \frac{p - \mu}{1 + p - \mu} \left(\frac{\alpha + \beta p + \gamma p^2}{\alpha + \beta(1 + p) + \gamma(1 + p)^2} \right)^n \times \\
 &\quad \times \sum_{k=1}^{\infty} \left[\frac{k + p - \mu}{p - \mu} |a_{p+k}| + \frac{k + p - \mu}{p - \mu} |b_{p+k}| \right] Y^n r^{p+1} \leq \\
 &\leq (1 + |b_p|)r^p + \left[\frac{p - \mu}{1 + p - \mu} - \frac{p + \mu}{1 + p - \mu} |b_p| \right] \left(\frac{\alpha + \beta p + \gamma p^2}{\alpha + \beta(1 + p) + \gamma(1 + p)^2} \right)^n r^{p+1},
 \end{aligned}$$

where Y is as defined earlier.

Theorem 2.3 is proved.

Theorem 2.4. Let $f \in \mathcal{H}_{\mu}^{n,p}(\alpha, \beta, \gamma)$. Then

$$\begin{aligned}
 |f'(z)| &\geq p(1 - |b_p|)r^{p-1} - \left[\frac{(1 + p)(p - \mu)}{1 + p - \mu} - \frac{(1 + p)(p + \mu)}{1 + p - \mu} |b_p| \right] \times \\
 &\quad \times \left(\frac{\alpha + \beta p + \gamma p^2}{\alpha + \beta(1 + p) + \gamma(1 + p)^2} \right)^n r^p
 \end{aligned}$$

or

$$\begin{aligned}
 |f'(z)| &\leq p(1 + |b_p|)r^{p-1} + \left[\frac{(1 + p)(p - \mu)}{1 + p - \mu} - \frac{(1 + p)(p + \mu)}{1 + p - \mu} |b_p| \right] \times \\
 &\quad \times \left(\frac{\alpha + \beta p + \gamma p^2}{\alpha + \beta(1 + p) + \gamma(1 + p)^2} \right)^n r^p.
 \end{aligned}$$

Proof is much similar to that of Theorem 2.2.

3. Application of fractional calculus. Given function $f(z)$ of the form (1). The fractional integral of order ϵ , $0 < \epsilon \leq 1$, is defined such that

$$D_z^{-\epsilon} f(z) = \frac{1}{\Gamma(\epsilon)} \int_0^z \frac{f(t)}{(z-t)^{1-\epsilon}} dt, \quad (17)$$

where $f(z)$ is analytic function in a simply connected region of z -plane containing the origin and the multiplicity of $(z-t)^{\epsilon-1}$ is removed by requiring $\log(z-t)$ to be real when $z-t > 0$.

Similarly, the fractional derivative of order ϵ , $0 \leq \epsilon < 1$, is given by

$$D_z^{\epsilon} f(z) = \frac{1}{\Gamma(1-\epsilon)} \frac{d}{dz} \int_0^z \frac{f(t)}{(z-t)^{\epsilon}} dt, \quad (18)$$

where $f(z)$ is as defined above and the multiplicity of $(z-t)^{-\epsilon}$ is removed by requiring $\log(z-t)$ to be real when $z-t > 0$. Interestingly both (17) and (18) have the series representations

$$D_z^{-\epsilon} f(z) = \frac{1}{\Gamma(2+\epsilon)} z^{\epsilon+1} + \sum_{k=2}^{\infty} \frac{\Gamma(k+1)}{\Gamma(k+1+\epsilon)} a_k z^{k+\epsilon}$$

and

$$D_z^\epsilon f(z) = \frac{1}{\Gamma(2-\epsilon)} z^{1-\epsilon} + \sum_{k=2}^{\infty} \frac{\Gamma(k+1)}{\Gamma(k+1-\epsilon)} a_k z^{k+\epsilon},$$

respectively (see [6, 7, 9, 11]).

Theorem 3.1. *Let $f(z)$ be of the form (3). If $f \in \mathcal{H}_\mu^{n,p}(\alpha, \beta, \gamma)$, then*

$$|D_z^{-\epsilon} f(z)| \leq \frac{\Gamma(p+1)|z|^{p+\epsilon}}{\Gamma(p+1+\epsilon)} \left\{ (1 + |b_p|) + \frac{p+1}{p+1+\epsilon} \left(\frac{p-\mu}{1+p-\mu} - \frac{p+\mu}{p+1+\epsilon} |b_p| \right) X^n |z| \right\}$$

and

$$|D_z^{-\epsilon} f(z)| \geq \frac{\Gamma(p+1)|z|^{p+\epsilon}}{\Gamma(p+1+\epsilon)} \left\{ (1 - |b_p|) - \frac{p+1}{p+1+\epsilon} \left(\frac{p-\mu}{1+p-\mu} - \frac{p+\mu}{1+p-\mu} |b_p| \right) X^n |z| \right\},$$

where

$$X = \left(\frac{\alpha + \beta p + \gamma p^2}{\alpha + \beta(1+p) + \gamma(1+p)^2} \right).$$

Proof. Following the representation of $D_z^{-\epsilon} f(z)$, we have

$$\begin{aligned} D_z^{-\epsilon} f(z) &= \frac{1}{\Gamma(\epsilon)} \int_0^z (z-t)^{-(1-\epsilon)} [f(t) + \overline{g(z)}] dt = \\ &= \frac{1}{\Gamma(\epsilon)} \left\{ \int_0^z (z-t)^{-(1-\epsilon)} \left(t^p + \sum_{k=1}^{\infty} a_{k+p} t^{k+p} \right) dt + \overline{\int_0^z (z-t)^{-(1-\epsilon)} \left(\sum_{k=0}^{\infty} a_{k+p} t^{k+p} \right) dt} \right\} = \\ &= \frac{\Gamma(p+1)}{\Gamma(p+1+\epsilon)} z^{\epsilon+p} + \sum_{k=1}^{\infty} \frac{\Gamma(k+p+1)}{\Gamma(k+p+1+\epsilon)} a_{k+p} z^{k+p+\epsilon} + \sum_{k=0}^{\infty} \frac{\Gamma(k+p+1)}{\Gamma(k+p+1+\epsilon)} b_{k+p} z^{k+p+\epsilon}. \end{aligned}$$

Then

$$\begin{aligned} \frac{\Gamma(p+1+\epsilon)}{\Gamma(p+1)} z^{-\epsilon} D_z^{-\epsilon} f(z) &= \\ &= z^p + \sum_{k=1}^{\infty} \frac{\Gamma(p+1+\epsilon)\Gamma(k+p+1)}{\Gamma(p+1)\Gamma(k+p+1+\epsilon)} a_{k+p} z^{k+p} + \sum_{k=0}^{\infty} \frac{\Gamma(p+1+\epsilon)\Gamma(k+p+1)}{\Gamma(p+1)\Gamma(k+p+1+\epsilon)} b_{k+p} z^{k+p}. \end{aligned}$$

Simple computation of the above yields

$$\begin{aligned} \left| \frac{\Gamma(p+1+\epsilon)}{\Gamma(p+1)} z^{-\epsilon} D_z^{-\epsilon} f(z) \right| &\leq \\ &\leq |z|^p + |b_p| |z|^p + \frac{p+1}{p+1+\epsilon} \left(\frac{p-\mu}{1+p-\mu} - \frac{p+\mu}{1+p-\mu} |b_p| \right) X^n |z|^{p+1}, \end{aligned}$$

where $X = \left(\frac{\alpha + \beta p + \gamma p^2}{\alpha + \beta(1+p) + \gamma(1+p)^2} \right)$. Therefore,

$$|D_z^{-\epsilon} f(z)| \leq \frac{\Gamma(p+1)|z|^{p+\epsilon}}{\Gamma(p+1+\epsilon)} \left\{ (1 + |b_p|) + \frac{p+1}{p+1+\epsilon} \left(\frac{p-\mu}{1+p-\mu} - \frac{p+\mu}{1+p-\mu} |b_p| \right) X^n |z| \right\}$$

and

$$|D_z^{-\epsilon} f(z)| \geq \frac{\Gamma(p+1)|z|^{p+\epsilon}}{\Gamma(p+1+\epsilon)} \left\{ (1 - |b_p|) - \frac{p+1}{p+1+\epsilon} \left(\frac{p-\mu}{1+p-\mu} - \frac{p+\mu}{1+p-\mu} |b_p| \right) X^n |z| \right\}.$$

Theorem 3.1 is proved.

Corollary 3.1. *Let $f(z)$ be of the form (3). If $f(z) \in \mathcal{H}_\mu^{n,1}(\alpha, \beta, \gamma)$, then*

$$|D_z^{-\epsilon} f(z)| \leq \frac{|z|^{1+\epsilon}}{\Gamma(2+\epsilon)} \left\{ (1 + |b_1|) + \frac{2}{2+\epsilon} \left(\frac{1-\mu}{2-\mu} - \frac{1+\mu}{2-\mu} |b_1| \right) X^n |z| \right\}$$

and

$$|D_z^{-\epsilon} f(z)| \geq \frac{|z|^{1+\epsilon}}{\Gamma(2+\epsilon)} \left\{ (1 - |b_1|) - \frac{2}{2+\epsilon} \left(\frac{1-\mu}{2-\mu} - \frac{1+\mu}{2-\mu} |b_1| \right) X^n |z| \right\},$$

where

$$X = \left(\frac{\alpha + \beta + \gamma}{\alpha + 2\beta + 4\gamma} \right).$$

Corollary 3.2. *Let $f(z)$ be of the form (3). If $f(z) \in \mathcal{H}_0^{n,1}(\alpha, \beta, \gamma)$, then*

$$|D_z^{-\epsilon} f(z)| \leq \frac{|z|^{1+\epsilon}}{\Gamma(2+\epsilon)} \left\{ (1 + |b_1|) + \frac{1}{2+\epsilon} (1 - |b_1|) X^n |z| \right\}$$

and

$$|D_z^{-\epsilon} f(z)| \geq \frac{|z|^{1+\epsilon}}{\Gamma(2+\epsilon)} \left\{ (1 - |b_1|) - \frac{1}{2+\epsilon} (1 - |b_1|) X^n |z| \right\},$$

where

$$X = \left(\frac{\alpha + \beta + \gamma}{\alpha + 2\beta + 4\gamma} \right).$$

Corollary 3.3. *Let $f(z)$ be of the form (3). If $f(z) \in \mathcal{H}_0^{0,1}(\alpha, \beta, \gamma)$, then*

$$|D_z^{-\epsilon} f(z)| \leq \frac{|z|^{1+\epsilon}}{\Gamma(2+\epsilon)} \left\{ (1 + |b_1|) + \frac{1}{2+\epsilon} (1 - |b_1|) |z| \right\}$$

and

$$|D_z^{-\epsilon} f(z)| \geq \frac{|z|^{1+\epsilon}}{\Gamma(2+\epsilon)} \left\{ (1 - |b_1|) - \frac{1}{2+\epsilon} (1 - |b_1|) |z| \right\}.$$

Theorem 3.2. *Let $f(z)$ be of the form (3). If $f \in \mathcal{H}_\mu^{n,p}(\alpha, \beta, \gamma)$, then*

$$|D_z^\epsilon f(z)| \leq \frac{\Gamma(p+1)|z|^{p-\epsilon}}{\Gamma(p+1-\epsilon)} \left\{ (1 + |b_p|) + \frac{p+1}{p+1-\epsilon} \left(\frac{p-\mu}{1+p-\mu} - \frac{p+\mu}{1+p-\mu} b_p \right) X^n |z| \right\}$$

and

$$|D_z^\epsilon f(z)| \geq \frac{\Gamma(p+1)|z|^{p-\epsilon}}{\Gamma(p+1-\epsilon)} \left\{ (1 - |b_p|) - \frac{p+1}{p+1-\epsilon} \left(\frac{p-\mu}{1+p-\mu} - \frac{p+\mu}{1+p-\mu} |b_p| \right) X^n |z| \right\},$$

where

$$X = \left(\frac{\alpha + \beta p + \gamma p^2}{\alpha + \beta(1+p) + \gamma(1+p)^2} \right).$$

Proof is similar to that of Theorem 3.1.

References

1. Y. Avici, E. Zlotkiewicz, *On harmonic univalent mappings*, Ann. Univ. Marie Curie-Sklodowska Sect. A, **44**, 1–7 (1990).
2. A. K. Bakhtin, I. V. Denega, *Extremal decomposition of the complex plane with free p-odes*, J. Math. Sci., **246**, № 1, 1–17 (2020).
3. N. E. Cho, H. M. Srivastava, *Argument estimates of certain analytic functions defined by a class of multiplier transformations*, Math. Comput. Modeling, **37**, № 1-2, 39–49 (2003).
4. J. Clunie, T. Sheil-Small, *Harmonic univalent functions*, Ann. Acad. Fenn. Math., **9**, 3–25 (1984).
5. I. V. Denega, *Estimate of the inner radii of non-overlapping domains*, J. Math. Sci., **242**, № 6, 787–795 (2019).
6. J. O. Hamzat, M. O. Olayiwola, *Application of fractional calculus on certain new subclasses of analytic function*, Int. J. Sci. Tech., **3**, Issue 10, 235–245 (2015).
7. Y. Komatu, *On analytic prolongation family of integral operators*, Mathematics (Cluj), **32(55)**, 141–145 (1990).
8. D. O. Makinde, J. O. Hamzat, A. M. Gbolagade, *A generalized multiplier transform on a univalent integral operator*, J. Contemp. Appl. Math., **9**, № 1, 24–31 (2019).
9. H. M. Srivastava, S. Owa, *Current topics in analytic function*, World Sci., Singapore (1992).
10. S. R. Swamy, *Inclusion properties of certain subclasses of analytic functions*, Int. Math. Forum, **7**, № 36, 1751–1760 (2012).
11. W. G. Atshan, *Fractional calculus on a subclass of Spiralike functions defined by Komatu operator*, Int. Math. Forum, **3**, № 32, 1587–1594 (2008).

Received 19.06.20