

A.I. IVANISIK

Taras Shevchenko National University of Kyiv, Faculty of Radio Physics,  
Electronics and Computer Systems

(4g, Academician Glushkov Ave., Kyiv 03127, Ukraine; e-mail: aivan@univ.kiev.ua)

**SCATTERING OF GINZBURG–FRANK  
AND CHERENKOV TYPES UNDER SELF-FOCUSING  
OF NANOSECOND LASER PULSES IN LIQUIDS<sup>1</sup>**

UDC 535.375

*We study the dynamics of nonlinear optical processes such as self-focusing, self-phase modulation, and stimulated Raman scattering in Kerr-liquids under the nanosecond laser pulse excitation. The results prove the existence of the transition Ginzburg–Frank-type effect, which promotes the appearance of new spectral components of the laser radiation at the medium boundary. The generation of extended anti-Stokes frequency-angular bands of stimulated Raman scattering is explained. When the velocity of a self-focusing focal spot matches the phase velocity of the non-linear polarization at the anti-Stokes Raman frequency and the phase velocity of the scattered axial radiation, the most intense frequency-angular bands appear. They are described by the equations typical of the Cherenkov radiation.*

*Keywords:* self-focusing, self-phase modulation, stimulated Raman scattering.

**1. Introduction**

The self-focusing (SF) of laser pulses in the nanosecond range in a Kerr medium leads to the movement of a focal spot [1]. The focal spot speed  $v_{fp}$  is defined by the laser pulse envelope. At the front and back of a pulse,  $v_{fp}$  takes positive and negative values and is not limited by the speed of light in vacuum [2].

In the practical aspect, SF creates a new situation – the dynamics of nonlinear optical processes such as the self-phase modulation (SPM) and the stimulated Raman scattering (SRS), which cannot be achieved within other technical methods.

Previously, we identified the stop point location of a focal spot [3], possibility of Cherenkov-type radiation of SRS under SF [4], effect of SF on angular spectra of SRS [5], angle-selective inverted SRS [6], frequency dependence of anti-Stokes SRS on the focal spot speed in the approach of “ideal thin lens”

[7], transition effect of SPM [8], and physical mechanism of anti-Stokes SRS of the Cherenkov type under SPM [9].

Now, it is possible to state the principles of the Ginzburg–Frank transition and the Cherenkov (or Vavilov–Cherenkov) superluminal scattering [10] under SF of nanosecond laser pulses in Kerr liquids.

**2. Consideration and Analysis**

A simplified scheme for describing the processes is presented in Fig. 1. The focal spot has velocity [2]  $v_{fp} = v_{fd}v_{gL}/(v_{fd} + v_{gL})$ , which can exceed the light speed  $c$  in vacuum ( $v_{fd}$  – velocity of focal  $z_f$  distance change,  $v_{gL}$  – group speed of laser radiation).

The laser spot has velocity  $v_{fp} = v_{fd} = 0$  for the top of a laser pulse. At this stop-point of a laser spot, classical SRS can be observed, which has asymmetric

© A.I. IVANISIK, 2018

<sup>1</sup> The paper was presented at the XXIII Galyna Puchkovska International School-Seminar “Spectroscopy of Molecules and Crystals”.




Fig. 1. Simplified scheme for describing the processes




Fig. 2. Experimental frequency-angular spectrum of the Cherenkov-type scattering (a) in the coordinates angle ( $\theta$ ) – wavelength ( $\lambda$ ) and frequency-spatial spectrum of the transition-type scattering (b) in the coordinates: lateral coordinate ( $x$ ) – wavelength ( $\lambda$ ) for a ruby laser at 20 ns, 0.5 J pulse in toluen

indicatrix for the parametric scattering. However, the indicatrix asymmetry is another question.

When the focal spot is closer to the exit of a cuvette, the Cherenkov-type superluminal scattering can be observed for parametric SRS. The experimental frequency-angular spectrum of the Cherenkov-type scattering is presented in Fig. 2, a for the first anti-Stokes (1AS) SRS in toluene under the excitation by a ruby laser in the coordinates: scattering angle ( $\theta$ ) – wavelength ( $\lambda$ ). As the focal spot crosses the exit boundary of a cuvette, Ginzburg–Frank transition-type scattering is observed for the laser radiation. An experimental frequency-spatial spectrum of the transition-type scattering is presented in Fig. 2, b for the ruby laser radiation ( $L$ ) in the coordinates: lateral coordinate ( $x$ ) – wavelength ( $\lambda$ ). The maximum Stokes frequency shifts in reverse centimeters ( $\text{cm}^{-1}$ ) are indicated in Fig. 2.

A general qualitative similarity for the axial scattering is observed. This is a result of the mutual SPM-effect of the laser radiation. The maximum Stokes frequency shift  $\Delta\nu_{1AS}$  for the first anti-Stokes SRS is about 3 times more than the laser shift  $\Delta\nu_L$ , because

$\nu_{1AS} = 2\nu_L - \nu_S$  and  $\Delta\nu_{1AS} = 2\Delta\nu_L + \Delta\nu_S = 3\Delta\nu_L$  (here  $\nu_S$  is the Stokes component frequency). The maximum Stokes frequency shift  $\Delta\nu_{2AS}$  for the second anti-Stokes SRS is about 7 times more than  $\Delta\nu_L$ , seeing  $\nu_{2AS} = 2\nu_{1AS} - \nu_L$ ,  $\Delta\nu_{2AS} = 2\Delta\nu_{1AS} + \Delta\nu_L = 7\Delta\nu_L$ .

For the maximal value  $\Delta\nu_L$  of the frequency Stokes-shift caused by the transition effect for the laser radiation, it is possible to derive the analytical expression:

$$\Delta\nu_L \approx \nu_L \Delta n_f \{z = L\} \frac{v_{fd} \{z = L\}}{c},$$

where  $z$  is a longitudinal coordinate,  $\Delta n_f \{z = L\}$  is an increment of the refractive index at the focal point at the distance  $z = L$  (at the medium boundary),  $v_{fd} \{z = L\}$  – a velocity of the focal point at the medium boundary (without considering the difference between times, which are necessary for the pulse fragments to reach the focal point), and  $c$  – the speed of light.

The maximum energy density of axial ( $\theta = 0$ ) 1AS radiation is located at the frequency determined by conditions, which are similar to those for the Cherenkov radiation: equality of the phase velocities of electromagnetic waves  $v_{ph} \{\omega\}$  at the frequency  $\omega$  and the phase velocity  $v_{ap0}$  of a polarization at the anti-Stokes Raman frequency  $\omega_a$ . The axial frequency shift in toluene is  $(\omega \{\theta = 0\} - \omega_a) / 2\pi c = -197 \text{ cm}^{-1}$ .

For  $\theta \neq 0$  and  $v_{fp} = v_{ap0}$ , the frequency-angular branches are related by the expression

$$\cos \theta \approx v_{ph}(\omega) / v_{ap0}$$

that gives a parabola for  $\cos \theta \approx 1 - \theta^2 / 2$ .

### 3. Conclusions

The Ginzburg–Frank (transition effect) – type and Cherenkov (superluminal effect) – type radiations are analyzed at the nanosecond laser pulse excitation in the spectra of a laser and SRS.

1. At the transition effect, the maximum frequency shift of the laser radiation appears, when the focal point of self-focusing intersects the exit boundary of the medium, and the phase delay of SFM before and after the focal point is uncompensated.

2. The SRS Cherenkov-type radiation appears at the coincidence of self-focusing focal point speed, the

phase velocity of a nonlinear polarization at the anti-Stokes Raman frequency and the phase velocity of scattered SRS-radiation. This is a result of the superluminal speed of the focal point and the velocity of a nonlinear polarization.

It will be of interest to analyze the mutual impact of the Ginzburg–Frank and Cherenkov effects on SRS.

1. Y.R. Shen. *The Principles of Nonlinear Optics* (Wiley-Interscience, 2003).
2. A. Ivanisik, P. Korotkov, G. Ponezha. Temporal dynamics of focal point location under self-focusing of nanosecond laser pulses. *Ukr. J. Phys. Opt.* **15**, 1 (2014).
3. A.I. Ivanisik, V.I. Malyi, G.V. Ponezha. The spatial-angular structure of the anti-Stokes emission under the stimulated Raman scattering of light in a Kerr fluid. *Optika i Spektroskopiya* **80**, 212 (1996).
4. A.I. Ivanisik, V.I. Malyi, G.V. Ponezha. Cherenkov-type radiation under conditions of Raman light scattering in self-focusing liquids. *Optics and Spectroscopy* **82**, 410 (1997).
5. A.I. Ivanisik, V.I. Malyi, G.V. Ponezha. On the influence of a self-focusing on the angular spectra of SRS. *Optika i Spektroskopiya* **85**, 88 (1998); *Optics and Spectroscopy* **85**, 78 (1998).
6. A.I. Ivanisik, V.I. Malyi, G.V. Ponezha. The spectral-angular manifestations of a competition of the Raman and parametric processes under SRS in self-focusing media. *Optika i Spektroskopiya* **85**, 512 (1998); *Optics and Spectroscopy* **85**, 469 (1998).
7. A.I. Ivanisik, G.V. Ponezha. Spectrum of anti-Stokes stimulated Raman scattering from the moving focal regions of

self-focusing. *Optika i Spektroskopiya* **90**, 699 (2001); *Optics and Spectroscopy* **90**, 625 (2001).

8. S.O. Dudka, A.I. Ivanisik, A.V. Konopatskiy, P.A. Korotkov. Transition effect at the medium vacuum interface under the self-phase modulation of a light pulse. *Ukr. J. Phys.* **51**, 140 (2006).
9. A.I. Ivanisik, O.Iu. Isaienko, P.A. Korotkov, G.V. Ponezha. Phase modulated parametric anti-Stokes stimulated Raman scattering of Cherenkov-type in self-focusing areas of exciting radiation. *Ukr. J. Phys.* **57**, 1000 (2012).
10. I.M. Frank. *Vavilov-Cherenkov Radiation: Theoretical Aspects* (Nauka, 1988).

Received 26.11.17

*A.I. Ivanisik*

РОЗСІЮВАННЯ ГІНЗБУРГА–ФРАНКА ТА  
ЧЕРЕНКОВСЬКОГО ТИПІВ ЗА САМОФОКУСУВАННЯ  
НАНОСЕКУНДНИХ ЛАЗЕРНИХ  
ІМПУЛЬСІВ У РІДИНАХ

Резюме

Досліджено динаміку нелінійно-оптичних процесів (самофокусування, фазова самомодуляція, вимушене комбінаційне розсіювання) у керівських рідинах за дії наносекундних лазерних імпульсів. Результати доводять наявність перехідного ефекту типу Гінзбурга–Франка, який породжує нові спектральні компоненти лазерного випромінювання на межі середовища. Пояснено генерацію протяжних частотно-кутових смуг вимушеного комбінаційного розсіювання. У випадку збігання швидкості фокальної точки з фазовою швидкістю нелінійної поляризації на антистоксовій комбінаційній частоті та фазовою швидкістю розсіяного аксіального випромінювання виникають найінтенсивніші смуги, які описуються рівняннями, характерними для черенковського випромінювання.