
<https://doi.org/10.15407/ujpe63.6.488>

A.M. PAVLYUK

Bogolyubov Institute for Theoretical Physics, Nat. Acad. of Sci. of Ukraine
(14b, Metrolohichna Str., Kyiv 03143, Ukraine; e-mail: pavlyuk@bitp.kiev.ua)

GENERALIZED EQUIDISTANT CHEBYSHEV POLYNOMIALS AND ALEXANDER KNOT INVARIANTS¹

We introduce the generalized equidistant Chebyshev polynomials $T^{(k,h)}$ of kind k of hyperkind h , where k, h are positive integers. They are obtained by a generalization of standard and monic Chebyshev polynomials of the first and second kinds. This generalization is fulfilled in two directions. The horizontal generalization is made by introducing hyperkind h and expanding it to infinity. The vertical generalization proposes expanding kind k to infinity with the help of the method of equidistant coefficients. Some connections of these polynomials with the Alexander knot and link polynomial invariants are investigated.

Keywords: Chebyshev polynomials, generalization, kind, hyperkind, equidistant coefficients, recurrence relation, knots and links, Alexander polynomial invariants.

1. Introduction

Knots and links appear in many areas of physics, particularly in electrodynamics, optics, liquid crystals, hydrodynamics [1–4]. The main characteristics of knots and links follow from the axioms of skein relation and normalization, which lead to polynomial invariants [5]. Each knot and link is described by some definite polynomial. The most known polynomial invariants are Alexander [6], Jones [7] and HOMFLY [8] ones.

The Chebyshev polynomials appear in different branches of mathematics, particularly in approximation theory, knot theory, combinatorics, differential equations, number theory [9]. As important mathematical instruments, they widely penetrate into different branches of physics. We exploit the Chebyshev polynomials and their generalizations for the description of knots and links [10–12]. To our mind, they have to play important role even as in the searching of new polynomial invariants.

In the present paper, we introduce the generalized equidistant Chebyshev polynomials $T^{(k,h)}$ of

kind k of hyperkind h , where k, h are positive integers. At the beginning, we unify the standard Chebyshev polynomials of the first kind and monic Chebyshev polynomials of the first kind with the help of one formula using the newly introduced notion of hyperkind h ($h = 1$ refers to all standard Chebyshev polynomials; $h = 2$ – to all monic Chebyshev polynomials). After that, a generalization means declaring h to be any positive integer and leads to generalized Chebyshev polynomials of the first kind. Analogously, one obtains generalized Chebyshev polynomials of the second kind. Thus, we fulfil the so-called horizontal generalization. The final step is making the vertical generalization, by using the method of equidistant coefficients. As a result, we expand the meanings of kind k to all positive integers and obtain the formula for generalized equidistant Chebyshev polynomials of kind k of hyperkind h .

¹ This paper was presented at the 3rd Walter Thirring International School on Fundamentals of Astroparticle and Quantum Physics (17–23 September, 2017, BITP, Kyiv, Ukraine).

2. Standard Chebyshev Polynomials

The standard Chebyshev polynomials of the *first kind* $T_n^{(1)}(x)$ are introduced as

$$T_n^{(1)}(x) = \cos(n\theta), \quad x = \cos \theta, \tag{1}$$

where n is a nonnegative integer. In our paper, the Chebyshev polynomials are characterized by two numbers $-(k, h)$. Here, the letter k stands for “*kind*”. In addition, we introduce a new notion of “*hy-perkind*”, for which we use the upper index h (in brackets); $h = 1$ refers to all standard Chebyshev polynomials. Thus, for all polynomials defined by (1), $(k, h) = (1, 1)$. In the case of monic Chebyshev polynomials considered in the next section, $h = 2$. The relation

$$\cos(n + 1)\theta + \cos(n - 1)\theta = 2 \cos \theta \cos n\theta \tag{2}$$

yields the recurrence relation for the standard Chebyshev polynomials of the first kind:

$$T_{n+1}^{(1)}(x) = 2xT_n^{(1)}(x) - T_{n-1}^{(1)}(x), \quad T_0^{(1)} = 1, \quad T_1^{(1)} = x, \tag{3}$$

which can be considered as an alternative definition of $T_n^{(1)}(x)$. Some examples of $T_n^{(1)}(x)$ obtained from (3) are as follows:

$$\begin{aligned} T_0^{(1)} &= 1, \quad T_1^{(1)} = x, \quad T_2^{(1)} = 2x^2 - 1, \\ T_3^{(1)} &= 4x^3 - 3x, \quad T_4^{(1)} = 8x^4 - 8x^2 + 1, \\ T_5^{(1)} &= 16x^5 - 20x^3 + 5x. \end{aligned} \tag{4}$$

The standard Chebyshev polynomials of the *second kind* $V_n^{(1)}(x)$ are defined in the following way ($k = 2, h = 1$):

$$V_n^{(1)}(x) = \frac{\sin(n + 1)\theta}{\sin \theta}, \quad x = \cos \theta. \tag{5}$$

The relation

$$\sin(n + 2)\theta + \sin n\theta = 2 \cos \theta \sin(n + 1)\theta \tag{6}$$

implies that definition (5) can be written in terms of the recurrence relation similar to (3):

$$V_{n+1}^{(1)}(x) = 2xV_n^{(1)}(x) - V_{n-1}^{(1)}(x), \quad V_0^{(1)} = 1, \quad V_1^{(1)} = 2x. \tag{7}$$

Some first standard Chebyshev polynomials of the second kind are:

$$\begin{aligned} V_0^{(1)} &= 1, \quad V_1^{(1)} = 2x, \quad V_2^{(1)} = 4x^2 - 1, \\ V_3^{(1)} &= 8x^3 - 4x, \quad V_4^{(1)} = 16x^4 - 12x^2 + 1, \\ V_5^{(1)} &= 32x^5 - 32x^3 + 6x. \end{aligned} \tag{8}$$

From the formula

$$2 \cos(n\theta) = \left(\frac{\sin(n + 1)\theta}{\sin \theta} - \frac{\sin(n - 1)\theta}{\sin \theta} \right), \tag{9}$$

one finds that definition (1) of $T_n^{(1)}(x)$ can be rewritten in the form

$$T_n^{(1)}(x) = \frac{1}{2} \frac{\sin(n + 1)\theta}{\sin \theta} - \frac{1}{2} \frac{\sin(n - 1)\theta}{\sin \theta}, \quad x = \cos \theta. \tag{10}$$

As follows from (9), the Chebyshev polynomials $T_n^{(1)}(x)$ and $V_n^{(1)}(x)$ are connected by the formula

$$2T_n^{(1)}(x) = V_n^{(1)}(x) - V_{n-2}^{(1)}(x), \quad n \geq 0. \tag{11}$$

Here, we mean that $V_{-1}^{(1)} = 0, V_{-2}^{(1)} = -1$, which follows from (7).

3. Monic Chebyshev Polynomials

The monic Chebyshev polynomials of the *first kind* $T_n^{(2)}(x)$, ($k = 1, h = 2$), are defined by

$$T_n^{(2)}(x) = 2 \cos(n\theta), \quad x = 2 \cos \theta. \tag{12}$$

The monic normalization means that all Chebyshev polynomials have the unit coefficients at the highest degree x^n .

The recurrence relation

$$T_{n+1}^{(2)}(x) = xT_n^{(2)}(x) - T_{n-1}^{(2)}(x), \quad T_0^{(2)} = 2, \quad T_1^{(2)} = x, \tag{13}$$

can be considered as an alternative definition of $T_n^{(2)}(x)$. Some first cases of $T_n^{(2)}(x)$ are:

$$\begin{aligned} T_0^{(2)} &= 2, \quad T_1^{(2)} = x, \quad T_2^{(2)} = x^2 - 2, \\ T_3^{(2)} &= x^3 - 3x, \quad T_4^{(2)} = x^4 - 4x^2 + 2, \\ T_5^{(2)} &= x^5 - 5x^3 + 5x. \end{aligned} \tag{14}$$

The monic Chebyshev polynomials of the *second kind* $V_n^{(2)}(x)$ are defined as ($k = 2, h = 2$)

$$V_n^{(2)}(x) = \frac{\sin(n + 1)\theta}{\sin \theta}, \quad x = 2 \cos \theta \tag{15}$$

or, in terms of the recurrence relation,

$$V_{n+1}^{(2)}(x) = xV_n^{(2)}(x) - V_{n-1}^{(2)}(x), \quad V_0^{(2)} = 1, \quad V_1^{(2)} = x. \tag{16}$$

The first monic Chebyshev polynomials of the second kind are:

$$\begin{aligned} V_0^{(2)} &= 1, \quad V_1^{(2)} = x, \quad V_2^{(2)} = x^2 - 1, \\ V_3^{(2)} &= x^3 - 2x, \quad V_4^{(2)} = x^4 - 3x^2 + 1, \\ V_5^{(2)} &= x^5 - 4x^3 + 3x. \end{aligned} \tag{17}$$

Definition (12) of $T_n^{(2)}(x)$ can be written in the form

$$T_n^{(2)}(x) = \frac{\sin(n+1)\theta}{\sin\theta} - \frac{\sin(n-1)\theta}{\sin\theta}, \quad x = 2\cos\theta. \tag{18}$$

The Chebyshev polynomials $T_n^{(2)}$ and $V_n^{(2)}$ are connected by the formula

$$T_n^{(2)} = V_n^{(2)} - V_{n-2}^{(2)}, \quad n \geq 0, \tag{19}$$

where $V_{-1}^{(2)} = 0$, $V_{-2}^{(2)} = -1$, being calculated from (16).

4. Horizontal Generalization of Chebyshev Polynomials: Expanding Hyperkind h to Infinity

The main idea for the horizontal generalization of Chebyshev polynomials consists of two steps. The first step is a unification of standard Chebyshev polynomials and monic Chebyshev polynomials of the same kinds. At the beginning, we unify ($k = 1, h = 1$) and ($k = 1, h = 2$) Chebyshev polynomials with the help of one formula; another formula unifies ($k = 2, h = 1$) and ($k = 2, h = 2$) Chebyshev polynomials. These formulas automatically lead to the second step of the horizontal generalization, by expanding the meanings of hyperkind h to infinity.

4.1. Generalized Chebyshev polynomials of the first kind

Thus, comparing (1) and (12), we write down the unified Chebyshev polynomials of the first kind $T_n^{(h)}(x)$ by the formula:

$$T_n^{(h)}(x) = h \cos(n\theta), \quad x = h \cos\theta, \tag{20}$$

for $h = 1$ and $h = 2$. From this formula, one obtains the generalized Chebyshev polynomials of the first

kind, by expanding h to infinity, i.e., h being any positive integer: $h = 1, 2, 3, 4, \dots$

Relation (2) written in the form

$$h \cos(n+1)\theta + h \cos(n-1)\theta = \frac{2}{h} h \cos\theta h \cos n\theta$$

yields the recurrence relation for the generalized Chebyshev polynomials of the first kind

$$\begin{aligned} T_{n+1}^{(h)}(x) &= \frac{2}{h} x T_n^{(h)}(x) - T_{n-1}^{(h)}(x), \\ T_0^{(h)} &= h, \quad T_1^{(h)} = x, \end{aligned} \tag{21}$$

which can be taken for the definition of $T_n^{(h)}(x)$.

Some examples of $T_n^{(h)}(x)$ obtained from (21) are:

$$\begin{aligned} T_0^{(h)} &= h, \quad T_1^{(h)} = x, \quad T_2^{(h)} = \frac{2}{h} x^2 - h, \\ T_3^{(h)} &= \frac{4}{h^2} x^3 - 3x, \quad T_4^{(h)} = \frac{8}{h^3} x^4 - \frac{8}{h} x^2 + h, \\ T_5^{(h)} &= \frac{16}{h^4} x^5 - \frac{20}{h^2} x^3 + 5x. \end{aligned} \tag{22}$$

From (20), (21), and (22), one has formulas for the standard Chebyshev polynomials of the first kind, if $h = 1$, and for the monic Chebyshev polynomials of the first kind in the case $h = 2$.

As an example, let us write (generalized) Chebyshev polynomials of the first kind of the third hyperkind. For ($k = 1, h = 3$), one has

$$T_n^{(3)}(x) = 3 \cos(n\theta), \quad 3 \cos\theta = x, \tag{23}$$

$$T_{n+1}^{(3)} = \frac{2}{3} x T_n^{(3)} - T_{n-1}^{(3)}, \quad T_0^{(3)} = 3, \quad T_1^{(3)} = x, \tag{24}$$

$$\begin{aligned} T_0^{(3)} &= 3, \quad T_1^{(3)} = x, \quad T_2^{(3)} = \frac{2}{3} x^2 - 3, \\ T_3^{(3)} &= \frac{4}{9} x^3 - 3x, \quad T_4^{(3)} = \frac{8}{27} x^4 - \frac{8}{3} x^2 + 3, \end{aligned} \tag{25}$$

$$T_5^{(3)} = \frac{16}{81} x^5 - \frac{20}{9} x^3 + 5x.$$

As follows from (9), definition (20) of $T_n^{(h)}(x)$ can be also written as

$$T_n^{(h)}(x) = \frac{h \sin(n+1)\theta}{2 \sin\theta} - \frac{h \sin(n-1)\theta}{2 \sin\theta}, \quad x = h \cos\theta. \tag{26}$$

4.2. Generalized Chebyshev polynomials of the second kind

Comparing (5) and (15), we obtain firstly unified Chebyshev polynomials of the second kind (for $h = 1$ and $h = 2$). Second, these formulas are used to introduce

the generalized ones by declaring the hyperkind h to be any positive integer.

Thus, the generalized Chebyshev polynomials of the second kind $V_n^{(h)}(x)$ are defined as

$$V_n^{(h)}(x) = \frac{\sin(n+1)\theta}{\sin\theta}, \quad x = h \cos\theta, \quad (27)$$

with h to be a positive integer. Relation (6) yields definition (27) in terms of the recurrence relation

$$V_{n+1}^{(h)} = \frac{2}{h}xV_n^{(h)} - V_{n-1}^{(h)}, \quad V_0^{(h)} = 1, \quad V_1^{(h)} = \frac{2}{h}x. \quad (28)$$

In view of (28), some first generalized Chebyshev polynomials of the second kind are as follows:

$$\begin{aligned} V_0^{(h)} &= 1, \quad V_1^{(h)} = \frac{2}{h}x, \quad V_2^{(h)} = \frac{4}{h^2}x^2 - 1, \\ V_3^{(h)} &= \frac{8}{h^3}x^3 - \frac{4}{h}x, \quad V_4^{(h)} = \frac{16}{h^4}x^4 - \frac{12}{h^2}x^2 + 1, \quad (29) \\ V_5^{(h)} &= \frac{32}{h^5}x^5 - \frac{32}{h^3}x^3 + \frac{6}{h}x. \end{aligned}$$

By substituting $h = 1$ into (29), one has standard Chebyshev polynomials of the second kind; if $h = 2$ – monic ones.

From formula (9), one finds that Chebyshev polynomials $T_n^{(h)}(x)$ and $V_n^{(h)}(x)$ are connected by the formula

$$2T_n^{(h)}(x) = h(V_n^{(h)}(x) - V_{n-2}^{(h)}(x)), \quad n \geq 0, \quad (30)$$

where $V_{-1}^{(h)} = 0, V_{-2}^{(h)} = -1$ from (28).

For example, (generalized) Chebyshev polynomials of the second kind of the third hyperkind, ($k = 2, h = 3$), are:

$$V_n^{(3)}(x) = \frac{\sin(n+1)\theta}{\sin\theta}, \quad 3 \cos\theta = x, \quad (31)$$

$$V_{n+1}^{(3)} = \frac{2}{3}xV_n^{(3)} - V_{n-1}^{(3)}, \quad V_0^{(3)} = 1, \quad V_1^{(3)} = \frac{2}{3}x, \quad (32)$$

$$V_0^{(3)} = 1, \quad V_1^{(3)} = \frac{2}{3}x, \quad V_2^{(3)} = \frac{4}{9}x^2 - 1,$$

$$V_3^{(3)} = \frac{8}{27}x^3 - \frac{4}{3}x, \quad V_4^{(3)} = \frac{16}{81}x^4 - \frac{12}{9}x^2 + 1, \quad (33)$$

$$V_5^{(3)} = \frac{32}{243}x^5 - \frac{32}{27}x^3 + 2x.$$

5. Vertical Generalization of Chebyshev Polynomials: Expanding Kind k to Infinity

The next direction of a generalization of Chebyshev polynomials is the vertical generalization. This means

expanding the meaning of kind k to infinity. The main idea for the vertical generalization is the *method of equidistant coefficients*.

We now introduce the following notation for generalized Chebyshev polynomials: $T_n^{(k,h)}(x)$. They are connected with the abovementioned polynomials in such way: $T_n^{(1,1)}(x) = T_n^{(1)}(x), T_n^{(1,2)}(x) = T_n^{(2)}(x), T_n^{(1,h)}(x) = T_n^{(h)}(x), T_n^{(2,1)}(x) = V_n^{(1)}(x), T_n^{(2,2)}(x) = V_n^{(2)}(x), T_n^{(2,h)}(x) = V_n^{(h)}(x)$. Now, we are going to generalize standard (and monic) Chebyshev polynomials by expanding the meaning of kind k .

5.1. Equidistant standard Chebyshev polynomials

The polynomials introduced here can be also called generalized Chebyshev polynomials of the first hyperkind or equidistant Chebyshev polynomials of the first hyperkind.

As an example, let us introduce the standard Chebyshev polynomials of the *third kind* $T_n^{(3,1)}(x)$, i.e., ($k = 3, h = 1$). Comparing the introductions of $T_n^{(1,1)}(x)$ by (3) and $T_n^{(2,1)}(x)$ by (7) (i.e., $T_n^{(1)}(x)$ and $V_n^{(1)}(x)$), one obtains, as a result of the natural generalization, the following definition:

$$\begin{aligned} T_{n+1}^{(3,1)}(x) &= 2xT_n^{(3,1)}(x) - T_{n-1}^{(3,1)}(x), \\ T_0^{(3,1)} &= 1, \quad T_1^{(3,1)} = 3x. \end{aligned} \quad (34)$$

Some examples of $T_n^{(3,1)}(x)$ are:

$$\begin{aligned} T_0^{(3,1)} &= 1, \quad T_1^{(3,1)} = 3x, \quad T_2^{(3,1)} = 6x^2 - 1, \\ T_3^{(3,1)} &= 12x^3 - 5x, \quad T_4^{(3,1)} = 24x^4 - 16x^2 + 1, \quad (35) \\ T_5^{(3,1)} &= 48x^5 - 44x^3 + 7x. \end{aligned}$$

By using (4), (8), and (35), it is easy to verify that

$$\begin{aligned} T_n^{(3,1)} - T_n^{(2,1)} &= T_n^{(2,1)} - T_n^{(1,1)}, \\ n &= 1, 2, 3, 4, 5, \end{aligned}$$

which demonstrates the property of equidistance of the coefficients.

By analogy, the definition of standard Chebyshev polynomials of *arbitrary kind* $T_n^{(k,1)}(x)$ is:

$$\begin{aligned} T_{n+1}^{(k,1)}(x) &= 2xT_n^{(k,1)}(x) - T_{n-1}^{(k,1)}(x), \\ T_0^{(k,1)} &= 1, \quad T_1^{(k,1)} = kx. \end{aligned} \quad (36)$$

Some first examples of $T_n^{(k,1)}(x)$ are:

$$\begin{aligned} T_0^{(k,1)} &= 1, T_1^{(k,1)} = kx, \\ T_2^{(k,1)} &= 2kx^2 - 1, \\ T_3^{(k,1)} &= 4kx^3 - (k+2)x, \\ T_4^{(k,1)} &= 8kx^4 - 4(k+1)x^2 + 1, \\ T_5^{(k,1)} &= 16kx^5 - 4(3k+2)x^3 + (k+4)x. \end{aligned} \tag{37}$$

The equidistance of the coefficients in this case is expressed by the fact that the difference

$$T_n^{(k+1,1)} - T_n^{(k,1)}$$

is independent of k for any n (due to the linearity of $T_n^{(k,1)}$ in k).

Putting $k = 1$ in (37), one has (4), $k = 2$ gives (8), and $k = 3 - (35)$.

Taking (10) and (5) in the form

$$\begin{aligned} T_n^{(1,1)}(x) &= \frac{1}{2} \frac{\sin(n+1)\theta}{\sin\theta} - \frac{1}{2} \frac{\sin(n-1)\theta}{\sin\theta}, \\ T_n^{(2,1)}(x) &= 1 \frac{\sin(n+1)\theta}{\sin\theta} + 0 \frac{\sin(n-1)\theta}{\sin\theta}, \\ x &= \cos\theta, \end{aligned} \tag{38}$$

and using the property of equidistant coefficients, one easily finds the formula for equidistant standard Chebyshev polynomials of the third kind (of the first hyperkind):

$$\begin{aligned} T_n^{(3,1)}(x) &= \frac{3}{2} \frac{\sin(n+1)\theta}{\sin\theta} + \frac{1}{2} \frac{\sin(n-1)\theta}{\sin\theta}, \\ x &= \cos\theta, \end{aligned} \tag{39}$$

and of arbitrary kind of the first hyperkind:

$$\begin{aligned} T_n^{(k,1)}(x) &= \frac{k}{2} \frac{\sin(n+1)\theta}{\sin\theta} + \frac{(k-2)}{2} \frac{\sin(n-1)\theta}{\sin\theta}, \\ x &= \cos\theta. \end{aligned} \tag{40}$$

It is easy to verify, by using (6), that (39) and (40) satisfy (34) and (36), correspondingly.

At last, we prove (40). Indeed, the general formula has two equidistant coefficients $\alpha^{(k,1)}$ and $\beta^{(k,1)}$ and looks as

$$T_n^{(k,1)}(x) = \alpha^{(k,1)} \frac{\sin(n+1)\theta}{\sin\theta} + \beta^{(k,1)} \frac{\sin(n-1)\theta}{\sin\theta}. \tag{41}$$

Introducing the difference of neighboring equidistant coefficients

$$\delta_1^{(h)} = \alpha^{(k+1,h)} - \alpha^{(k,h)}, \quad \delta_2^{(h)} = \beta^{(k+1,h)} - \beta^{(k,h)},$$

which are independent of k , we have

$$\begin{aligned} \alpha^{(k,1)} &= \alpha^{(1,1)} + \delta_1^{(1)}(k-1) = \alpha^{(1,1)} + (\alpha^{(2,1)} - \alpha^{(1,1)}) \times \\ &\times (k-1) = \frac{1}{2} + \left(1 - \frac{1}{2}\right)(k-1) = \frac{k}{2}, \\ \beta^{(k,1)} &= \beta^{(1,1)} + \delta_2^{(1)}(k-1) = \beta^{(1,1)} + (\beta^{(2,1)} - \beta^{(1,1)}) \times \\ &\times (k-1) = -\frac{1}{2} + \left(0 - \left(-\frac{1}{2}\right)\right)(k-1) = \frac{(k-2)}{2}. \end{aligned}$$

5.2. Equidistant monic Chebyshev polynomials

They can be also called generalized Chebyshev polynomials of the second hyperkind or equidistant Chebyshev polynomials of the second hyperkind.

To introduce the monic Chebyshev polynomials of the third kind $T_n^{(3,2)}(x)$, where ($k = 3, h = 2$), one must compare the introductions of $T_n^{(1,2)}(x)$ by (13) and $T_n^{(2,2)}(x)$ by (16):

$$\begin{aligned} T_{n+1}^{(3,2)}(x) &= xT_n^{(3,2)}(x) - T_{n-1}^{(3,2)}(x), \\ T_0^{(3,2)} &= 0, T_1^{(3,2)} = x. \end{aligned} \tag{42}$$

Below, we give some examples of $T_n^{(3,2)}(x)$:

$$\begin{aligned} T_0^{(3,2)} &= 0, T_1^{(3,2)} = x, T_2^{(3,2)} = x^2, \\ T_3^{(3,2)} &= x^3 - x, T_4^{(3,2)} = x^4 - 2x^2, \\ T_5^{(3,2)} &= x^5 - 3x^3 + x. \end{aligned} \tag{43}$$

Let us define monic Chebyshev polynomials of arbitrary kind $T_n^{(k,2)}(x)$:

$$\begin{aligned} T_{n+1}^{(k,2)}(x) &= xT_n^{(k,2)}(x) - T_{n-1}^{(k,2)}(x), \\ T_0^{(k,2)} &= 3 - k, T_1^{(k,2)} = x. \end{aligned} \tag{44}$$

The first examples of $T_n^{(k,2)}(x)$ are as follows:

$$\begin{aligned} T_0^{(k,2)} &= 3 - k, T_1^{(k,2)} = x, \\ T_2^{(k,2)} &= x^2 + (k-3), T_3^{(k,2)} = x^3 + (k-4)x, \\ T_4^{(k,2)} &= x^4 + (k-5)x^2 - (k-3), \\ T_5^{(k,2)} &= x^5 + (k-6)x^3 - (2k-7)x. \end{aligned} \tag{45}$$

Comparing (18) and (15) in the form

$$\begin{aligned} T_n^{(1,2)}(x) &= 1 \frac{\sin(n+1)\theta}{\sin\theta} - 1 \frac{\sin(n-1)\theta}{\sin\theta}, \\ T_n^{(2,2)}(x) &= 1 \frac{\sin(n+1)\theta}{\sin\theta} + 0 \frac{\sin(n-1)\theta}{\sin\theta}, \\ x &= 2 \cos\theta, \end{aligned} \tag{46}$$

and using equidistant coefficients, one obtains the formula for equidistant monic Chebyshev polynomials of arbitrary kind:

$$T_n^{(k,2)}(x) = \frac{\sin(n+1)\theta}{\sin\theta} + (k-2) \frac{\sin(n-1)\theta}{\sin\theta}, \quad (47)$$

$$x = 2 \cos \theta.$$

5.3. Equidistant Chebyshev polynomials of kind k of hyperkind h

Taking (26) and (27) in the form

$$T_n^{(1,h)}(x) = \frac{h}{2} \frac{\sin(n+1)\theta}{\sin\theta} - \frac{h}{2} \frac{\sin(n-1)\theta}{\sin\theta},$$

$$T_n^{(2,h)}(x) = 1 \frac{\sin(n+1)\theta}{\sin\theta} + 0 \frac{\sin(n-1)\theta}{\sin\theta}, \quad (48)$$

$$x = h \cos \theta,$$

and using the property of equidistant coefficients, one easily finds the formula for equidistant standard Chebyshev polynomials of arbitrary kind:

$$T_n^{(k,h)}(x) = \left((k-1) - (k-2) \frac{h}{2} \right) \frac{\sin(n+1)\theta}{\sin\theta} +$$

$$+ \left((k-2) \frac{h}{2} \right) \frac{\sin(n-1)\theta}{\sin\theta}, \quad x = h \cos \theta. \quad (49)$$

Indeed, the general formula has two equidistant coefficients $\alpha^{(k,h)}$ and $\beta^{(k,h)}$ and looks as

$$T_n^{(k,h)}(x) = \alpha^{(k,h)} \frac{\sin(n+1)\theta}{\sin\theta} + \beta^{(k,h)} \frac{\sin(n-1)\theta}{\sin\theta}, \quad (50)$$

where

$$\alpha^{(k,h)} = \alpha^{(1,h)} + \delta_1^{(h)}(k-1) =$$

$$= \alpha^{(1,h)} + (\alpha^{(2,h)} - \alpha^{(1,h)})(k-1) = (k-1) - (k-2) \frac{h}{2},$$

$$\beta^{(k,h)} = \beta^{(1,h)} + \delta_2^{(h)}(k-1) =$$

$$\beta^{(1,h)} + (\beta^{(2,h)} - \beta^{(1,h)})(k-1) = (k-2) \frac{h}{2}.$$

From (21) and (49), we have

$$T_{n+1}^{(k,h)}(x) = \frac{2}{h} x T_n^{(k,h)}(x) - T_{n-1}^{(k,h)}(x),$$

$$T_0^{(k,h)}(x) = (k-1) - (k-2)h = A, \quad (51)$$

$$T_1^{(k,h)}(x) = \left((k-1) \frac{2}{h} - (k-2) \right) x = Bx,$$

which can be the definition of $T_n^{(k,h)}(x)$.

At last, from (51), we have some first examples of $T_n^{(k,h)}(x)$:

$$T_0^{(k,h)} = A, \quad T_1^{(k,h)} = Bx, \quad T_2^{(k,h)} = \frac{2}{h} Bx^2 - A,$$

$$T_3^{(k,h)} = \frac{4}{h^2} Bx^3 - \left(\frac{2}{h} A + B \right) x, \quad (52)$$

$$T_4^{(k,h)} = \frac{8}{h^3} Bx^4 - \left(\frac{4}{h^2} A + \frac{4}{h} B \right) x^2 + A,$$

$$T_5^{(k,h)} = \frac{16}{h^4} Bx^5 - \left(\frac{8}{h^3} A + \frac{12}{h^2} B \right) x^3 + \left(\frac{4}{h} A + B \right) h.$$

From (49), we obtain

$$T_n^{(k,h)} = (k-1) T_n^{(2,h)} - (k-2) T_n^{(1,h)}, \quad (53)$$

and

$$T_n^{(k,h)}(x) = \left((k-1) - (k-2) \frac{h}{2} \right) T_n^{(2,h)} +$$

$$+ \left((k-2) \frac{h}{2} \right) T_{n-2}^{(2,h)}, \quad x = h \cos \theta. \quad (54)$$

6. Chebyshev Polynomials and Alexander Knot Invariants

The Alexander polynomials $\Delta(q)$ are defined by the Alexander skein relation for knots and links [6], (see notations in [13]):

$$\Delta_+(q) - \Delta_-(q) = (q^{\frac{1}{2}} - q^{-\frac{1}{2}}) \Delta_O(q), \quad \Delta_{\text{unknot}} = 1. \quad (55)$$

The Alexander polynomial invariants for torus knots $T(n, l)$ are given by the formula [14, 15]

$$\Delta_{n,l}(q) = \frac{(q^{\frac{nl}{2}} - q^{-\frac{nl}{2}})(q^{\frac{1}{2}} - q^{-\frac{1}{2}})}{(q^{\frac{n}{2}} - q^{-\frac{n}{2}})(q^{\frac{l}{2}} - q^{-\frac{l}{2}})}, \quad (56)$$

where n and l are coprime positive integers. For $l = 2$, formula (56) gives the expression for torus knots $T(n, 2)$:

$$\Delta_{n,2}^{(K)}(q) = \frac{q^{\frac{n}{2}} + q^{-\frac{n}{2}}}{q^{\frac{1}{2}} + q^{-\frac{1}{2}}}, \quad n = 2m - 1, \quad (57)$$

where m are positive integers: $m = 1, 2, 3, \dots$. Here, (K) stands for a “knot” to be distinguished from a “link” (L) . Putting $q = e^{2i\theta}$ in (57), we get

$$\Delta_{n,2}^{(K)}(q) = \frac{\cos n\theta}{\cos \theta}, \quad 2 \cos \theta = q^{\frac{1}{2}} + q^{-\frac{1}{2}}. \quad (58)$$

It easy to see that relations (20) and (58) yield

$$\frac{T_n^{(1,h)}(x)}{x} = \frac{\cos n\theta}{\cos \theta} = \Delta_{n,2}^{(K)}(q), \quad 2 \cos \theta = q^{\frac{1}{2}} + q^{-\frac{1}{2}} \quad (59)$$

for odd n . This formula says firstly that the Alexander polynomials $\Delta_{n,2}^{(K)}(q)$ are invariants (not depending on h) of all generalized Chebyshev polynomials of the first kind. Second, it gives the recipe for obtaining the corresponding Chebyshev polynomials (for odd n) from the Alexander polynomials $\Delta_{n,2}^{(K)}(q)$:

$$T_n^{(1,h)}(x) = x \Delta_{n,2}^{(K)}(q), \quad x = \frac{h}{2}(q^{\frac{1}{2}} + q^{-\frac{1}{2}}). \quad (60)$$

The corresponding formula for torus links $L(n, 2)$ (in the case of even n) [13] are:

$$\Delta_{n,2}^{(L)}(q) = \frac{q^{\frac{n}{2}} - q^{-\frac{n}{2}}}{q^{\frac{1}{2}} + q^{-\frac{1}{2}}}, \quad n = 2m. \quad (61)$$

It can be shown that, for odd n ,

$$T_n^{(2,h)}(x) = \frac{2x}{h} \frac{\Delta_{n+1,2}^{(L)}(q)}{(q^{\frac{1}{2}} - q^{-\frac{1}{2}})}, \quad x = \frac{h}{2}(q^{\frac{1}{2}} + q^{-\frac{1}{2}}). \quad (62)$$

At last, we give some Alexander polynomials, which can be calculated from (57) and (61) and can be used for the direct verification of (60) and (62):

$$\begin{aligned} \Delta_{1,2}^{(K)}(q) &= 1, \quad \Delta_{2,2}^{(L)}(q) = q^{\frac{1}{2}} - q^{-\frac{1}{2}}, \\ \Delta_{3,2}^{(K)}(q) &= q - 1 + q^{-1}, \\ \Delta_{4,2}^{(L)}(q) &= q^{\frac{3}{2}} - q^{\frac{1}{2}} + q^{-\frac{1}{2}} - q^{-\frac{3}{2}}, \\ \Delta_{5,2}^{(K)}(q) &= q^2 - q + 1 - q^{-1} + q^{-2}. \end{aligned} \quad (63)$$

7. Concluding Remarks

We have proposed a rather simple generalization of Chebyshev polynomials, which follows immediately from the attempt to unify standard and monic Chebyshev polynomials of both kinds with the help of one formula. It is believed that these polynomials are connected with some knot and link polynomial invariants. Some connections of them with Alexander polynomial invariants are presented here. We believe that a deeper investigation in this direction will bring interesting results.

The present work was partially supported by the National Academy of Sciences of Ukraine (project No. 0117U000238).

1. M.F. Atiyah. *The Geometry and Physics of Knots* (Cambridge Univ. Press, 1990).
2. L.H. Kauffman. *Knots and Physics* (World Scientific, 2001).
3. L.H. Kauffman (editor). *The Interface of Knots and Physics*. AMS Short Course Lecture Notes, a subseries of Proc. Symp. App. Math. **51** (AMS, 1996).
4. E. Radu, M.S. Volkov. Stationary ring solitons in field theory – knots and vortons. *Phys. Rep.* **468** (4), 101 (2008).
5. J.H. Conway. An enumeration of knots and links. In: *Computational Problems in Abstract Algebra* (Pergamon, 1970).
6. J.W. Alexander. Topological invariants of knots and links. *Trans. Amer. Math. Soc.* **30**, 275 (1928).
7. V.F.R. Jones. A polynomial invariant for knots and links via von Neumann algebras *Bull. AMS* **12**, 103 (1985).
8. P. Freyd, D. Yetter, J. Hoste, W.B.R. Lickorish, K. Millet, A. Ocneanu. A new polynomial invariant of knots and links. *Bull. AMS* **12**, 239 (1985).
9. T.J. Rivlin. *Chebyshev Polynomials: From Approximation Theory to Algebra and Number Theory* (Wiley, 1990).
10. A.M. Gavriliuk, A.M. Pavlyuk. On Chebyshev polynomials and torus knots. *Ukr. J. Phys.* **55**, 129 (2010).
11. A.M. Gavriliuk, A.M. Pavlyuk. Alexander polynomial invariants of torus knots $T(n,3)$ and Chebyshev polynomials. *Ukr. J. Phys.* **56**, 680 (2011).
12. A.M. Pavlyuk. On $T(n, 4)$ torus knots and Chebyshev polynomials. *Ukr. J. Phys.* **57**, 439 (2012).
13. A.M. Pavlyuk. Polynomial invariants of torus knots and (p, q) -calculus. *Algebras, Groups and Geometries* **31**, 175 (2014).
14. D. Rolfsen. *Knots and Links* (Publish or Perish, 1976).
15. W.B.R. Lickorish. *An Introduction to Knot Theory* (Springer, 1997).

Received 05.02.18

A.M. Павлюк

УЗАГАЛЬНЕНІ ЕКВІДИСТАНТНІ ПОЛІНОМИ
ЧЕБИШОВА І ВУЗЛОВІ ІНВАНІАНТИ АЛЕКСАНДЕРА

Резюме

Вводяться узагальнені еквідистантні поліноми Чебишова $T^{(k,h)}$ роду k із гіперроду h , де k, h – додатні цілі числа. Вони отримані шляхом узагальнення стандартних поліномів Чебишова першого і другого родів та монічних поліномів Чебишова першого і другого родів. Це узагальнення виконується в двох напрямках. Горизонтальне узагальнення здійснюється шляхом введення поняття гіперроду h , і розширення множини його значень до нескінченності. Вертикальне узагальнення передбачає розширення множини значень роду k до нескінченності методом еквідистантних коефіцієнтів. Досліджено деякі зв'язки введених поліномів з вузловими інваріантами Александра.