
<https://doi.org/10.15407/ujpe65.5.412>

L.N. CHRISTOPHOROV

Bogolyubov Institute for Theoretical Physics, Natl. Acad. Sci. Ukraine
(14b, Metrologichna Str., Kyiv 03143, Ukraine; e-mail: lchrist@bitp.kiev.ua)

ON THE VELOCITY OF ENZYMATIC REACTIONS IN MICHAELIS–MENTEN-LIKE SCHEMES (ENSEMBLE AND SINGLE-MOLECULE VERSIONS)

In searching non-standard ways of conformational regulation, various Michaelis–Menten-like schemes attract relentless attention, resulting in sometimes too sophisticated considerations. With the example of monomeric enzymes possessing an only binding site, we define the minimal schemes capable of bearing peculiar regulatory properties like “cooperativity” or substrate inhibition. The simplest ways of calculating the enzymatic reaction velocity are exemplified, either in the ensemble or single-molecule case.

Keywords: enzymatic reactions, Michaelis–Menten schemes, monomeric enzymes, conformational regulation, reaction velocity.

1. Introduction

The Michaelis–Menten (MM) scheme [1] serves as a basis for the enzymatic kinetics for more than a century [2, 3]. Early attempts to search for internal mechanisms of regulating the enzyme activity, rooted in the conformational lability, and corresponding deviations from the classical MM kinetics (see [4] and references therein) were practically ignored for a prolonged period [5]. Since the beginning of the 2000s, however, the situation has been changing dramatically [3], and nowadays there is no lack of theoretical papers devoted to various MM-like schemes. This is mainly conditioned by implementation of the single-molecule (SM) methods into enzymology [6], and by similarities to heterogeneous catalysis [7, 8]. Undoubtedly, this activity will be only intensified. That is why it is expedient to try assessing the intermediate achievements on this way in order to outline its further direction more thor-


oughly, and to avoid unnecessary deflections and over-complications.

With this purpose, the present work is restricted by the most basic example – precisely, by reactions of monomeric enzymes with an only binding site. In this generic case, it is especially clear which new possibilities arise due to conformational fluctuations of the enzyme, how transparent their physical nature is, how complex their (theoretical) analysis should be, and eventually, how new the results of the latter are. In Section II, the difference in the methods of calculations of the reaction velocity in the ensemble and SM versions – the point that often lacks for proper attention – is elucidated. In Section III, the minimal MM-like schemes which bear the characteristic regulation phenomena (non-monotonic dependence of the velocity on the rate of substrate release, cooperativity, and substrate inhibition) caused by the presence of conformational channels are defined, and the simplest, as distinct from frequently used, algorithm of calculating this ve-

locity is pointed out. Section IV contains concluding remarks.

2. The Classical Scheme (Ensemble and SM Versions)

This starting point of enzymatic kinetics reads:



where a is the rate constant of binding substrate S to enzyme E , forming the enzyme-substrate complex ES ; b is the rate constant of the backward “unproductive” substrate release, and R is that of the catalytic stage of converting substrate S into product P .

The kinetic equations to scheme (1) are based on the mass action law. For example, for concentration $[ES]$ of the complex, one has:

$$d[ES]/dt = -(b + R)[ES] + a[E], \quad (2)$$

where the rate constant a is in fact of the pseudo-first order, as the binding reaction is bimolecular, and $a \equiv k[S]$, where k is the second-order constant. Non-linearity of those equations is avoided due to the assumption that usually $[S]$ is much greater than $[E]$ (at least at the starting stages of the reaction) or maintained constant during many turnovers. Then Eq. (2) added with the normalization condition $[E_t] = [E] + [ES]$, where $[E_t]$ is the total enzyme concentration, turns into

$$d[ES]/dt = -(b + R + a)[ES] + a[E_t], \quad (3)$$

so that the stationary complex concentration is

$$[ES]_{\text{st}} = \frac{a[E_t]}{b + R + a} = \frac{[E_t][S]}{[S] + K_M}, \quad (4)$$

where $K_M = (b + R)/k$, the known Michaelis’ constant. Of prime interest is the stationary velocity v of product formation, $d[P]/dt$, which, according to Eqs. (1, 4), reads

$$v = R[ES]_{\text{st}} = \frac{Ra[E_t]}{a + b + R} = \frac{R[E_t][S]}{[S] + K_M}. \quad (5)$$

The hyperbolic dependence $v([S])$ (5) is the main result of scheme (1) and a primary test for investigations into any enzymatic reaction. With the progress in these investigations, the deviations from

this dependence have been revealed even for some monomeric enzymes with the only binding site. This has eventually led to the development of various schemes of such kind of biochemical regulation, crucial for many physiological processes, see, e.g., [9]. All of them are related to the splitting of the pathway of scheme (1) by introducing a set of (at least two) conformational channels of the reaction, see Section III. So far, we concentrate on the analysis of scheme (1), when the reaction is traced on a single enzyme – i.e., in a way that nowadays becomes dominant. In this case, one and the same enzyme macromolecule works in a serial regime, converting a substrate into a product consecutively, one by one. Then the obtained arrays of the turnover times (up to tens of thousands of turnovers [10]) undergo statistical processing. The prime characteristic is the probability density $f(t)$ of the first passage time, i.e. of that from the beginning to the end of a conversion act. Correspondingly, the kinetic equations related to scheme (1) are formulated not for concentrations but for probabilities $P_i(t)$ of the residence in states $i \in \{E, ES\}$. The quantity of the product released in the interval $(t, t + dt)$ is $dP = f(t)dt$; therefore, $f(t) = dP/dt = RP_{ES}(t)$, where $P_{ES}(t)$ can be found from the set

$$\begin{cases} \frac{dP_E}{dt} = -aP_E + bP_{ES}, \\ \frac{dP_{ES}}{dt} = aP_E - (b + R)P_{ES}, \end{cases} \quad (6)$$

with initial conditions $P_E(0) = 1$, $P_{ES}(0) = 0$.

At this point, it is necessary to make important remarks in order to avoid some misunderstandings occurring even in the best recent papers. Precisely, despite the similarities in appearance of, say, Eq. (2) and the second equation in (6), one has to bear in mind the obvious difference between the *stationary* problem formulation for the ensemble version and *non-stationary* one for the SM version. In the latter, in particular, there is no normalization conservation, since $P_E(t) + P_{ES}(t) = 1$ holds for $t = 0$ only, as it is obvious from Eq. (6). Moreover, $P_{E,ES}(t \rightarrow \infty) \rightarrow 0$ (in the first of Eqs. (6), the term RP_{ES} of renewing P_E due to the return of the enzyme to its free state is absent, because this stage should not contribute to $f(t)$ by definition, while it is mandatory in the stationary ensemble version!). Nevertheless, in some works (e.g., [11–13]), wishing to avoid this rather natural difference, they introduce the stage $E^0 \xrightarrow{\delta} E$ ad-




Fig. 1. Generic schemes for the effects caused by introducing different conformational channels into the classical MM scheme (see the text)

ditional to (1) and preserve the unnecessary normalization $\sum_i P_i(t) = 1$ ($i = E, ES, E^0 \dots$) with subsequent elimination of this stage in not too consistent ways. Consequently, there appear artificial and needless contradictions between the very set of kinetic equations and normalization conditions, confusing and complicating the derivation of the correct expression for $f(t)$ considerably.

Thus, solving set (6), one can obtain an explicit expression for $f(t)$ as a difference of two time exponents which is given in many works, see, e.g., [11]. The mean first passage time is $\langle t \rangle = \int_0^\infty t f(t) dt$, and the inverse quantity $1/\langle t \rangle$ serves as an analog of the reaction velocity v . To calculate $1/\langle t \rangle$, it is in fact sufficient to know the Laplace transform $\tilde{f}(s) = \int_0^\infty f(t) e^{-st} dt$, as $\langle t^m \rangle = (-1)^m d\tilde{f}(s)/ds|_{s=0}$. The Laplace transformation of set (6) gives $\tilde{P}_{ES}(s) = a[s^2 + s(a + b + R) + aR]^{-1}$. Then $\tilde{f}(s) = R\tilde{P}_{ES}(s)$, and $\langle t \rangle = (a + b + R)/aR$, so that

$$\frac{1}{\langle t \rangle} = \frac{Ra}{a + b + R}. \quad (7)$$

Comparing this with the ensemble result (5), we see their complete identity, if to adopt $v/[E_t] = 1/\langle t \rangle$. This identity has been checked experimentally [10], and Eq. (7) was named “single-molecule Michaelis–Menten equation” [11]. As it can be seen in the literature, its validity (which is considered as a version of the ergodic hypothesis) is not called in question even under complications of scheme (1) with conformational sub-states. Now, we are prepared to deal with them consistently.

3. Minimal Schemes of the Effects Caused by the Presence of Different Conformational Channels

It is advisable to confine ourselves to two channels – at least until it is desirable to stay within discrete (sub)states and understand the content of new ef-

fects within not too cumbersome analytical expressions. There exist high-quality generalizations for an arbitrary number of channels (see, e.g., [11, 14]), but they are too overloaded with mathematics, while the characteristic effects are explained within two-channel schemes all the same.

Actually, there are three such effects, and each of them can be realized within its minimal scheme, see Fig. 1.

Scheme (a) was recently proposed in [15]. It illustrates the effect of non-monotonic dependence of the velocity on the rate constant b of “unproductive” substrate release under conditions that the catalytic rate constants r and R are essentially different. The effect takes place even in the absence of conformational fluctuations, and the hyperbolic dependence $v([S])$ still holds. The latter can be replaced with a sigmoid dependence $v([S])$ in schemes of the type (b) [16, 17], if conformation E_1 of the free enzyme is more stable, but conformation E_2 binds the substrate better ($A > a$). If the return to less active sub-state E_1 is sufficiently slow (α is small), then, with $[S]$ growing, the enzyme stays in more active sub-state E_2 increasingly longer. This eventually results in imitation of cooperativity (in fact, this is a simplified version of Rabin’s scheme [4, 18]). Lastly, scheme (c), as a more general one, contains the effects listed above as its particular cases, added with the possibility of substrate inhibition.

3.1. Direct calculations of the velocity in the ensemble and SM versions

In the ensemble version, the stationary velocity can be easily found from the corresponding set of linear algebraic equations, with one of them being the conservation condition of the total enzyme concentration. In the SM version, however, one has to be more careful with calculation of this velocity and manifestation of the validity of $v/[E_t] = 1/\langle t \rangle$. Let us illustrate this, starting from the example of simplified scheme (c). To avoid cumbersome formulae, set $\beta = 0$, and also $A = a$, $B = b$. Then the only distinction of the channels is the difference in the catalytic rates, $R > r$:



In the ensemble version, the corresponding set of kinetic equations reads:

$$\begin{cases} \frac{d[E_1]}{dt} = -(a + \alpha)[E_1] + \alpha[E_2] + (b + R)[ES_1], \\ \frac{d[E_2]}{dt} = \alpha[E_1] - (a + \alpha)[E_2] + (b + r)[ES_2], \\ \frac{d[ES_1]}{dt} = a[E_1] - (b + R)[ES_1], \\ [E_1] + [E_2] + [ES_1] + [ES_2] = [E_t], \end{cases} \quad (9)$$

with its stationary solution

$$\begin{aligned} [E_1]_{st} &= [E_2]_{st} = [E_t] \frac{(b + R)(b + r)}{a(2b + R + r) + 2(b + R)(b + r)}, \\ [ES_1]_{st} &= [E_t] \frac{a(b + r)}{a(2b + R + r) + 2(b + R)(b + r)}, \\ [ES_2]_{st} &= [E_t] \frac{a(b + R)}{a(2b + R + r) + 2(b + R)(b + r)}, \end{aligned} \quad (10)$$

so that

$$\begin{aligned} v &= R[ES_1]_{st} + r[ES_2]_{st} = \\ &= [E_t] \frac{a[R(b + r) + r(b + R)]}{a(2b + R + r) + 2(b + R)(b + r)}. \end{aligned} \quad (11)$$

One can see that Michaelis' hyperbola still holds here (since $a = k[S]$), but there appears the effect of non-monotonicity of $v(b)$, provided that the inequality

$$k[S] > \frac{2rR(R + r)}{(R - r)^2} \quad (12)$$

is satisfied (a similar inequality was obtained in [15] for the scheme in Fig. 1, a).

Now, consider the SM version of scheme (8). Taking the remarks after Eq. (6) into account, the correct equations to this *non*-stationary problem read:

$$\begin{cases} \frac{dP_{E_1}}{dt} = -(a + \alpha)P_{E_1} + \alpha P_{E_2} + bP_{ES_1}, \\ \frac{dP_{E_2}}{dt} = \alpha P_{E_1} - (a + \alpha)P_{E_2} + bP_{ES_2}, \\ \frac{dP_{ES_1}}{dt} = aP_{E_1} - (b + R)P_{ES_1}, \\ \frac{dP_{ES_2}}{dt} = aP_{E_2} - (b + r)P_{ES_2}, \end{cases} \quad (13)$$

but the initial conditions should be distributed now between sub-states E_1 and E_2 . So far, designate them as $P_{E_1}(0) = n_1$, $P_{E_2}(0) = n_2$, and $P_{ES_1}(0) = P_{ES_2}(0) = 0$. Obviously, $n_1 + n_2 = 1$.

Applying the Laplace transformation to set (13) and taking into account that now $f(t) = RP_{ES_1}(t) + rP_{ES_2}(t)$, after some straightforward algebra, one has:

$$\frac{1}{\langle t \rangle} = \frac{\Delta^2}{R(\Delta_{ES_1}\Delta' - \Delta'_{ES_1}\Delta) + r(\Delta_{ES_2}\Delta' - \Delta'_{ES_2}\Delta)}, \quad (14)$$

where

$$\begin{cases} \Delta = a[ab(R + r) + (a + 2\alpha)rR], \\ \Delta' = a[(a + b + 2\alpha)(R + r) + 2\alpha b + 2rR] + \\ + 2\alpha(b + r)(b + R), \\ \Delta_{ES_1} = a[arn_1 + \alpha(b + r)], \\ \Delta'_{ES_1} = a[n_1(a + b + r) + \alpha], \\ \Delta_{ES_2} = a[aRn_2 + \alpha(b + R)], \\ \Delta'_{ES_2} = a[n_2(a + b + R) + \alpha]. \end{cases} \quad (15)$$

To find the necessary initial values of n_1 and n_2 , we note that, in the serial working regime of a single enzyme, the ratio n_1/n_2 is equal to that of stationary flows of the returns to sub-states E_1 and E_2 after the catalytic stage. That is, $n_1/n_2 = R[ES_1]_{st}/r[ES_2]_{st}$, where $[ES_{1,2}]_{st}$ are taken from Eqs. (10) of the stationary ensemble version (in fact, this was noted, for example, in [19], although sometimes n_1 and n_2 are erroneously called steady-state probabilities [11, 12]). According to (10), $n_1/n_2 = R(b + r)/r(b + R)$, then

$$\begin{aligned} n_1 &= \frac{R(b + r)}{r(b + R) + R(b + r)}, \\ n_2 &= \frac{r(b + R)}{r(b + R) + R(b + r)}. \end{aligned} \quad (16)$$

Substituting (16) into Eqs. (14), (15) and comparing the result with Eq. (11), one can eventually make sure that the identity $v/[E_t] = 1/\langle t \rangle$ really holds.

Adding conformational fluctuations of the complex and possible distinctions between the binding (unbinding) rates ($A \neq a$ and $B \neq b$, see Fig. 1, c) into consideration does not change the calculation algorithm and eventually leads to the following expression:

$$\begin{aligned} v/[E_t] &= G/H, \\ G &= \alpha[ar(B + R) + AR(b + r)] + \\ &+ \alpha\beta(A + a)(r + R) + \beta Aa(r + R), \end{aligned}$$




Fig. 2. (a) Non-monotonic $v(b)$ in the scheme in Fig. 1, a. $a = 10$. Curve 1: $R = r = 3$; curve 2: $R = 10, r = 1$. (b) Weak cooperativity in the scheme in Fig. 1, b with the return after the catalytic stage to sub-state E_2 . The flexion of $v(x)$ is negligible unless one looks at the concave in the inset for extremely small x (see the text below Eq. (20)). Here, $\alpha = 1, k_1 = a/[S] = 0.1, k_2 = A/[S] = 1, b = 10, R = 100$. (c) Substrate inhibition in the scheme in Fig. 1, c, according to Eq. (17). $\alpha = 10, \beta = 1, k_1 = A/[S] = 10, k_2 = a/[S] = 1, b = 10, B = 1, r = 1, R = 10$

$$\begin{aligned}
 H &= \alpha [a(B + R) + A(b + r) + 2(B + R)(b + r)] + \\
 &+ 2\alpha\beta (A + a + B + b + R + r + \alpha) + \\
 &+ \beta [2Aa + a(B + R) + A(b + r)], \tag{17}
 \end{aligned}$$

which, again, coincides with $1/\langle t \rangle$ obtained with the help of the Laplace transform $\hat{f}(s)$, thereby confirming this peculiar ergodic hypothesis (expression (17) coincides, to the accuracy of designations, with that obtained for $1/\langle t \rangle$ in Ref. [11], albeit in a much more complex way, via fictitious E^0 and δ , so that it was called approximate [12], while being in fact rigorous). In the general case, Eq. (17) does not follow the classical MM-dependence on $[S]$ any longer. The latter, however, still holds (at least, with rather good precision) in many limiting cases [11, 14]. On the whole, as mentioned above, the scheme in Fig. 1, c) and Eq. (17) admit realization of practically all the interesting deviations from the standard MM kinetics which are related to regulatory abilities rooted in the presence of (multiple) conformational channels: non-monotonic $v(b)$, “cooperativity”, and substrate inhibition, see Fig. 2.

3.2. Fractional polynomials

From the formal point of view, all these effects can be analyzed within the so-called fractional polynomials, i.e., expressions of the type $F_n(x)/G_n(x)$, where F_n and G_n are polynomials of the n -th order, with n depending on the number of intermediates and discrete conformational channels in an MM-like scheme (see, e.g., [4, 20]). Henceforth, x stands for substrate concentration $[S]$. In particular, for the scheme in

Fig. 1, c) $n = 2$, and $v(x)$ reads:

$$v(x) = v(\infty) \frac{x^2 + px}{x^2 + qx + c}, \tag{18}$$

where p, q , and c are positive combinations of the rate constants involved, as Eq. (17) adopts this form after replacements $A \rightarrow k_1x, a \rightarrow k_2x$. Generalizations of Eq. (18) for more complex schemes can be found in [14], although they seemingly do not lead to principally new regulation mechanisms. The general conclusions of such analyses are quite transparent, since one can arrive at them by simply equating the derivative $v'(x)$ or $v''(x)$ to zero. The presence of a positive root of the equation $v'(x) = 0$ testifies to the possibility of substrate inhibition. In our case, this condition is reduced to the quadratic equation $(q - p)x^2 + 2cx + pc = 0$ which has a positive root, if only $q < p$. In terms of parameters of the scheme in Fig. 1, c, this condition reads

$$\frac{k_1(b + r) - k_2(B + R)}{k_1(b + r) + k_2(B + R)} > \frac{\beta(R + r)}{\alpha(R - r)}, \tag{19}$$

and the greater the difference between the catalytic rates, the better the validity of Eq. (19) (see an example given in Fig. 2, c).

Possible cooperativity (i.e., flexion of $v(x)$) is related to the presence of positive roots of the equation $v''(x) = 0$. Here, the latter is cubic,

$$(q - p)x^3 + 3cx^2 + 3pcx + c(pq - c) = 0. \tag{20}$$

According the known Descartes rule, Eq. (20) can have positive roots, if only changes of signs of the

coefficients in its l.h.s. take place, i.e., if the following alternative is satisfied: $p > q$, or $p < q$, but $c > pq$. It is hardly appropriate to write down these inequalities in terms of the parameters of Eq. (17); an illustrative example is given in Fig. 2, *b*). One can notice that cooperativity is barely seen (the corresponding Hill coefficient would only slightly exceed 1), as it is a commonplace in models with discrete conformations [16, 17]. No wonder that, in many cases, the classic MM hyperbola holds practically true [11]. Examples of “phase diagrams” of regimes of cooperativity or substrate inhibition are given in [14].

As for a non-monotonic dependence of $v(b)$ on the rate of “unproductive” substrate release – the effect that attracted attention quite recently [21], albeit in an unnecessarily sophisticated way – its simplest scheme is shown in Fig. 1, *a* [15], see also Fig. 2, *a*. It includes no conformational fluctuations at all. This effect is also possible within more general scheme in Fig. 1, *c*, even in its simplified version (8) ($A = a$, $B = b$, $\beta = 0$). With b as an argument, Eq. (11) has even simpler form than (18):

$$v(b) \sim \frac{b + [2rR/(R+r)]}{b^2 + (R+r+a)b + rR + a(R+r)/2},$$

and the analysis of the equation $dv(b)/db = 0$ shows that its positive root exists under the condition

$$kx > \frac{2rR(R+r)}{(R-r)^2} \quad (21)$$

(it is equivalent to Eq. (12), since $a = kx$ here) which requires a sufficiently high substrate concentration; the greater the difference $R - r$, the lower the concentration is required. Then, staying too long in the less-active conformational sub-state ES_2 , the enzyme can return to its free state E with a chance to perform the catalytic act in a more effective way via sub-state ES_1 [15]. An analysis (again, somewhat over-complicated and involving unnecessary E^0 and δ) of this effect in another version of the scheme in Fig. 1 *a*) was given in the recent paper [13].

4. Concluding Remarks

Our main goal is to distinguish the minimal schemes of enzymatic reactions that lead to deviations from the classic MM kinetics and are a signal for the relevant above-listed effects of conformation regulation.

We point out the simplest and straightforward algorithm of calculation of the reaction velocity, avoiding unneeded complications that occur in the current literature. The reactions of monomeric enzymes with a single binding site fit these goals best.

We elucidate the important difference in the problem formulation in its ensemble and SM versions, despite the similarity of the corresponding kinetic equations. It is the difference that ensures the identity $v/[E_t] = 1/\langle t \rangle$ even in the presence of several conformational pathways of the reaction. Consequently, if interested in the velocity only, one can restrict her/himself to the stationary ensemble version, where calculations are quite trivial. Not much harder is to find $\hat{f}(s)$ in the *non*-stationary SM version in order to calculate, if needed, the higher moments $\langle t^m \rangle$ and disorder parameters.

For the demonstrative purposes, it is sufficient to involve two conformational pathways. Considering multiple pathways makes only relative sense, since, apart from producing extremely bulky expressions, it can hardly be used in practice because of too many parameters involved.

In addition, one should bear in mind that the schemes described above, as well as the overwhelming majority of similar schemes in the literature, are those of the classical *linear* kinetics based on the mass action law¹, with *constant* coefficients (rate constants). They were, are, and will be an important tool of the enzyme kinetics. Their simplicity is their undisputable merit, as well as the fact that they help to clarify the physical nature/mechanisms of the effects – but mainly qualitatively. Since, as it is well known from, for example, extensive investigations into ligand binding to myoglobin [22] or electron transfer in photosynthetic reaction centers [23], the rate “constants” of protein reactions often change in time and are distributed along “perpendicular” (i.e., structural) coordinates [16, 24]. This entails *essentially* non-exponential kinetics and *nonlinear* effects of structural memory/feedback based on an adequate treatment of the substrate-conformational interactions and inevitable, *essentially* non-equilibrium flux conditions of the enzyme functioning [25]. All these eventually lead to the molecular self-organization effects and changes of the functional regimes of the

¹ Even if the kinetic equations are formulated for the probabilities in the SM version.

enzyme by means of non-equilibrium phase transitions. We have proposed and developed a proper concept quite long ago (for more details, see, e.g., [17] and references therein).

Apart from enzymes, the MM-like schemes concern many transport processes, absorption on soft substrates, and especially nanocatalysis [7, 8], so that nowadays they are in intensive use. That is why the above-presented analysis of their features seems important for the modern theory of conformational regulation.

The present work is performed within the project 0116U002067 of the NAS of Ukraine.

1. L. Michaelis, M.L. Menten. Die Kinetik der Invertinwirkung. *Biochem. Zeitschrift* **49**, 333 (1913).
2. *A Century of Michaelis–Menten Kinetics* (Special issue, ed. by A. Cornish-Bowden, C.P. Whitham). *FEBS Lett.* **587**, 2711 (2013).
3. A. Cornish-Bowden. One hundred years of Michaelis–Menten kinetics. *Perspective in Science* **4**, 3 (2015).
4. A. Cornish-Bowden, M.L. Cárdenas. Cooperativity in monomeric enzymes. *J. Theor. Biol.* **124**, 1 (1987).
5. M.L. Cárdenas. Michaelis and Menten equation and the long road to the discovery of cooperativity. *FEBS Lett.* **587**, 2767 (2013).
6. Y.P. Lu. Sizing up single-molecule enzymatic conformational dynamics. *Chem. Soc. Rev.* **43**, 1118 (2014).
7. R. Ye, X. Mao, X. Sun, P. Chen. Analogy between enzyme and nanoparticle catalysis: A single-molecule perspective. *ACS Catal.* **9**, 1985 (2019).
8. M. Panigrahy, A. Kumar, S. Chowdhury, A. Dua. Unraveling mechanisms from waiting time distributions in single-nanoparticle catalysis. *J. Chem. Phys.* **150**, 204119 (2019).
9. V.J. Hilser, J.A. Anderson, H.N. Motlagh. Allostery vs “allokairy”, *Proc. Natl. Acad. Sci. USA* **112**, 11430 (2015).
10. B.P. English, W. Min, A.M. van Oijen, K.T. Lee, G. Luo, H. Sun, B.J. Cherayil, S.C. Kou, X.S. Xie. Ever-fluctuating single enzyme molecules: Michaelis–Menten equation revisited. *Nat. Chem. Biol.* **2**, 87 (2006).
11. S.C. Kou, B.J. Cherayil, W. Min, B.P. English, X.S. Xie. Single-molecule Michaelis–Menten equations. *J. Phys. Chem. B* **109**, 19068 (2005).
12. A. Kumar, H. Maity, A. Dua. Parallel versus off-pathway Michaelis–Menten mechanism for single-enzyme kinetics of a fluctuating enzyme. *J. Phys. Chem. B* **119**, 8490 (2015).
13. D. Singh, S. Chaudhury. Theoretical study of the conditional non-monotonic off rate dependence of catalytic reaction rates in single enzymes in the presence of conformational fluctuations. *Chem. Phys.* **523**, 150 (2019).
14. D.E. Piephoff, J. Wu, J. Cao. Conformational nonequilibrium kinetics: Generalized Michaelis–Menten equation. *J. Phys. Chem. Lett.* **8**, 3619 (2017).
15. L.N. Christophorov. Influence of substrate unbinding on kinetics of enzymatic catalysis. *Rep. Natl. Acad. Sci. Ukraine (Dopovidi)* **1**, 40 (2019).
16. L.N. Christophorov, V.N. Kharkyanen. Synergetic mechanisms of structural regulation of the electron-transfer and other reactions of biological macromolecules. *Chem. Phys.* **319**, 330 (2005).
17. L.N. Christophorov. Enzyme functioning: Along the lines of non-equilibrium phase transitions. *AIP Advances* **8**, 125326 (2018).
18. B.R. Rabin. Co-operative effects in enzyme catalysis: A possible kinetic model based on substrate-induced conformational isomerization. *Biochem. J.* **102**, 22c (1967).
19. J. Cao. Event-averaged measurements of single-molecule kinetics. *Chem. Phys. Lett.* **327**, 38 (2000).
20. W. Ferdinand. The interpretation of non-hyperbolic rate curves for two-substrate enzymes. *Biochem. J.* **98**, 278 (1966).
21. S. Reuveni, M. Urbakh, J. Klafter. Role of substrate unbinding in Michaelis–Menten enzymatic reactions. *Proc. Natl. Acad. Sci. USA* **111**, 4391 (2014).
22. N. Agmon, J.J. Hopfield. CO-binding to heme proteins: A model for barrier height distribution and slow conformational changes. *J. Chem. Phys.* **79**, 2042 (1983).
23. Yu.M. Barabash, N.M. Berezetskaya, L.N. Christophorov, A.O. Goushcha, V.N. Kharkyanen. Effects of structural memory in protein reactions. *J. Chem. Phys.* **116**, 4339 (2002).
24. N. Agmon, J.J. Hopfield. Transient kinetics of chemical reactions with bounded diffusion perpendicular to the reaction coordinate: Intramolecular processes with slow conformational changes. *J. Chem. Phys.* **78**, 6947 (1983).
25. L.N. Christophorov. Dichotomous noise with feedback and charge-conformation interactions. *J. Biol. Phys.* **22**, 197 (1996).

Received 14.11.19

Л.М. Хрестофоров

ПРО ШВИДКІСТЬ ФЕРМЕНТАТИВНИХ
РЕАКЦІЙ У МІХАЕЛІС–МЕНТЕН-ПОДІБНИХ
СХЕМАХ. АНСАМБЛЕВА
ТА ОДНОМОЛЕКУЛЯРНА ВЕРСІЇ

Резюме

У пошуках нестандартних шляхів конформаційної регуляції активно використовують різноманітні, іноді переускладнені, версії схеми Міхаеліса–Ментен. На прикладі мономерних ензимів з одним місцем зв'язування визначено мінімальні схеми, що уможливають особливі регуляторні властивості типу “кооперативності”, субстратного інгібування тощо. Проілюстровано найпростіші шляхи розрахунку швидкості ензиматичної реакції в ансамблі або на рівні поодинокій молекули ензиму.