

Jerzy Dydak

Coverings and fundamental groups: a new approach

Classical fundamental groups behave reasonably well for Poincaré spaces (i.e., semi-locally simply connected spaces). One has a construction of the universal covering for such spaces. For arbitrary spaces it is a different matter.

We define monodromy groups $\pi(p, b_0)$ for any map $p : E \rightarrow B$ with the path lifting property and any $b_0 \in B$. p is called a \mathcal{P} -covering, where \mathcal{P} is a class of Peano spaces (i.e., connected and locally path connected spaces), if it has existence and uniqueness of lifts of maps $f : X \rightarrow B$ for any $X \in \mathcal{P}$. For any B there is the maximal \mathcal{P} -covering $p_{\mathcal{P}} : B_{\mathcal{P}} \rightarrow B$ and its monodromy group is called the \mathcal{P} -fundamental group of (B, b_0) . In case of \mathcal{P} consisting of all disk-hedgehogs we construct a universal covering theory of all spaces in analogy to the classical covering theory of Poincaré spaces.

1. INTRODUCTION

The traditional approach of defining the fundamental group first and then constructing universal coverings works well only for the class of Poincaré spaces. For general spaces there were several attempts to define generalized coverings (see [1], [3], and [12]), yet there is no general theory so far that covers all path connected spaces. In this paper we plan to remedy that by changing the order of things: we define the universal covering first and its group of deck transformations is the new fundamental group of the base space.

The basic idea is that a non-trivial loop ought to be detected by a covering (not by extension over the unit disk): a loop is non-trivial if there is a covering such that some lift of the loop is a non-loop.

So it remains to define coverings: the most natural class is the class of maps that have unique disk lifting property. To make the theory work one needs to add the assumption that path components of pre-images of open sets form a basis of the total space.

2. COVERINGS AND DECK TRANSFORMATIONS

Maps are synonymous with continuous functions.

Definition 2.1. Let \mathcal{P} be a class of spaces. A map $p : E \rightarrow B$ has **\mathcal{P} -lifting property** if for any $e_0 \in E$ and any map $f : (X, x_0) \rightarrow (B, p(e_0))$, where $X \in \mathcal{P}$, there is a map $g : X \rightarrow E$ such that $p \circ g = f$ and $g(x_0) = e_0$.

p is a **\mathcal{P} -covering** (or a **\mathcal{P} covering**) if it has the \mathcal{P} -lifting property and all lifts are unique. That means $g = h$ if $g, h : X \rightarrow E$, $p \circ g = p \circ h$, and $g(x_0) = h(x_0)$ for some $x_0 \in X \in \mathcal{P}$.

Of special interest are arc-coverings (\mathcal{P} consists of the unit interval I), disk-coverings (\mathcal{P} consists of the unit disk D^2), and hedgehog-coverings (see 3.1 for the definition of hedgehogs).

Definition 2.2. A topological space X is an **lpc-space** if it is locally path-connected. X is a **Peano space** if it is locally path-connected and connected.

Problem 2.3. Suppose $p : E \rightarrow D^2$ is an arc-covering for some Peano space E . Is p a homeomorphism?

The most fundamental example of a covering is that of the identity function $id : P(X) \rightarrow X$ from the **Peanification** $P(X)$ of X to X (see [3]). $P(X)$ is obtained from X by changing its topology to the one whose basis consist of path-components of open sets in X . $id : P(X) \rightarrow X$ is a \mathcal{P} -covering for the class \mathcal{P} of all Peano spaces.

Proposition 2.4. If $p : E \rightarrow B$ is an arc-covering and E is path-connected, then the fibers of p are T_1 spaces.

Proof. A space F is T_1 if each point is closed in it. Equivalently, for any two different points $a, b \in F$ there is an open subset of F containing a but not b .

Suppose $e_0, e_1 \in p^{-1}(b_0)$ are two different points such that every neighborhood of e_0 contains e_1 . Choose a path α from e_0 to e_1 in E . Consider the loop β obtained from α by changing the value at 1 from e_1 to e_0 . Notice β is continuous ($\beta^{-1}(U) = \alpha^{-1}(U)$ for all open subsets U of E) and is a lift of the same path as α , yet ending at a different point, a contradiction. \square

2.1. The monodromy group. Suppose $p : E \rightarrow B$ is an arc-covering and $b_0 \in B$. Any loop α at b_0 induces a function from the fiber $F = p^{-1}(b_0)$ to itself that we denote by $x \rightarrow \alpha \cdot x$. Namely, we lift α to $\tilde{\alpha}$ starting at x and we put $\alpha \cdot x = \tilde{\alpha}(1)$. Notice the function $x \rightarrow \alpha \cdot x$ is a bijection: it inverse is $x \rightarrow \alpha^{-1} \cdot x$, where $\alpha^{-1}(t)$ is defined as $\alpha(1-t)$ (in other words, α^{-1} is the reverse of α). We say that α acts on F . Notice the composition of α acting on F and β acting on F is the action of the concatenation $\alpha * \beta$ on F . The basic idea is to identify any two loops that act on F the same way.

Definition 2.5. Suppose $p : E \rightarrow B$ is an arc-covering and $b_0 \in B$. The **monodromy group** $\pi(p, b_0)$ of p at b_0 is the set of equivalence classes of loops in B at b_0 : $\alpha \sim \beta$ if and only for any two lifts $\tilde{\alpha}$ (of α) and $\tilde{\beta}$ (of β) one has $\tilde{\alpha}(1) = \tilde{\beta}(1)$ if $\tilde{\alpha}(0) = \tilde{\beta}(0)$. The group operation is induced by concatenation: $[\alpha] \cdot [\beta] := [\alpha * \beta]$.

Remark 2.6. Notice the above equivalence of loops can be easily extended to the concept of equivalence of paths in B starting at b_0 . We will use that equivalence throughout the paper. In particular, by $\alpha \cdot x$ we mean $\tilde{\alpha}(1)$, where $\tilde{\alpha}$ is the lift of α starting at x .

Notice $[\alpha]$ is the trivial element of $\pi(p, b_0)$ if and only if all its lifts are loops.

Notice that, if p is a disk-covering, then any null-homotopic loop of (B, b_0) represents the trivial element of $\pi(p, b_0)$ and there is a natural homomorphism $\pi_1(B, b_0) \rightarrow \pi(p, b_0)$ that is surjective.

It is easy to show that $\pi(p, b_0)$ and $\pi(p, b_1)$ are isomorphic just as in the case of classical fundamental groups of spaces.

2.2. The deck transformation group.

Definition 2.7. Given a map $p : E \rightarrow B$ its **deck transformation group** $DTG(p)$ is the group of homeomorphisms $h : E \rightarrow E$ such that $p \circ h = p$.

Proposition 2.8. *If $p : E \rightarrow B$ is an arc-covering and E is path-connected, then the group of deck transformations $DTG(p)$ of p acts freely on E .*

Proof. Suppose $g(e) = e$ for a deck transformation g . For any $x \in E$ pick a path α from e to x . Both α and $g \circ \alpha$ are lifts of $p \circ \alpha$ originating at e . Therefore $x = \alpha(1) = (g \circ \alpha)(1) = g(x)$ and $g \equiv id_E$. \square

Definition 2.9. An arc-covering $p : E \rightarrow B$ is **regular** if for any loop α in B all its lifts are either all loops or all non-loops. This is the same as saying that $\pi(B, b_0)$ acts freely on the fiber $F = p^{-1}(b_0)$.

Notice that, if B is path-connected, regularity of p depends only on loops at a specific point. If no loop at $b_0 \in B$ has mixed lifts, then no loop at another point $b \in B$ has mixed lifts.

Proposition 2.10. *If $p : E \rightarrow B$ is a regular arc-covering and E is path-connected, then for any $e_0 \in E$ there is a natural monomorphism $DTG(p) \rightarrow \pi(p, b_0)$, $b_0 = p(e_0)$. The monomorphism is an isomorphism if $DTG(p)$ acts transitively on the fibers of p .*

Proof. For any $h \in DTG(p)$ choose a path α_h in E from e_0 to $h(e_0)$. Since p is a regular arc-covering, the equivalence class $[p \circ \alpha_h]$ does not depend on the choice of α_h . If $g \in DTG(p)$, then $\alpha_g * g(\alpha_h)$ is a path from e_0 to $g(h(e_0))$ and $[p(\alpha_g * g(\alpha_h))] = [p(\alpha_g) * p(\alpha_h)]$, so it is indeed a homomorphism.

If $DTG(p)$ acts transitively on the fibers of p and $[\alpha] \in \pi(p, b_0)$, then lift α to $\tilde{\alpha}$ and pick a deck transformation h such that $h(\tilde{\alpha}(0)) = h(\tilde{\alpha}(1))$. Notice h is mapped to $[\alpha]$. \square

Problem 2.11. *Characterize continuous group actions G on a Peano space E such that the projection $p : E \rightarrow E/G$ is an arc-covering.*

Problem 2.12. *Characterize continuous group actions G on a Peano space E such that the projection $p : E \rightarrow E/G$ is a disk-covering.*

3. HEDGEHOG COVERINGS

Definition 3.1. A **directed wedge** (see [3]) is the wedge

$$(Z, z_0) = \bigvee_{s \in S} (Z_s, z_s)$$

of pointed Peano spaces indexed by a directed set S and equipped with the following topology (all wedges in this paper are considered with that particular topology):

- (1) $U \subset Z \setminus \{z_0\}$ is open if and only if $U \cap Z_s$ is open for each $s \in S$,
- (2) U is an open neighborhood of z_0 if and only if there is $t \in S$ such that $Z_s \subset U$ for all $s > t$ and $U \cap Z_s$ is open for each $s \in S$.

A **arc-hedgehog** is a directed wedge $(Z, z_0) = \bigvee_{s \in S} (Z_s, z_s)$ such that each (Z_s, z_s)

is homeomorphic to $(I, 0)$. The **standard arc-hedgehog** is the arc-hedgehog over the set of natural numbers N .

A **disk-hedgehog** is a directed wedge $(Z, z_0) = \bigvee_{s \in S} (Z_s, z_s)$ such that each Z_s is homeomorphic to the 2-disk D^2 .

A typical construction of an arc-hedgehog and its map to a space X is the following:

Proposition 3.2. *Let $x_0 \in X$. Suppose $\{\alpha_V : I_V = [0, 1] \rightarrow X\}_{V \in S}$ is a family of paths in X indexed by a basis S of open neighborhoods V of x_0 in X . If $\alpha_V(I) \subset V$ and $\alpha_V(0) = x_0$ for all $V \in S$ and S is ordered by inclusion ($U \leq V$ means $V \subset U$), then the natural function $\alpha = \bigvee_{V \in S} \alpha_V : \bigvee_{V \in S} (I_V, 0) \rightarrow X$ is continuous.*

Proof. $\alpha^{-1}(U)$ is certainly open if $x_0 \notin U$. If $x_0 \in U$, then $I_V \subset \alpha^{-1}(U)$ for all $V \subset U$, so α is indeed continuous. \square

Corollary 3.3. *Suppose $f : Y \rightarrow X$ is a function from an lpc-space Y . f is continuous if $f \circ g$ is continuous for every map $g : Z \rightarrow Y$ from an arc-hedgehog Z to Y .*

Proof. Assume U is open in X and $x_0 = f(y_0) \in U$. Suppose for each path-connected neighborhood V of y_0 in Y there is a path $\alpha_V : (I, 0) \rightarrow (V, y_0)$ such that $\alpha_V(1) \notin f^{-1}(U)$. Notice the wedge $\alpha = \bigvee_{V \in S} \alpha_V$ is a map from an arc-hedgehog to Y by 3.2 (here S is the family of all path-connected neighborhoods of y_0 in Y). Hence $h = f \circ g$ is continuous and there is $V \in S$ so that $I_V \subset h^{-1}(U)$. That means $f(\alpha_V(I)) \subset U$, a contradiction. \square

Remark 3.4. If X is first countable (it has a countable basis at each point) in 3.2 or Y is first countable in 3.3, then it is sufficient to use the standard arc-hedgehog only.

Theorem 3.5. *If $p : E \rightarrow B$ is an arc-covering, then the following conditions are equivalent:*

- a. p is an arc-hedgehog covering,
- b. given an open subset U of E containing e_0 , there is a neighborhood V of b_0 in B such that the path component of $p^{-1}(V)$ containing e_0 is a subset of U .

Proof. a) \implies b). Suppose, for every neighborhood V of b_0 in B , there is a path α_V in $p^{-1}(V)$ joining e_0 with a point in $E \setminus U$. The function $\alpha = \bigvee_{V \in S} \alpha_V :$

$H = \bigvee_{V \in S} I_V \rightarrow E$ is continuous as $p \circ \alpha$ is continuous and α is the only possible lift of $p \circ \alpha$ at e_0 . However, the point-inverse of U under α contains e_0 but none of I_V is contained in it, a contradiction.

b) \implies a). Suppose $\alpha = \bigvee_{s \in S} \alpha_s : \bigvee_{s \in S} I_s \rightarrow B$ is a map of an arc-hedgehog with the base-point mapped to $b_0 = p(e_0)$. The only possible lift β of α must be obtained by lifting each α_s separately. The only issue is the continuity of β at the base-point. Given a neighborhood U of e_0 in E , pick a neighborhood V of b_0 in B with the property that the path component P of $p^{-1}(V)$ containing e_0 is a subset of U . Pick an open subset W of the base-point of H satisfying $W \subset \alpha^{-1}(V)$ so that W is path-connected. Notice $\beta(W) \subset P \subset U$, which means β is continuous at the base-point of H . \square

Corollary 3.6. *If B is first countable and $p : E \rightarrow B$ is an arc-covering with E being a Peano space, then p is an arc-hedgehog covering.*

Proof. Suppose $b_0 \in B$ and $\{U_n\}$ is a decreasing basis of neighborhoods of b_0 in B . Given $e \in F = p^{-1}(b_0)$ and a neighborhood V of e in E , assume that for every $n \geq 1$ there is a path α_n in $p^{-1}(U_n)$ joining e to a point $e_n \in E \setminus V$. Consider the infinite concatenation $p(\alpha_1) * p(\alpha_1^{-1}) * p(\alpha_2) * p(\alpha_2^{-1}) * \dots$ which we assume ends at b_0 . The lift γ of β starting at e cannot be a loop as $\gamma^{-1}(V)$ does not contain any e_n . So it ends at a different point of F . Pick a neighborhood W of $\gamma(1)$ not containing e (see 2.4). $\gamma^{-1}(W)$ is a neighborhood of 1 in $[0, 1]$. Therefore infinitely many paths α_n lie in W , a contradiction. \square

Corollary 3.7. *If $p : E \rightarrow B$ is an arc-hedgehog covering and E is a Peano space, then the fibers of p are regular (T_3 -spaces) 0-dimensional spaces.*

Proof. By 2.4, fibers of p are T_1 -spaces, so, given $x \notin A$ in a fiber F (and A being closed in F), there is an open neighborhood V of $p(x) = p(y)$ such that the path component W of $p^{-1}(V)$ containing x does not intersect A . The restriction $W \cap F$ of W to F is an open-closed subset of F containing x and missing A . \square

Corollary 3.8. *Arc-hedgehog coverings $p : E \rightarrow B$ are open if both E and B are locally path-connected.*

Proof. Suppose U is open in E and $e_0 \in U$. Put $b_0 = p(e_0)$ and $F = p^{-1}(b_0)$. By 3.5 there is a path-connected neighborhood V of b_0 such that the path-component of e_0 in $p^{-1}(V)$ is a subset of U . Therefore $V \subset p(U)$ (connect e_0 with a path to any point in V and then lift the path - it must be contained in U). \square

Here is an important supplement to 2.10:

Theorem 3.9. *Suppose $p : E \rightarrow B$ is an arc-hedgehog covering. If E is a Peano space, then p is regular if and only if the deck transformation group $DTG(p)$ acts transitively on the fibers of p .*

Proof. If $DTG(p)$ acts transitively on the fibers of p , then for any two lifts α and β of the same loop in B there is a deck transformation h such that $h \circ \alpha = \beta$. Hence they are either both loops or both non-loops.

Suppose p is regular and $e_1, e_2 \in E$ with $p(e_1) = e_2$. Given $x \in E$ choose a path α_x in E from e_1 to x and let β_x be the path from e_2 to $h(x)$ with the property $p \circ \alpha_x = p \circ \beta_x$. Notice $h(x)$ does not depend on the choice of α_x as p is regular.

The reason h is continuous is that $h \circ f$ is continuous for any map f from an arc-hedgehog to E . Since analogous construction creates the inverse of h , it is a homeomorphism. \square

Proposition 3.10. *Suppose $p : E \rightarrow B$ is an arc-hedgehog covering of Peano spaces. If B is metrizable, then E is metrizable.*

Proof. Denote r -balls in B centered at b by $B(b, r)$. Define $d(x, y)$ as the infimum of $r > 0$ such that there is a path α from x to y in E with $p(\alpha([0, 1])) \subset B(p(x), r) \cap B(p(y), r)$. Clearly, d is symmetric. Also, $d(x, y) = 0$ implies $x = y$. Indeed, $p(x) = p(y)$ and $x \neq y$ would imply existence of a neighborhood U of $p(x)$ in B such that no path in U can be lifted to a path from x to y (see 3.5).

The proof of the Triangle Inequality is left to the reader.

Given $x \in U$, U open in E , find an $r > 0$ such that the path component of $p^{-1}(B(p(x), r))$ containing x is contained in U (see 3.5). Therefore the r -ball of metric d centered in x is contained in U .

Consider the r -ball $B_d(x, r)$ in d centered at $x \in E$. Look at the path-component U of $p^{-1}(B(p(x), r/2))$ containing x . It must be contained in $B_d(x, r)$ which completes the proof. \square

Proposition 3.11. *If $p : E \rightarrow B$ an arc-hedgehog covering, E is Peano, and B has a countable basis at b_0 , then $F = p^{-1}(b_0)$ is a Baire space.*

Proof. Let $\{U_n\}$ be a basis of open sets at b_0 that forms a decreasing sequence. We plan to show that, given a decreasing sequence $\{V_n\}$ of path-components V_n of $p^{-1}(U_n)$, the intersection $F \cap \bigcap_{n=1}^{\infty} V_n$ is not empty. By induction, pick points $e_n \in V_n$ and paths α_n in V_n joining e_n with e_{n+1} . The infinite concatenation $p(\alpha_1) * p(\alpha_2) * \dots$ (its end-point is declared to be b_0) is a path α in U_1 . Lift α starting at e_1 and notice the end-point of the lift belongs to $F \cap \bigcap_{n=1}^{\infty} V_n$. \square

Remark 3.12. Combining the proofs of 3.10 and 3.11 one can show E is completely metrizable if B is completely metrizable and both E and B are Peano spaces.

Definition 3.13. Suppose $p : E \rightarrow B$ is an arc-hedgehog covering of Peano spaces. p is **trivial** at b_0 if there is a connected neighborhood U of b_0 in B such that p maps each component of $p^{-1}(U)$ homeomorphically onto U .

Theorem 3.14. *Suppose $p : E \rightarrow B$ is a regular arc-hedgehog covering of Peano spaces. p is trivial at b_0 if and only if the fiber $F = p^{-1}(b_0)$ contains an isolated point.*

Proof. One direction is obvious, so assume F has an isolated point $e \in F$. Choose a connected neighborhood U of b_0 in B such that the path component V of $p^{-1}(U)$ containing e does not intersect $F \setminus \{e\}$ (see 3.5). Notice p maps V homeomorphically onto U . Indeed, $p(V) = U$ (lift a path from b_0 to any $x \in U$ starting from e to arrive at $y \in V$ such that $p(y) = x$) and $p|_V$ has to be injective: if $p(y) = p(z) = b$ for two different points $y, z \in V$, then there is a path β in V from y to z such that $p \circ \beta$ is a loop and picking a path γ from e to y in V results in a loop $p(\gamma) * p(\beta) * p(\gamma^{-1})$ in U that has a lift in V starting at e and ending at a different point contrary to $V \cap F = \{e\}$.

Consider another component W of $p^{-1}(U)$. Using 3.9 one can see there is a deck transformation h such that $h(V) = W$. Therefore $p|_W : W \rightarrow U$ is a homeomorphism as well. \square

Proposition 3.15. *If $p : E \rightarrow B$ is an arc-hedgehog covering, then p is a disk-hedgehog covering if and only if it is a disk covering.*

Proof. It only suffices to consider the case p is a disk-covering (the other implication is obvious). Given a map $f : H \rightarrow B$ from a disk-hedgehog to B and given $e \in E$ in the fiber of p over the base-point there is only one candidate for the lift of f . That candidate must be continuous as otherwise we would generate a map from an arc-hedgehog to B that has no lift at e . \square

4. THE WHISKER TOPOLOGY

In this section we are generalizing the whisker topology that was introduced in [3] in a special case.

Definition 4.1. Let B be a space and $b_0 \in B$. Suppose \sim is an equivalence relation on the set of loops in B at b_0 which induces a group structure on the set of equivalence classes via $[\alpha] \cdot [\beta] := [\alpha * \beta]$ with the constant loop at b_0 being the neutral element and $[\alpha]^{-1} = [\alpha^{-1}]$ for all loops α, β at b_0 .

The above can be summarized as follows:

1. $\alpha \sim \beta$ and $\gamma \sim \omega$ implies $\alpha * \gamma \sim \beta * \omega$ for all loops $\alpha, \beta, \gamma, \omega$ at b_0 ,
2. $\alpha * \alpha^{-1} \sim \text{const}$ and $\alpha \sim \alpha * \text{const}$ for all loops α , where α^{-1} is the reversed path of α .

The above equivalence relation can be extended to an equivalence relation on the set of all paths in B originating at b_0 : $\alpha \sim \beta$ means $\alpha(1) = \beta(1)$ and $\alpha * \beta^{-1} \sim \text{const}$.

By the **whisker topology** on the space $P(B, b_0, \sim)$ of equivalence classes $[\alpha]$ we mean the topology with the basis $N([\alpha], U)$, U an open set in B containing $\alpha(1)$, consisting of all $[\beta]$ such that $\beta \sim \alpha * \gamma$ for some path γ in U

Theorem 4.2.

- a. $P(B, b_0, \sim)$ is a Peano space and the end-point projection $p : P(B, b_0, \sim) \rightarrow B$ has arc-lifting property.
- b. p is an arc-hedgehog covering if and only if it is an arc-covering.
- c. p is a disk-hedgehog covering if and only if it is an arc-covering and $\alpha \sim \text{const}$ for every loop α at b_0 that is null-homotopic.

Proof. a. Notice $\lambda \in N([\alpha], U) \cap N([\beta], V)$ implies $N([\lambda], U \cap V) \subset N([\alpha], U) \cap N([\beta], V)$, so it is indeed a topology.

Given α at any point of B let α_t be the path equal to α on the interval $[0, t]$ and then being a constant path. If γ is a path in U originating at $\alpha(1)$, then each $[\alpha * \gamma_t] \in N([\alpha], U)$ and $t \rightarrow [\alpha * \gamma_t]$ is continuous (indeed, the inverse of $N([\alpha * \gamma_t], V)$ contains the interval around t that is mapped under γ to V). That means $P(B, b_0, \sim)$ is a Peano space. At the same time it implies p has arc-lifting property.

b. Suppose p is an arc-covering, U is open in B , and α is a path in B starting at b_0 and ending at a point in U . It suffices to show $N([\alpha], U)$ is the path component of $p^{-1}(U)$ containing $[\alpha]$. Suppose $\tilde{\gamma}$ is a path in $p^{-1}(U)$ starting at $[\alpha]$. Put $\gamma = p \circ \tilde{\gamma}$ and notice $t \rightarrow [\alpha * \gamma_t]$ is another lift of γ . Thus $\tilde{\gamma}(t) = [\alpha * \gamma_t]$ for all t proving that $\tilde{\gamma}$ is a path in $N([\alpha], U)$. In view of 3.5, p is an arc-hedgehog covering.

c. Assume p is an arc-covering and $\alpha \sim \text{const}$ for every loop α at b_0 that is null-homotopic. In view of b) and 3.15 it suffices to show p is a disk-covering.

Suppose $f : D^2 \rightarrow B$ and α is a path in B from b_0 to $f(d)$ for some $d \in D^2$. Given $x \in D^2$ let β_x be a path in D^2 from d to x . Define $g(x) \in P(B, b_0, \sim)$ as $g(x) = [\alpha * (f \circ \beta_x)]$ and notice $g(x)$ does not depend on the choice of β_x . Given a map $u : H \rightarrow D^2$ from an arc-hedgehog, $g \circ u$ is the lift of $f \circ u$, hence it is continuous. Therefore g is continuous. \square

Here is an inner description of arc-hedgehog coverings:

Theorem 4.3. *Suppose E is a Peano space. If $p : E \rightarrow B$ is an arc-hedgehog covering and $b_0 \in B$, then p is equivalent to the end-point projection $P(B, b_0, \sim) \rightarrow B$, where $P(B, b_0, \sim)$ is equipped with the whisker topology.*

Proof. Pick $e_0 \in E$ with $p(e_0) = b_0$ and declare two paths α and β in B originating at b_0 equivalent if $\alpha \cdot b_0 = \beta \cdot b_0$.

Given a point $x \in E$ choose a path α_x in E from e_0 to x and define $h : E \rightarrow P(B, e_0)$ by $h(x) = [p \circ \alpha_x]$.

Since $h^{-1}(N([\alpha_x], U))$ is the path-component of $p^{-1}(U)$ containing x , it is open in E and h is continuous.

If U is an open neighborhood of x in E , choose an open neighborhood V of $p(x)$ with the property that the path component W of x in $p^{-1}(V)$ is contained in U . Notice $N([\alpha_x], V) \subset h(W) \subset h(U)$, so h is open. Since h is bijective, it is a homeomorphism. \square

5. SUPRENUMS OF COVERINGS

Two coverings $p_1 : E_1 \rightarrow B$ and $p_2 : E_2 \rightarrow B$ are said to be **equivalent** if there is a homeomorphism $h : E_1 \rightarrow E_2$ satisfying $p_2 \circ h = p_1$. It turns out there is a set of coverings over B such that any disk-hedgehog covering over B is equivalent to one from that set. In that sense we may talk about the set of all disk-hedgehog coverings over B .

In this section we define a partial order on the set of all disk-hedgehog coverings over a fixed path-connected space B and we show this set has a maximum. That maximum plays the role of the universal covering space.

Definition 5.1. Suppose E_1, E_2 are Peano spaces and $p_1 : E_1 \rightarrow B, p_2 : E_2 \rightarrow B$ are disk-hedgehog coverings. We define the inequality $(p_1, e_1) \geq (p_2, e_2)$ of pointed coverings as follows: $p_1(e_1) = p_2(e_2)$ and there is a continuous function $f : E_1 \rightarrow E_2$ satisfying $p_2 \circ f = p_1$ and $f(e_1) = e_2$.

We define the inequality of unpointed coverings $p_1 \geq p_2$ as follows: for every points $e_1 \in E_1$ and $e_2 \in E_2$ such that $p_1(e_1) = p_2(e_2)$ we have $(p_1, e_1) \geq (p_2, e_2)$.

Lemma 5.2. *If $(p_1, e_1) \geq (p_2, e_2)$ and $(p_2, e_2) \geq (p_1, e_1)$, then there is a homeomorphism $h : E_2 \rightarrow E_1$ such that $h(e_2) = e_1$ and $p_1 \circ h = p_2$.*

Proof. Choose continuous functions $f : E_1 \rightarrow E_2$ and $g : E_2 \rightarrow E_1$ such that $p_2 \circ f = p_1, p_1 \circ g = p_2$ and $f(e_1) = e_2, g(e_2) = e_1$. As $p_1 \circ (g \circ f) = p_1$ and $(g \circ f)(e_1) = e_1$, we get $g \circ f = \text{id}_{E_1}$. Similarly, $f \circ g = \text{id}_{E_2}$. \square

Lemma 5.3. *If p_1 is a regular disk-hedgehog covering and $(p_1, e_1) \geq (p_2, e_2)$, then $p_1 \geq p_2$.*

Proof. Choose a continuous function $f : E_1 \rightarrow E_2$ such that $p_2 \circ f = p_1$. Notice f is surjective. Given $x_1 \in E_1$ and $x_2 \in E_2$ satisfying $p_1(x_1) = p_2(x_2)$ choose a deck transformation $h : E_1 \rightarrow E_1$ so that $h(x_1) \in f^{-1}(x_2)$ (see 3.9). Put $g = f \circ h$ and notice $p_2 \circ g = p_1$, $g(x_1) = x_2$. \square

Corollary 5.4. $p \geq p$ if and only if p is regular.

Proof. In view of 5.3 it suffices to show p is regular if $p \geq p$. That follows from 3.9 as any $f : E \rightarrow E$ satisfying $p \circ f = p$ must be a homeomorphism. \square

Definition 5.5. Suppose $\{p_s : E_s \rightarrow B\}_{s \in S}$ is a family of disk-hedgehog coverings of Peano spaces over a path-connected B and $e_s \in E_s$ so that $p_s(e_s) = b_0$ for all $s \in S$. (p, e) is the **supremum** of $\{(p_s, e_s)\}_{s \in S}$ if $(p, e) \geq (p_s, e_s)$ for all $s \in S$ and (p, e) is the smallest pointed covering with that property.

Definition 5.6. Suppose $\{p_s : E_s \rightarrow B\}_{s \in S}$ is a family of disk-hedgehog coverings of Peano spaces over a path-connected B and $e_s \in E_s$ so that $p_s(e_s) = b_0$ for all $s \in S$.

The **Peano fibered product** of $\{(p_s, e_s)\}_{s \in S}$ is the pair (p, e) , where $p : E \rightarrow B$, $e = \{e_s\}_{s \in S}$, and E is the Peanification of the path-component of e in the subset of $\prod_{s \in S} E_s$ consisting of points $\{x_s\}_{s \in S}$ such that $p_s(x_s) = p_t(x_t)$ for all $s, t \in S$. The projection p is defined by $p(\{x_s\}_{s \in S}) = p_t(x_t)$ for any $t \in S$.

Proposition 5.7. *Peano fibered product of a family of pointed disk-hedgehog coverings is the supremum of that family.*

Proof. If $q : E' \rightarrow B$ and $(q, e') \geq (p_s, e_s)$ for all $s \in S$, then there are maps $g_s : E' \rightarrow E_s$ so that $q = p_s \circ g_s$ and $g_s(e') = e_s$ for each $s \in S$. The collection $\{g_s\}_{s \in S}$ induces a map $g : E' \rightarrow E$ satisfying $g(e') = e$ and $p \circ g = q$. Thus $(q, e') \geq (p, e)$.

Suppose $b_0 = p(\{e_s\}_{s \in S})$, $\{e_s\}_{s \in S} \in E$, and $f : (H, 0) \rightarrow (B, b_0)$ is a map from a disk-hedgehog. Create lifts $f_s : (H, 0) \rightarrow (E_s, e_s)$ of f with respect to p_s . That defines a map $f : H \rightarrow E$ by $f(x) = \{f_s(x)\}_{s \in S}$ that is a lift of f with respect to p .

That proves existence of lifts - a proof of uniqueness is obvious. \square

Proposition 5.8. *If $p : E \rightarrow B$ is a disk-hedgehog covering and $e_0 \in E$, then the Peano fibered product of all $p : (E, e) \rightarrow (B, p(e_0))$, e ranging over all points in the fiber F of p containing e_0 , is regular.*

Proof. Suppose α is a loop in B at $b_0 = p(e_0)$ such that for some $\{x_e\}_{e \in F}$ in the fiber of q , $\alpha \cdot \{x_e\}_{e \in F} = \{x_e\}_{e \in F}$. That means $\alpha \cdot x_e = x_e$ for all $e \in F$.

Since both $\{x_e\}_{e \in F}$ and $\{e\}_{e \in F}$ can be joined by a path in the Peano fibered product, there is a loop β at b_0 in B such that $\beta \cdot \{e\}_{e \in F} = \{x_e\}_{e \in F}$. Thus $\beta \cdot e = x_e$ and $(\alpha * \beta) \cdot e = \beta \cdot e$ for all $e \in F$. Plugging in $\beta^{-1} \cdot e \in F$ for e in the equation $(\alpha * \beta) \cdot e = \beta \cdot e$ gives $\alpha \cdot e = e$ for all $e \in F$. That implies $\alpha \cdot \{y_e\}_{e \in F} = \{y_e\}_{e \in F}$ for all $\{y_e\}_{e \in F}$ in the fiber of q , i.e. q is regular. \square

Notice the Peano fibered product of all $z \rightarrow z^n$ is the covering $t \rightarrow \exp(2\pi ti)$ of reals over the unit circle.

Corollary 5.9. *Every path-connected space B has a maximal disk-hedgehog covering among those with total space being Peano. It is a regular covering.*

Proof. Pick b_0 and consider the space of paths $P(B, b_0)$ in B starting at b_0 . For every disk-hedgehog covering $p : E \rightarrow B$, E is an image of a function from

$P(B, b_0)$ obtained by lifting paths (the lifts start at $e_0 \in p^{-1}(b_0)$). That means there is a set $\{p_s : E_s \rightarrow B\}_{s \in S}$ of disk-hedgehog coverings with the property that for any disk-hedgehog covering $p : E \rightarrow B$ there is $s \in S$ and a homeomorphism $h : E \rightarrow E_s$ such that $p = p_s \circ h$. We only consider disk-hedgehog with Peano total space. Take the Peano fibered product of $\{p_s : E_s \rightarrow B\}_{s \in S}$. It must be a regular disk-hedge covering but it is easier to use 5.8 and produce the maximal covering that is regular. \square

6. HEDGEHOG FUNDAMENTAL GROUP

Definition 6.1. Given a path-connected space B and $b_0 \in B$ define the **hedgehog fundamental group** $\pi(B, b_0)$ of (B, b_0) as the monodromy group $\pi(p, b_0)$, where $p : E \rightarrow B$ is the maximal disk-hedgehog covering over B .

Proposition 6.2. Any map $f : B_1 \rightarrow B_2$ of path-connected spaces induces a natural homomorphism from $\pi(B, b_1)$ to $\pi(B_2, f(b_1))$.

Proof. Let $f(b_1) = b_2$. Consider the maximum disk-hedgehog covering $p_2 : E_2 \rightarrow B_2$ and pick $e_2 \in p_2^{-1}(b_2)$. Take the path-component of (b_1, e_2) in $\{(x, y) \in B_1 \times E_2 \mid f(x) = p_2(y)\}$, Peanify it to get E and let $q : E \rightarrow B_1$ be the projection onto the first coordinate. Notice q is a disk-hedgehog covering. Let $p : E_1 \rightarrow E$ be the maximum disk-hedgehog covering over E . Notice $p_1 = q \circ p$ is the maximum disk-hedgehog covering over E_1 . If a loop α in B_1 at b_1 has all lifts to E_1 that are loops, then all lifts of α to E must be loops. Given a lift β in E_2 of $f \circ \alpha$, the map $t \rightarrow (\alpha(t), \beta(t))$ is a lift of α in E . As it is a loop, β must be a loop as well. Consequently, if two loops in B_1 at b_1 are similar, so are their images in B_2 which is sufficient to conclude there is a natural homomorphism from $\pi(B, b_1)$ to $\pi(B_2, f(b_1))$. \square

Proposition 6.3. If $p : E \rightarrow B$ is a regular disk-hedgehog covering and $p(e_0) = b_0$, then one has a natural exact sequence

$$1 \rightarrow \pi(E, e_0) \rightarrow \pi(B, b_0) \rightarrow \pi(p, b_0) \rightarrow 1$$

Proof. Choose a maximal disk-hedgehog covering $p_1 : E_1 \rightarrow E$ over E , where E_1 is a Peano space. Notice $p \circ p_1$ is a maximal disk-hedgehog covering over B .

The kernel of $\pi(B, b_0) \rightarrow \pi(p, b_0)$ consists exactly of loops whose all lifts to E are loops. In particular, the kernel is contained in the image of $\pi(E, e_0) \rightarrow \pi(B, b_0)$. Obviously, the image of $\pi(E, e_0) \rightarrow \pi(B, b_0)$ is contained in that kernel.

Any loop in E at e_0 that becomes trivial in $\pi(B, b_0)$ must have all lifts in E_1 as loops. That means $\pi(E, e_0) \rightarrow \pi(B, b_0)$ is a monomorphism. \square

Theorem 6.4. Suppose $p : E \rightarrow B$ is a disk-hedgehog covering of path connected spaces. Suppose $f : X \rightarrow B$ is a map from a Peano space, $x_0 \in X$ and $e_0 \in E$ with $f(x_0) = b_0 = p(e_0)$. f has a lift $g : (X, x_0) \rightarrow (E, e_0)$ if and only if the image of $\pi(X, x_0) \rightarrow \pi(B, b_0)$ is contained in the image of $\pi(E, e_0) \rightarrow \pi(B, b_0)$.

Proof. Only one implication is of interest, so assume the image of $\pi(X, x_0) \rightarrow \pi(B, b_0)$ is contained in the image of $\pi(E, e_0) \rightarrow \pi(B, b_0)$.

Given a point $x \in X$ pick a path α_x in X from x_0 to x and define $g(x)$ as $\alpha_x \cdot e_0$. $g(x)$ does not depend on the choice of α_x : choosing a different path β_x leads to a loop γ in E at e_0 such that $[\beta_x * \alpha_x^{-1}] = [p \circ \gamma]$ in $\pi(B, b_0)$. Therefore $\beta_x \sim (p \circ \gamma) * \alpha_x$ and $\beta_x \cdot e_0 = ((p \circ \gamma) * \alpha_x) \cdot e_0 = \alpha_x \cdot e_0 = g(x)$.

Given any map $q : H \rightarrow X$ from a disk-hedgehog H to X , the composition $g \circ q : H \rightarrow E$ is the only possible lift of $f \circ q$, hence it is continuous. By 3.3, g is continuous. \square

Corollary 6.5. *Suppose $p : E \rightarrow B$ is a disk-hedgehog covering with E being Peano and $e_0 \in E$. $\pi(E, e_0) = 0$ if and only if $p : E \rightarrow B$ is the maximal disk hedgehog-covering over B .*

Proof. If p is maximal, then E does not admit any non-trivial disk-hedgehog covering and $\pi(E, e_0) = 0$. If $\pi(E, e_0) = 0$, then given any other disk-hedgehog covering $q : E_1 \rightarrow B$ there is a lift $g : E \rightarrow E_1$ of q proving p is maximal. \square

7. COMPARISON TO THE CLASSICAL FUNDAMENTAL GROUP

As the natural homomorphism $\pi_1(B, b_0) \rightarrow \pi(B, b_0)$ is an epimorphism, there are two natural questions:

Problem 7.1. *Characterize the kernel of $\pi_1(B, b_0) \rightarrow \pi(B, b_0)$ for path-connected spaces B .*

Problem 7.2. *Characterize path-connected spaces B such that $\pi_1(B, b_0) \rightarrow \pi(B, b_0)$ is an isomorphism.*

Since the identity map $P(B) \rightarrow B$ from the Peanification of B to B induces isomorphisms of both the classical fundamental group and the hedgehog fundamental group, we will consider both Problems 7.1 and 7.2 for Peano B spaces only. In particular, we differ with [10] in that regard.

Recall B is **shape injective** if the natural homomorphism $\pi_1(B, b_0) \rightarrow \tilde{\pi}_1(B, b_0)$ from the classical fundamental group to the Čech fundamental group is a monomorphism. Papers [11], [7, Corollary 1.2 and Final Remark], [6], and [9] contain results that various classes of spaces are shape injective. We will generalize the concept of shape injectivity as follows:

Definition 7.3. B is **residually Poincaré** if for every loop α in B that is not null-homotopic there is a map $f : B \rightarrow P$ such that P is a Poincaré space and $f \circ \alpha$ is not null-homotopic.

Proposition 7.4. *If B is residually Poincaré, then $\pi_1(B, b_0) \rightarrow \pi(B, b_0)$ is an isomorphism.*

Proof. Clearly, it is so if B is a Poincaré space as it has the classical universal cover that is simply connected. Given a non-trivial element $[\alpha] \in \pi_1(B, b_0)$ choose $f : B \rightarrow (P, p_0)$ such that $f \circ \alpha$ is not null-homotopic. If α represents the neutral element of $\pi(B, b_0)$, then $[f \circ \alpha]$ is neutral in $\pi(P, p_0) = \pi_1(P, p_0)$, a contradiction. \square

Theorem 7.5. *Suppose \mathcal{U} is an open cover of a paracompact space B consisting of path-connected sets. If, for each $x \in B$, the inclusion $st(x, \mathcal{U}) \rightarrow B$ of the star of \mathcal{U} at x induces the trivial homomorphism of $\pi(st(x, \mathcal{U}), x) \rightarrow \pi(B, x)$, then $\pi(B, b_0)$ is isomorphic to the fundamental group of the nerve of \mathcal{U} for all $b_0 \in B$.*

Proof. Pick $V_0 \in \mathcal{U}$ containing b_0 . For each $V \in \mathcal{U}$ pick $b_V \in V$ ($b_V = b_0$ if $V = V_0$).

Define a map α from the 1-skeleton of the nerve $N(\mathcal{U})$ to B as follows: each vertex V of the nerve is mapped to b_V and each edge VW is mapped to a path α_{VW} in $V \cup W$ joining b_V and b_W .

Given an edge-path in the nerve from V_0 to V followed by a loop around a triangle that belongs to the nerve, then followed back by the path-edge results in a loop that is mapped to the star $st(b_V, \mathcal{U})$ of b_V in \mathcal{U} , hence α induces a homomorphism j from $\pi_1(N(\mathcal{U}), V_0)$ to $\pi(B, b_0)$.

Given a loop λ in B at b_0 , we can represent it as the concatenation of paths γ_i , $0 \leq i \leq n$, such that the carrier of γ_i is contained in $V(i) \in \mathcal{U}$, and $V(0) = V_0 = V(n)$. Pick a path ω_i in $V(i)$ joining $\gamma_i(1)$ and $b_{V(i)}$. Notice each γ_i is equivalent to $\omega_{i-1} * \alpha_{V(i-1)V(i)} * \omega_i^{-1}$, so replacing it by that path results in a loop in the image of j that is equivalent to λ . That proves j is an epimorphism.

To show it is a monomorphism, assume there is an edge-loop in the nerve that is mapped to a loop in B being trivial in $\pi(B, b_0)$. Choose a partition of unity $\phi : B \rightarrow N(\mathcal{U})$ sending b_0 to $V_0 \in \mathcal{U}$. The composition of $j : \pi(N(\mathcal{U})) \rightarrow \pi(B, b_0)$ and the homomorphism induced by ϕ is the identity.

Indeed, for each $V \in \mathcal{U}$ choose a path β_V in $N(\mathcal{U})$ from $\phi(b_V)$ to V that lies in the open star $st(V)$ of V in $N(\mathcal{U})$. Notice, if $V \cap W \neq \emptyset$, then $\beta_V * VW * \beta_W^{-1} * (\phi(\alpha_{VW}))^{-1}$ lies in the union $st(V) \cup st(W)$ of open stars in $N(\mathcal{U})$. As their intersection is contractible, the union is simply connected and the composition of $j : \pi(N(\mathcal{U})) \rightarrow \pi(B, b_0)$ and the homomorphism induced by ϕ is the identity. \square

Corollary 7.6. *If B is a paracompact Peano space and $\pi(B, b)$ is discrete for all $b \in B$, then for every sufficiently small open cover \mathcal{U} of B , $\pi(B, b)$ is isomorphic to the fundamental group of the nerve of \mathcal{U} for all $b \in B$.*

Proof. By 3.14 every point $b \in B$ has a path-connected neighborhood U_b such that the maximal disk-hedgehog covering $p : E \rightarrow B$ has a section over U_b . That implies $\pi(U_b, b) \rightarrow \pi(B, b)$ is trivial. Choose a star-refinement \mathcal{V} of $\{U_b\}_{b \in B}$ and apply 7.5 to any refinement \mathcal{U} of \mathcal{V} . \square

Let us show that the analog of the famous result of Shelah [19] (see also [18]) stating that the fundamental group of a Peano continuum is finitely generated if it is countable not only holds for the hedgehog fundamental groups but it also has a much simpler proof.

Corollary 7.7. *Suppose B is a Peano continuum. If $\pi(B, b_0)$ is countable for some $b_0 \in B$, then it is finitely presented.*

Proof. Consider the maximal disk-hedgehog covering $p : E \rightarrow B$. 3.11 says its fibers are Baire spaces. Since they are countable, they must be discrete. Apply 7.6. \square

Let's turn to Problem 7.1. First, let us show that every small loop belongs to the kernel of $\pi_1(B, b_0) \rightarrow \pi(B, b_0)$. It shows that the hedgehog fundamental group eliminates some of the pathologies of the classical fundamental group.

Recall (see [21]) that a loop α at b_0 in B is called **small** if it can be homotoped relative to b_0 into any neighborhood U of b_0 in B .

Proposition 7.8. *Suppose B is path-connected. If $p : E \rightarrow B$ is a disk-hedgehog covering, then $[\alpha]$ is the neutral element of $\pi(p, b_0)$ for every small loop α at b_0 .*

Proof. We may assume E is Peano by switching to its Peanification. Suppose α is a small loop at b_0 in B so that $[\alpha]$ is not the neutral element of $\pi(p, b_0)$. There is a lift $\tilde{\alpha}$ of α with $e_0 = \tilde{\alpha}(0) \neq e_1 = \tilde{\alpha}(1)$.

Choose a path-connected neighborhood U of b_0 in B such that the path component V of e_0 in $p^{-1}(U)$ is different from path-component W of e_1 in $p^{-1}(U)$. Suppose there is a loop β in U homotopic to α rel. b_0 in B . Its lift $\tilde{\beta}$ would join e_0 and e_1 , a contradiction. \square

Let's consider a more general question than 7.1: Characterize kernels of $\pi_1(B, b_0) \rightarrow \pi(p, b_0)$, where $p : E \rightarrow B$ is a disk hedgehog covering over a Peano space B .

As in [20, p.81], given an open cover \mathcal{U} of X , $\pi(\mathcal{U}, x_0)$ is the subgroup of $\pi_1(X, x_0)$ generated by elements of the form $[\alpha * \gamma * \alpha^{-1}]$, where γ is a loop in some $U \in \mathcal{U}$ and α is a path from x_0 to $\gamma(0)$.

Lemma 7.9. *Suppose $P(B, b_0, \sim)$ has a whisker topology such that $\alpha \sim \text{const}$ implies $[\alpha] \in \pi(\mathcal{U}, b_0)$ for some open cover \mathcal{U} of B . If $\beta(t)$, $t \in [0, 1]$, are paths in $P(B, b_0, \sim)$ forming a lift of a path γ starting at $[\alpha]$, then $\beta(t) * \gamma_t^{-1} * \alpha^{-1} \in \pi(\mathcal{U}, b_0)$ for all $t \in [0, 1]$.*

Proof. Let $S = \{t \in [0, 1] \mid \beta(t) * \gamma_t^{-1} * \alpha^{-1} \in \pi(\mathcal{U}, b_0)\}$. Clearly, $0 \in S$. It suffices to show that for any $t \in S$, $t < 1$, there is $s > t$ such that $[t, s] \subset S$ and that S contains its supremum. Given $t \in S$, $t < 1$, pick $V \in \mathcal{U}$ containing $\gamma(t)$ and choose a closed interval $W = [t, u]$ in $[0, 1]$, $u > t$, such that $\beta(s) \in N(\beta(s), V)$ for $s \in W$ and $\gamma(s) \in V$ for $s \in W$. Therefore, given $s \in W$, there is a path ω in V satisfying $\beta(s) \sim \beta(t) * \omega$. Notice ω joins $\gamma(t)$ and $\gamma(s)$.

The loop $\lambda = \omega * (\gamma|_{[t, s]})^{-1}$ lies in V and $\beta(s) * \gamma_s^{-1} * \alpha^{-1} \sim \beta(t) * \omega * \gamma_s^{-1} * \alpha^{-1} \sim \beta(t) * \lambda * (\gamma|_{[t, s]}) * \gamma_s^{-1} * \alpha^{-1} \sim \beta(t) * \lambda * \gamma_t^{-1} * \alpha^{-1} \sim (\beta(t) * \gamma_t^{-1} * \alpha^{-1}) * \alpha * \gamma_t * \lambda * \gamma_t^{-1} * \alpha^{-1}$ and the last loop belongs to $\pi(\mathcal{U}, b_0)$.

The same argument proves that the supremum of S belongs to S (we only used that s and t are sufficiently close). \square

Proposition 7.10. *Let B be a Peano space. If $p : E \rightarrow B$ is a covering projection, then the kernel of $\pi_1(B, b_0) \rightarrow \pi(p, b_0)$ contains $\pi(\mathcal{U}, b_0)$, where \mathcal{U} consists of all open subsets U of B that are evenly covered.*

Given an open cover \mathcal{U} of B , the set of covering projections $q : E \rightarrow B$ for which each $U \in \mathcal{U}$ is evenly covered has a maximum p and the kernel of $\pi_1(B, b_0) \rightarrow \pi(p, b_0)$ is exactly $\pi(\mathcal{U}, b_0)$.

Proof. Obviously, elements of the form $[\alpha * \gamma * \alpha^{-1}]$, where γ is a loop in some $U \in \mathcal{U}$ and α is a path from b_0 to $\gamma(0)$ in E that is a loop, so they are trivial in $\pi(p, b_0)$.

Consider the end-point projection $p : P(B, b_0, \sim) \rightarrow B$ ($\alpha \sim \beta$ if and only if $[\alpha * \beta^{-1}] \in \pi(\mathcal{U}, b_0)$). It is a classical covering with each member of \mathcal{U} being evenly covered (see [3] or use 7.9 to deduce it has unique path-lifting property and then construct sections over members of \mathcal{U}). Notice the kernel of $\pi_1(B, b_0) \rightarrow \pi(p, b_0)$ is exactly $\pi(\mathcal{U}, b_0)$. Indeed, if a loop γ in B lifts to a loop in $P(B, b_0, \sim)$, then 7.9 says the loop must belong to $\pi(\mathcal{U}, b_0)$.

Given any classical covering projection $q : E \rightarrow B$ with each member of \mathcal{U} being evenly covered one can construct $f : P(B, b_0, \sim) \rightarrow E$ such that $q \circ f = p$ by lifting paths. That proves maximality of p . \square

Definition 7.11. The intersection of all $\pi(\mathcal{U}, b_0)$, \mathcal{U} ranging over all open covers of B , is called the **Spanier group** of (B, b_0) (see [10]).

By a **medium loop** we mean a loop α at b_0 that is not small and its homotopy class $[\alpha]$ belongs to the Spanier group. By a **big loop** we mean a loop α at b_0 that is neither medium nor small.

Proposition 7.12. *Let B be a Peano space. If p is the supremum of all classical coverings over B , then the kernel of $\pi_1(B, b_0) \rightarrow \pi(p, b_0)$ is exactly the Spanier group.*

Proof. Consider the end-point projection $p : P(B, b_0, \sim) \rightarrow B$ ($\alpha \sim \beta$ if and only if $[\alpha * \beta^{-1}] \in \pi(\mathcal{U}, b_0)$ for all open covers \mathcal{U} of B). Use 7.9 to deduce it has unique path-lifting property and then use 4.2 to show it is a disk-hedgehog covering. Notice the kernel of $\pi_1(B, b_0) \rightarrow \pi(p, b_0)$ is exactly the Spanier group. Indeed, if a loop γ in B lifts to a loop in $P(B, b_0, \sim)$, then 7.9 says the loop must belong to $\pi(\mathcal{U}, b_0)$ for all open covers \mathcal{U} of B .

Given any classical covering projection $q : E \rightarrow B$ with each member of \mathcal{U} being evenly covered one can construct $f : P(B, b_0, \sim) \rightarrow E$ such that $q \circ f = p$ by lifting paths.

Suppose $q : E \rightarrow B$, E Peano, is a disk-hedgehog covering with $q(e) = b_0$ and maps $f_{\mathcal{U}} : E \rightarrow P(B, b_0, \sim_{\mathcal{U}})$ such that $f_{\mathcal{U}} \circ p_{\mathcal{U}} = q$ for each open cover \mathcal{U} of B . Here $\alpha \sim_{\mathcal{U}} \beta$ if and only if $[\alpha * \beta^{-1}] \in \pi(\mathcal{U}, b_0)$ and $p_{\mathcal{U}}$ is the end-point projection.

Given $x \in E$ and two paths α_x, β_x from e to x , the loop $\alpha_x * \beta_x^{-1}$ must belong to the Spanier group as it can be factored through all $P(B, b_0, \sim_{\mathcal{U}})$, therefore the function $f(x) = [\gamma \alpha_x]$ (γ a fixed loop at b_0 in B) is well-defined and is continuous as p is an arc-hedgehog covering. As $p \circ f = q$, $q \geq p$. That proves maximality of p . \square

Corollary 7.13. *The kernel of $\pi_1(B, b_0) \rightarrow \pi(B, b_0)$ contains all small loops and is contained in the union of small loops and medium loops.*

Let us show how direct wedge can be used to construct interesting spaces.

First of all, one can change the topology of the standard arc-hedgehog $\bigvee_{n \in \mathbb{N}} (I_n, 0_n)$ by requiring open neighborhoods of the base-point to contain all but finitely many 0_n 's (instead of all but finitely many I_n 's) and get a connected space that is not locally connected (a modified topologist's sine curve).

Second, one can change the topology of the standard disk-hedgehog $\bigvee_{n \in \mathbb{N}} (D_n^2, 0_n)$ by requiring open neighborhoods of the base-point to contain all but finitely many ∂D_n^2 's (instead of all but finitely many D_n^2 's) and get a space with properties similar to Harmonic Archipelago [2]: every loop is small.

It is easy to construct examples of medium loops by connecting two Harmonic Archipelagos by an arc. However, there is a more interesting example of Fischer-Zastrow [12] that can be used for that purpose. What is not clear is if that example does not become trivial once we kill all small loops.

Problem 7.14. *Construct a medium loop in a Peano space that does not belong to the normalizer of all small loops.*

REFERENCES

- [1] V. Berestovskii, C. Plaut, *Uniform universal covers of uniform spaces*, Topology Appl. 154 (2007), 1748–1777.
- [2] W.A.Bogley, A.J.Sieradski, *Universal path spaces*, <http://oregonstate.edu/~bogleyw/#research>

- [3] N.Brodskiy, J.Dydak, B.Labuz, A.Mitra, *Covering maps for locally path-connected spaces*, <http://front.math.ucdavis.edu/0801.4967>
- [4] J.W. Cannon, G.R. Conner, *On the fundamental groups of one-dimensional spaces*, *Topology and its Applications* 153 (2006), 2648–2672.
- [5] J. Dydak and J. Segal, *Shape theory: An introduction*, *Lecture Notes in Math.* 688, 1–150, Springer Verlag 1978.
- [6] K. Eda, *The fundamental groups of one-dimensional spaces and spatial homomorphisms*, *Topology and Its Applications*, 123 (2002) 479–505.
- [7] K. Eda and K. Kawamura, *The fundamental group of one-dimensional spaces*, *Topology and Its Applications*, 87 (1998) 163–172.
- [8] P.Fabel, *Metric spaces with discrete topological fundamental group*, *Topology and its Applications* 154 (2007), 635–638.
- [9] H. Fischer and C.R. Guilbault, *On the fundamental groups of trees of manifolds*, *Pacific Journal of Mathematics* 221 (2005) 49–79.
- [10] H.Fischer, D.Repovš, Z.Virk, A.Zastrow *On semilocally simply connected spaces*, *Topology and its Applications* 158(2011), 397–408
- [11] H. Fischer, A. Zastrow, *The fundamental groups of subsets of closed surfaces inject into their first shape groups*, *Algebraic and Geometric Topology* 5 (2005) 1655–1676.
- [12] H.Fischer, A.Zastrow, *Generalized universal coverings and the shape group*, *Fundamenta Mathematicae* 197 (2007), 167–196.
- [13] P.J. Hilton, S. Wylie, *Homology theory: An introduction to algebraic topology*, Cambridge University Press, New York 1960 xv+484 pp.
- [14] Sze-Tsen Hu, *Homotopy theory*, Academic Press, New York and London, 1959.
- [15] E. L. Lima, *Fundamental groups and covering spaces*, AK Peters, Natick, Massachusetts, 2003.
- [16] S. Mardešić and J. Segal, *Shape theory*, North-Holland Publ.Co., Amsterdam 1982.
- [17] J. R. Munkres, *Topology*, Prentice Hall, Upper Saddle River, NJ 2000.
- [18] J.Pawlikowski, *The fundamental group of a compact metric space*, *Proceedings of the American Mathematical Society*, 126 (1998), 3083–3087.
- [19] S. Shelah, *Can the fundamental group of a nice space be e.g. the rationals*, *Abstracts Amer.Math. Soc.* 5 (1984), 217.
- [20] E. Spanier, *Algebraic topology*, McGraw-Hill, New York 1966.
- [21] Z. Virk, *Homotopical Smallness and Closeness*, *Topology and its Applications* 158(2011), 360–378

UNIVERSITY OF TENNESSEE, KNOXVILLE, TN 37996
E-mail address: dydak@math.utk.edu