

Let us consider the weighted graph $(G_{X, \tilde{r}}, \rho)$ with the vertex set $V(G_{X, \tilde{r}}) = \Omega_{\infty, \tilde{r}}^X$, the edge set $E(G_{X, \tilde{r}})$ such that

$$(\{u, v\} \in E(G_{X, \tilde{r}})) \Leftrightarrow (u \text{ and } v \text{ are mutually stable and } u \neq v),$$

and the weight $\rho: E(G_{X, \tilde{r}}) \rightarrow \mathbb{R}^+$ defined by formula (1).

Definition 1. The *pretangent spaces* (to (X, d) at infinity w.r.t. \tilde{r}) are the maximal cliques $\Omega_{\infty, \tilde{r}}^X$ of $G_{X, \tilde{r}}$ with metrics determined with the help of (1).

Recall that a *clique* in a graph G is a set $A \subseteq V(G)$ such that every two distinct points of A are adjacent. A clique C in G is maximal if $C \subseteq A$ implies $C = A$ for every clique A in G .

Define a subset $\alpha_0 = \alpha_0(X, \tilde{r})$ of the set $\tilde{\mathbf{X}}_{\infty, \tilde{r}}$ by the rule

$$(\tilde{z} \in \alpha_0) \Leftrightarrow (\tilde{z} \in \tilde{\mathbf{X}}_{\infty, \tilde{r}} \text{ and } \tilde{d}_{\tilde{r}}(\tilde{z}) = 0). \tag{2}$$

Then α_0 is a common point of all pretangent spaces $\Omega_{\infty, \tilde{r}}^X$. This means, in particular, that the graph $G_{X, \tilde{r}}$ is connected.

Let $(n_k)_{k \in \mathbb{N}} \subset \mathbb{N}$ be infinite and strictly increasing. Denote, by \tilde{r}' , the subsequence $(r_{n_k})_{k \in \mathbb{N}}$ of the $\tilde{r} = (r_n)_{n \in \mathbb{N}}$ and, for every $\tilde{x} = (x_n)_{n \in \mathbb{N}} \in \tilde{\mathbf{X}}_{\infty, \tilde{r}}$, write $\tilde{x}' := (x_{n_k})_{k \in \mathbb{N}}$. It is clear that $\lim_{k \rightarrow \infty} d(x_{n_k}, p) = \infty$ and $\tilde{d}_{\tilde{r}'}(\tilde{x}') = \tilde{d}_{\tilde{r}}(\tilde{x})$ for every $\tilde{x} \in \tilde{\mathbf{X}}_{\infty, \tilde{r}}$. Moreover, if $\tilde{y} \in \tilde{\mathbf{X}}_{\infty, \tilde{r}}$ and $\lim_{n \rightarrow \infty} \frac{d(x_n, y_n)}{r_n}$ exists, then

$$\lim_{k \rightarrow \infty} \frac{d(x_{n_k}, y_{n_k})}{r_{n_k}} = \lim_{n \rightarrow \infty} \frac{d(x_n, y_n)}{r_n}. \tag{3}$$

Let us define $\tilde{\mathbf{X}}_{\infty, \tilde{r}'}$ and ρ' similarly to $\tilde{\mathbf{X}}_{\infty, \tilde{r}}$ and, respectively, to ρ , and let $\pi: \tilde{\mathbf{X}}_{\infty, \tilde{r}} \rightarrow \Omega_{\infty, \tilde{r}}^X$, $\pi': \tilde{\mathbf{X}}_{\infty, \tilde{r}'} \rightarrow \Omega_{\infty, \tilde{r}'}^X$ be the natural projections $\pi(x) := \{\tilde{y} \in \tilde{\mathbf{X}}_{\infty, \tilde{r}} : \rho(\tilde{x}, \tilde{y}) = 0\}$, $\pi'(x) := \{\tilde{y} \in \tilde{\mathbf{X}}_{\infty, \tilde{r}'} : \rho'(\tilde{x}, \tilde{y}) = 0\}$, and let $\varphi_{\tilde{r}'}(\tilde{x}) := \tilde{x}'$ for all $\tilde{x} \in \tilde{\mathbf{X}}_{\infty, \tilde{r}}$. Then there is an embedding $em': \Omega_{\infty, \tilde{r}}^X \rightarrow \Omega_{\infty, \tilde{r}'}^X$ of the weighted graph $(G_{X, \tilde{r}}, \rho)$ in the weighted graph $(G_{X, \tilde{r}'}, \rho')$ such that the diagram

$$\begin{array}{ccc} \tilde{\mathbf{X}}_{\infty, \tilde{r}} & \xrightarrow{\varphi_{\tilde{r}'}} & \tilde{\mathbf{X}}_{\infty, \tilde{r}'} \\ \pi \downarrow & & \downarrow \pi' \\ \Omega_{\infty, \tilde{r}}^X & \xrightarrow{em'} & \Omega_{\infty, \tilde{r}'}^X \end{array}$$

is commutative. Since em' is an embedding of weighted graphs, $em'(C)$ is a clique in $G_{X, \tilde{r}'}$ if C is a clique in $G_{X, \tilde{r}}$. Furthermore, (3) implies that the restrictions $em'|_{\Omega_{\infty, \tilde{r}}^X}$ are isometries of the pretangent spaces $\Omega_{\infty, \tilde{r}}^X$ on the metric spaces $em'(\Omega_{\infty, \tilde{r}}^X)$.

Definition 2. A pretangent space $\Omega_{\infty, \tilde{r}}^X$ is *tangent* if the clique $em'(\Omega_{\infty, \tilde{r}}^X)$ is maximal for every infinite, strictly increasing sequence $(n_k)_{k \in \mathbb{N}} \subset \mathbb{N}$.

Example 1. Let E be a finite-dimensional Euclidean space and let $X \subseteq E$ be such that the Hausdorff distance $d_H(E, X)$ is finite. Then, for every scaling sequence \tilde{r} , all pretangent spaces $\Omega_{\infty, \tilde{r}}^X$ are tangent and isometric to E .

In conclusion of this brief introduction, it should be noted that there exist other techniques, which allow one to investigate the asymptotic properties of metric spaces at infinity. As examples, we mention only the ballean theory [2] and the Wijsman convergence [3–5].

1. Finiteness. In this section, we study the conditions, under which pretangent spaces are finite.

Theorem 1. Let (X, d) be an unbounded metric space, $p \in X$, $n \geq 2$, and let

$$F_n(x_1, \dots, x_n) := \begin{cases} \frac{\min_{1 \leq k \leq n} d(x_k, p) \prod_{1 \leq k < l \leq n} d(x_k, x_l)}{\left(\max_{1 \leq k \leq n} d(x_k, p) \right)^{\frac{n(n-1)}{2} + 1}}, & \text{if } (x_1, \dots, x_n) \neq (p, \dots, p), \\ 0, & \text{if } (x_1, \dots, x_n) = (p, \dots, p). \end{cases}$$

Then the inequality $|\Omega_{\infty, \tilde{r}}^X| \leq n$ holds for every pretangent space $\Omega_{\infty, \tilde{r}}^X$ if and only if $\lim_{x_1, \dots, x_n \rightarrow \infty} F_n(x_1, \dots, x_n) = 0$.

Note that, for every unbounded metric space (X, d) , there is a pretangent space $\Omega_{\infty, \tilde{r}}^X$ consisting of at least two points.

Corollary 1. Let (X, d) be an unbounded metric space and let α_0 be a point defined by (2) for every scaling sequence \tilde{r} . Then the following statements are equivalent:

- (i) The graph $G_{X, \tilde{r}}$ is a star with the center α_0 for every scaling sequence \tilde{r} ;
- (ii) The limit relation $\lim_{x_1, x_2 \rightarrow \infty} F_2(x_1, x_2) = 0$ holds.

Let us consider now the problem of existence of finite tangent spaces.

Definition 3. Let $E \subseteq \mathbb{R}^+$. The porosity of E at infinity is the quantity

$$p(E, \infty) := \limsup_{h \rightarrow \infty} \frac{l(\infty, h, E)}{h} \tag{4}$$

where $l(\infty, h, E)$ is the length of the longest interval in the set $[0, h] \setminus E$. The set E is strongly porous at infinity if $p(E, \infty) = 1$.

The standard definition of the porosity at a point can be found in [6].

For a metric space (X, d) and $p \in X$, we write $S_p(X) := \{d(x, p) : x \in X\}$.

Theorem 2. Let (X, d) be an unbounded metric space, $p \in X$. The following statements are equivalent:

- (i) The set $S_p(X)$ is strongly porous at infinity;
- (ii) There is a single-point tangent space $\Omega_{\infty, \tilde{r}}^X$;
- (iii) There is a finite tangent space $\Omega_{\infty, \tilde{r}}^X$;
- (iv) There is a compact tangent space $\Omega_{\infty, \tilde{r}}^X$;
- (v) There is a bounded, separable tangent space $\Omega_{\infty, \tilde{r}}^X$.

Some results similar to Theorem 1 and Theorem 2 can be found in [7] and [8] respectively.

2. Boundedness. Let $\tilde{\tau} = (\tau_n)_{n \in \mathbb{N}} \subset \mathbb{R}$. We say that $\tilde{\tau}$ is eventually increasing if the inequality $\tau_{n+1} \geq \tau_n$ holds for sufficiently large n . For $E \subseteq \mathbb{R}^+$, we write \tilde{E}_∞^i for the set of eventually

increasing sequences $\tilde{\tau} \subset E$ with $\lim_{n \rightarrow \infty} \tau_n = \infty$. Denote also, by \tilde{I}_E^i , the set of all sequences of open intervals $(a_n, b_n) \subseteq \mathbb{R}^+$ meeting the following conditions:

- Each (a_n, b_n) is a connected component of the set $\text{Int}(\mathbb{R}^+ \setminus E)$;
- $(a_n)_{n \in \mathbb{N}}$ is eventually increasing;

$$\lim_{n \rightarrow \infty} a_n = \infty \text{ and } \lim_{n \rightarrow \infty} \frac{b_n - a_n}{b_n} = 1.$$

Define an equivalence \asymp on the set of sequences of strictly positive numbers as follows. Let $\tilde{a} = (a_n)_{n \in \mathbb{N}}$ and $\tilde{\gamma} = (\gamma_n)_{n \in \mathbb{N}}$. Then $\tilde{a} \asymp \tilde{\gamma}$ if there are some constants $c_1, c_2 > 0$ such that $c_1 a_n < \gamma_n < c_2 a_n$ for every $n \in \mathbb{N}$.

Definition 4. Let $E \subseteq \mathbb{R}^+$ and let $\tilde{\tau} \in \tilde{E}_\infty^i$. The set E is $\tilde{\tau}$ -strongly porous at infinity if there is a sequence $((a_n, b_n))_{n \in \mathbb{N}} \in \tilde{I}_E^i$ such that $\tilde{\tau} \asymp \tilde{a}$ where $\tilde{a} = (a_n)_{n \in \mathbb{N}}$. The set E is completely strongly porous at infinity if E is $\tilde{\tau}$ -strongly porous at infinity for every $\tilde{\tau} \in \tilde{E}_\infty^i$.

Note that every completely strongly porous at infinity set is strongly porous at infinity but not conversely.

Definition 5. Let (X, d) be an unbounded metric space and let $p \in X$. A scaling sequence \tilde{r} is normal if \tilde{r} is eventually increasing and there is $\tilde{x} \in \tilde{X}_{\infty, \tilde{r}}$ such that

$$\lim_{n \rightarrow \infty} \frac{d(x_n, p)}{r_n} = 1.$$

Write $\mathfrak{S}_n(X)$ for the set of all pretangent spaces $\Omega_{\infty, \tilde{r}}^X$ with normal scaling sequences \tilde{r} . Under which conditions is the family $\mathfrak{S}_n(X)$ uniformly bounded?

Recall that a family \mathfrak{S} of a metric spaces (Y, d_Y) is uniformly bounded if

$$\sup\{\text{diam} Y : Y \in \mathfrak{S}\} < \infty$$

If all metric spaces (Y, d_Y) are pointed with marked points $p_Y \in Y$ and

$$\inf\{d_Y(p_Y, y) : y \in Y \setminus \{p_Y\}, Y \in \mathfrak{S}\} > 0,$$

then we say that \mathfrak{S} is uniformly discrete (w.r.t. the points p_Y).

The following theorem is an analog of Theorem 3.11 and Theorem 4.1 from [9].

Theorem 3. Let (X, d) be an unbounded metric space and let $p \in X$. Then the following statements are equivalent:

- (i) The family $\mathfrak{S}_n(X)$ is uniformly bounded;
- (ii) $\mathcal{S}_p(X)$ is completely strongly porous at infinity;
- (iii) The family $\mathfrak{S}_n(X)$ is uniformly discrete w.r.t. the points α_0 defined by (2).

If $\mathfrak{S}_n(X)$ is uniformly bounded, then every pretangent space $\Omega_{\infty, \tilde{r}}^X$ is bounded, but the converse, in general, does not hold.

Definition 6. The set $E \subseteq \mathbb{R}^+$ is w -strongly porous at infinity if, for every sequence $\tilde{\tau} \in \tilde{E}_\infty^i$, there is a subsequence $\tilde{\tau}'$, for which E is $\tilde{\tau}'$ -strongly porous at infinity.

The following theorem gives a boundedness criterion for pretangent spaces.

Theorem 4. Let (X, d) be an unbounded metric space and let $p \in X$. All pretangent spaces to X at infinity are bounded if and only if the set $\mathcal{S}_p(X)$ is w -strongly porous at infinity.

The example of ω -strongly porous at infinity set $E \subseteq \mathbb{R}^+$ which is not completely strongly porous at infinity, can be obtained as a modification of Example 2.10 [10].

Using Theorem 4, we can obtain a criterion of existence of an unbounded pretangent space. The conditions of such type are important for the development of a theory of pretangent spaces of the second order or more (i.e., pretangent spaces to pretangent spaces, pretangent spaces to pretangent spaces to pretangent spaces and so on).

Let (Y, d_Y) be a metric space. Then for every $K \subseteq Y$ and $y \in Y$, we write $\text{dist}(y, K) = \inf \{d_Y(y, x) : x \in K\}$. The following definition can be found in [11].

Definition 7. Let $(K_n)_{n \in \mathbb{N}}$ be a sequence of subsets of (Y, d_Y) . The set

$$\text{Lim inf}_{n \rightarrow \infty} K_n := \{y \in Y : \lim_{n \rightarrow \infty} \text{dist}(y, K_n) = 0\}$$

is the Kuratowski lower limit of $(K_n)_{n \in \mathbb{N}}$ in (Y, d_Y) .

For $A \subseteq \mathbb{R}$ and $t \in \mathbb{R}$, we set $tA := \{ta : a \in A\}$.

Theorem 5. Let (X, d) be an unbounded metric space and let $p \in X$. Then the following statements are equivalent:

(i) There exists an unbounded pretangent space $\Omega_{\infty, \tilde{r}}^X$;

(ii) There exists a scaling sequence $\tilde{r} = (r_n)_{n \in \mathbb{N}}$ such that the Kuratowski lower limit

$$\text{Lim inf}_{n \rightarrow \infty} \frac{1}{r_n} S_p(X)$$

is an unbounded subset of \mathbb{R}^+ ;

(iii) The set $S_p(X)$ is not ω -strongly porous at infinity.

Let $(\tilde{r}_m)_{m \in \mathbb{N}}$ be a sequence of scaling sequences. Then for every $m \in \mathbb{N}$ and every unbounded metric space (X, d) we define a pretangent space $\Omega_{\infty, (\tilde{r}_1, \dots, \tilde{r}_m)}^X$ by the following inductive rule: $\Omega_{\infty, (\tilde{r}_1)}^X := \Omega_{\infty, \tilde{r}_1}^X$ if $m = 1$ and

$$\Omega_{\infty, (\tilde{r}_1, \dots, \tilde{r}_m)}^X := \Omega_{\infty, \tilde{r}_m}^X(\Omega_{\infty, (\tilde{r}_1, \dots, \tilde{r}_{m-1})}^X)$$

if $m \geq 2$.

Using Theorem 5, we can obtain the following corollary.

Corollary 2. Let (X, d) be an unbounded metric space and let $p \in X$. If the equality $p(S_p(X), \infty) = 0$ holds, then there is a sequence $(\tilde{r}_m)_{m \in \mathbb{N}}$ of scaling sequences such that the pretangent space $\Omega_{\infty, (\tilde{r}_1, \dots, \tilde{r}_m)}^X$ is unbounded for every $m \in \mathbb{N}$.

The research was supported by the grant of the State Fund for Fundamental Research (project F71/20570).

REFERENCES

1. Roe, J. (2003). Lectures on coarse geometry. University Lecture Series 31. Providence, RI: American Mathematical Society.
2. Protasov, I. & Zarichnyi, M. (2007). General Asymptology. Math. Studies Monograph Series, Vol. 12. Lviv: VNTL Publishers.
3. Lechiki, A. & Levi, S. (1987). Wijsmann convergence in the hyperspace of a metric space. Bull. Unione Mat. Ital. 1-B, pp. 439-452.
4. Wijsman, R. A. (1964). Convergence of sequences of convex sets, cones and functions. Bull. Amer. Math. Soc. 70, pp. 186-188.

5. Wijsman, R. A. (1966). Convergence of sequences of convex sets, cones and functions II. Trans. Amer. Math. Soc., 123, No 1, pp. 32-45.
6. Thomson, B. S. (1985). Real Functions. Lecture Notes in Mathematics, 1170. Berlin etc.: Springer.
7. Abdullayev, F., Dovgoshey, O. & Küçükaslan, M. (2010). Compactness and boundedness of tangent spaces to metric spaces. Beitr. Algebra Geom., 51, No 2, pp. 547-576.
8. Dovgoshey, O. & Martio, O. (2011). Tangent spaces to general metric spaces. Rev. Roumaine Math. Pures. Appl., 56, No 2, pp. 137-155.
9. Bilet, V., Dovgoshey, O. & Küçükaslan, M. (2013). Uniform boundedness of pretangent spaces, local constancy of metric derivatives and strong right upper porosity at a point. J. Anal., 21, pp. 31-55.
10. Bilet, V. & Dovgoshey, O. (2014). Boundedness of pretangent spaces to general metric spaces. Ann. Acad. Sci. Fenn., 39, No 4, pp. 73-82.
11. Aubin, J.-P. & Frankowska, H. (1990). Set-Valued Analysis. Boston, Basel, Berlin: Birkhäuser.

Received 11.04.2017

V.V. Билет, О.А. Довгошей

Институт прикладной математики и механики НАН Украины, Слов'янськ
E-mail: viktoriibilet@gmail.com, oleksiy.dovgoshey@gmail.com

АСИМПТОТИЧНА ПОВЕДІНКА МЕТРИЧНИХ ПРОСТОРІВ НА НЕСКІНЧЕННОСТІ

Запропоновано новий секвенціальний підхід до дослідження структури метричних просторів у нескінченно віддаленій точці. Знайдено критерії скінченності та обмеженості метричних просторів на нескінченності.

Ключові слова: асимптотична обмеженість метричного простору, асимптотична скінченність метричного простору, збіжність метричних просторів, сильна пористість у точці.

V.V. Bilet, A.A. Dovgohey

Институт прикладной математики и механики НАН Украины, Славянск
E-mail: viktoriibilet@gmail.com, oleksiy.dovgoshey@gmail.com

АСИМПТОТИЧЕСКОЕ ПОВЕДЕНИЕ МЕТРИЧЕСКИХ ПРОСТРАНСТВ НА БЕСКОНЕЧНОСТИ

Предложен новый секвенциальный подход к исследованию структуры метрических пространств в бесконечно удаленной точке. Найден критерии конечности и ограниченности метрических пространств на бесконечности.

Ключевые слова: асимптотическая ограниченность метрического пространства, асимптотическая конечность метрического пространства, сходимість метрических пространств, сильная пористость в точке.