

THE GENERALIZED STRATIGRAPHIC SCHEME OF THE JURASSIC OF WESTERN UKRAINE N.M. Zhabina¹, V.Ye. Shlapinsky², M.G. Prykhodko³, O.V. Anikeyeva⁴, D.V. Machalsky⁵

(Recommended by doctor of geological and mineralogical sciences M.M. Ivanik)

¹ *Institute of Geological Sciences of NAS of Ukraine, Kyiv, Ukraine, E-mail: zhabinanatalia@gmail.com
Doctor of geological sciences, senior research scientist.*

² *Institute of Geology and Geochemistry of Combustible Minerals of NAS of Ukraine, Lviv, Ukraine,
E-mail: vlash.ukr@gmail.com
Candidate of geological sciences, senior research scientist.*

³ *Zahidukrgeologiya DP NAK «Nadra Ukraine», Zakarpatska GRE, Berehovo, Ukraine,
E-mail: geomukpart@mail.ru
Chief of search and shooting party.*

⁴ *Institute of Geology and Geochemistry of Combustible Minerals of NAS of Ukraine, Lviv, Ukraine,
E-mail: geolena@ukr.net
Candidate of geological sciences, senior research scientist.*

⁵ *Scientific and Geological Center DP NAK «Nadra Ukraine» Zahidukrgeologiya, Lviv, Ukraine,
E-mail: dijumast@ukr.net
Candidate of geological sciences, leading geologist.*

In Western Ukraine the Jurassic deposits are known in the Carpathians, Transcarpathian Deep, Carpathian Foredeep and Western part of the East European platform. The Generalized stratigraphic scheme of the Lower, Middle and Upper Jurassic of these regions is represented. The current stratigraphic scheme is supplemented and detailed according to modern data, analysis and re-interpretation of many-years investigations these deposits by different specialists. The characteristics of stratigraphic units: distribution, lithological and paleontological composition, position in section, thicknesses and criteria for dating are given. The most complete and precise sequence of the Jurassic stages is determined on the territory of Carpathian Foredeep and adjacent part of the platform. In other regions these deposits are present fragmentarily. The stratigraphic units are mainly dated by the complexes of macro- and microfossils. In Pre-carpathians predominantly there are palynocomplexes for dating the Lower Jurassic deposits, mollusks, foraminifers and palynocomplexes for Middle Jurassic, and foraminifers, mollusks and tintinnids for Upper Jurassic. In Carpathians there are ammonites and belemnites for dating the Lower and Middle Jurassic deposits, and ammonites, tintinnids, dinocysts for Upper Jurassic. Correlation of the Jurassic stratigraphic units of the Western Ukraine is done.

Key words: stratigraphy, Jurassic, Ukrainian Carpathians, Transcarpathian deep, Carpathian foredeep, East European Platform.

ЗВЕДЕНА СТРАТИГРАФІЧНА СХЕМА ЮРИ ЗАХОДУ УКРАЇНИ

Жабіна Н.М.¹, Шлапінський В.Є.², Приходько М.Г.³, Анікеєва О.В.⁴, Мачальський Д.В.⁴

(Рекомендовано д-ром геол.-мінерал. наук М.М. Іваніком)

¹ *Інститут геологічних наук НАН України, Київ, Україна, E-mail: zhabinanatalia@gmail.com
Доктор геологічних наук, старший науковий співробітник.*

² *Інститут геології і геохімії горючих копалин НАН України, Львів, Україна, E-mail:
vlash.ukr@gmail.com
Кандидат геологічних наук, старший науковий співробітник.*

³ *ДП «Західукргеологія», Закарпатська ГРЕ, Берегове, Закарпатська область, Україна,
E-mail: geomukpart@mail.ru
Начальник пошуково-знімальної партії.*

⁴ *Інститут геології і геохімії горючих копалин НАН України, Львів, Україна, E-mail: geolena@ukr.net
Кандидат геологічних наук, старший науковий співробітник.*

⁵ *Науково-геологічний центр ДП НАК «Надра України» «Западукргеологія», Львів, Україна,
E-mail: dijumast@ukr.net
Кандидат геологічних наук, провідний геолог.*

На Заході України юрські відклади поширені у Карпатах, фундаменті Закарпатського та Передкарпатського прогинів та на західній окраїні Східно-Європейської платформи. Представлено зведену схему стратиграфії нижньо-, середньо- та верхньоярських відкладів Західного регіону України. Чинну стратиграфічну схему доповнено і деталізовано на основі аналізу новітніх даних, узагальнення та переінтерпретації результатів багаторічних досліджень цих відкладів різними спеціалістами. Наведено характеристику стратиграфічних підрозділів: поширення, літологічний та палеонтологічний склад, потужності, стратиграфічне положення та обґрунтування віку. Найбільш повну і чітку послідовність ярусів юрської системи встановлено на території Передкарпатського прогину та прилеглою краю платформи, в інших регіонах ці відклади присутні фрагментарно. Стратиграфічні підрозділи датовані переважно за комплексами макро- та мікрофосилій. У Передкарпатті стратиграфічні підрозділи нижньої юри датовані за палинокомплексами, середньої юри – за моллюсками, форамініферами та палинокомплексами, верхньої юри – за форамініферами, моллюсками, тинтинідами. В Карпатах відклади нижньої та середньої юри датовано переважно за амонітами і белемнітами, верхньої юри – за амонітами, тинтинідами, диноцистами. У фундаменті Закарпатського прогину ярусний поділ на теперішній час не встановлено, виділені товщі датуються на підставі віку моллюсків, радіолярій та тинтинід. Здійснено кореляцію стратиграфічних підрозділів юри в межах Західного регіону України.

Ключові слова: стратиграфія, юра, Українські Карпати, Закарпатський прогин, Передкарпатський прогин, Східно-Європейська платформа.

СВОДНАЯ СТРАТИГРАФИЧЕСКАЯ СХЕМА ЮРЫ ЗАПАДА УКРАИНЫ

Жабина Н.М.¹, Шлапинский В.Е.², Приходько М.Г.³, Аникеева Е.В.⁴, Мачальский Д.В.⁵

(Рекомендовано д-ром геол.-минерал. наук М.М. Иваником)

¹ *Институт геологических наук НАН Украины, Киев, Украина, E-mail: zhabinanatalia@gmail.com
Доктор геологических наук, старший научный сотрудник.*

² *Институт геологии и геохимии горючих ископаемых НАН Украины, Львов, Украина,
E-mail: vlash.ukr@gmail.com
Кандидат геологических наук, старший научный сотрудник.*

³ *ДП «Западукргеология», Закарпатская ГРЭ, Берегово, Закарпатская область, Украина,
E-mail: geomukpart@mail.ru
Начальник поисковосъемочной партии.*

⁴ *Институт геологии и геохимии горючих ископаемых НАН Украины, Львов, Украина,
E-mail: geolena@ukr.net
Кандидат геологических наук, старший научный сотрудник.*

⁵ *Научно-геологический центр ДП НАК «Недра Украины» «Западукргеология», Львов, Украина,
E-mail: dijumast@ukr.net
Кандидат геологических наук, ведущий геолог.*

На Западе Украины юрские отложения распространены в Карпатах, фундаменте Закарпатского и Предкарпатского прогибов, на западной окраине Восточно-Европейской платформы. Представлена сводная схема стратиграфии нижне-, средне- и верхнеюрских отложений Западного региона Украины. Действующая стратиграфическая схема дополнена и детализирована на основании анализа новейших данных, обобщения и переинтерпретации результатов многолетних исследований этих отложений многими специалистами. Приведена характеристика стратиграфических подразделений: распространение, литологический и палеонтологический состав, мощности, стратиграфическое положение и обоснование возраста. Наиболее полная и четкая последовательность ярусов юрской системы установлена на территории Предкарпатского прогиба и прилегающей окраины платформы, в других регионах эти отложения присутствуют фрагментарно. Стратиграфические подразделения датированы преимущественно по комплексам макро- и микрофосиллий. В Предкарпатье стратиграфические подразделения нижней юры датированы по палинокомплексам, средней юры – по моллюскам, фораминиферам и палинокомплексам, верхней юры – по фораминиферам, моллюскам, тинтиннидам. В Карпатах отложения нижней и средней юры датированы преимущественно по аммонитам и белемнитам, верхней юры – по аммонитам, тинтиннидам, диноцистам. В фундаменте Закарпатского прогиба ярусное деление в настоящее время не установлено, выделенные толщи датируются на основании возраста моллюсков и тинтиннид. Проведена корреляция стратиграфических подразделений юры в пределах Западного региона Украины.

Ключевые слова: стратиграфия, юра, Украинские Карпаты, Закарпатский прогиб, Предкарпатский прогиб, Восточно-Европейская платформа.

Introduction

In the West of Ukraine the Jurassic deposits are known in Folded Carpathians Area, basements of Carpathian Foredeep and Transcarpathian Depression and in Western part of the East-European platform. In Carpathians these deposits outcrop in Pieniny Zone, in Marmarosh Massif (Bilopotik Subnapper), in Kamyanopotik and Porkulets Nappers. In Transcarpathian Depression basement, the Jurassic rocks were opened by the wells in the Uzhgorod-Solotvino and Pripannonian zones (Fig. 1) [Жабина и др., 2015]. In the territory of Carpathian Foredeep and adjacent part of the East-European platform (Stryi Jurassic depression) Jurassic deposits outcrop locally in Dniester Valley and were opened by numerous wells.




Fig. 1. Tectonic zonation of the Western Ukraine (after data of Lviv Branch UkrDGRI and DP Zahidukrgeologia) Legend: East-European platform: I-1A – Kovel Ledge, I-1B – Volyno-Podolian Monocline, I-1C – Lviv Paleozoic Deep, I-2 – West-European platform; Carpathian Foredeep: II – Bilche-Volitsa zone, III – Sambir Napper, IV – Borislav-Pokutian; Outer (Flish) Carpathians (nappers): V – Skiby, VI – Krosno, VII – Chornogory, VIII – Dukla, IX – Porculets (Burkut), X – Rakhiv, XI – Magura, XII – Kamyanopotik; Median Carpathians: XIII – Marmarosh Massif (subnappers: b – Bilopotik, d – Delovets), XIV – Vezhany Napper, XV – Monastrets Napper; XVI – Pieniny Zone; Inner Carpathians (Transcarpathian Deep): XVII – Uzhgorod-Solotvino Zone, XVIII – Pripannonian Zone (subzones: a – Berehovo, b – Vyshkiv)

Many eminent geologists were studied geology and stratigraphy of Jurassic deposits in the Western Ukraine – A. Alth, O.S. Vyalov, V.V. Glushko, V.I. Slavin, V.J. Dobrynina, S.S. Kruglov, S.E. Smirnov, P.Yu. Lozyniak, etc. Geology of Carpathians and Transcarpathian Depression was the subject of research of F. Hauer, F. Herbich, L. Szajnocha, L. Loczy, A. Hezell, H. Zapalowicz, V. Uhlig, Z. Sujkowski, V. Spacek, D. Andrusov, A. Matejka, Z. Pazdro, M.G. Lomize, M.R. Ladyzhensky, E.K. Lazarenko, V.Ye. Khain, J.O. Kulchytsky, O.V. Maksimov, M.A. Beyer, A.G. Zhurakovsky, A.A. Voloshin, Z.F. Zhigunova, A.K. Boyko, S.L. Byzova, V.G. Chernov, B.D. Pukach, E.M. Titov, A.V. Zobkov, M.J. Petrashkevich, A.A. Matveyeva, V.A. Danilenko, N.F. Udich, and many others. Geology of Ukrainian Precarpathians were studied by Fr. Bieniasz, K. Glazewski, O.O. Leszczynsky, J.M. Sandler, O.M. Anastasieva, V.M. Utrobin, A.P. Sen, M. Palkin, V.S. Burov, I.B. Vishnyakov, V.M. Markovsky, Yu.K. Ovcharenko, Yu.R. Karpenchuk, R.T. Trushkevich, O.V. Samarska, T.S. Izotova, and many others. As a result of many-years investigations, the various regional stratigraphic schemes were created [Бирюлев и др., 1974; Дулуб и др., 1986; Гаврилишин, 1993; Стратиграфические..., 1993; Дулуб и др., 2003; Жабина, Анিকেева, 2007]. Modernized stratigraphic schemes of the Lower-Middle Jurassic and of the Upper Jurassic of Ukrainian Carpathians [Стратиграфія..., 2014] are the synthesis of mentioned schemes.

This work represents a Generalized correlative scheme of stratigraphy of the Jurassic deposits of Western Ukraine, detailed and supplemented with biostratigraphic and lithostratigraphic data.

Materials and methods

The proposed generalized stratigraphic scheme of the Jurassic of the Western Ukraine is based on the correlation of Jurassic deposits in the regions of Carpathian belt and the western edge of the East European platform. The current stratigraphic scheme [Стратиграфія..., 2014] is supplemented by data about the Jurassic deposits in Kamennopotik and Porkulets Nappers of Outer (Flish) Carpathians and in the Marmarosh Massif according to results of investigations, which were done by V.V. Kuzovenko, V.E. Shlapinsky, M.G. Prykhodko, B.M. Matskiv in Carpathians. The stratigraphy of the Pieniny Zone is detailed on the basis of analysis, comparison and re-interpretation of published data of V.I. Slavin, S.S. Kruglov, V.I. Havrylyshyn, G.J. Krymgolts,

T.D. Kalenichenko, L.V. Linetska, M.A. Beer, A. Wierzbowski, M. Krobicki, B. Matyja, D. Rehakova, J. Schlogl, M. Barski. The age of the stratigraphy units is justified by their detailed paleontological characteristic with the analysis of data of study of various group of fossils by many researchers. There are **ammonites** (V.I. Slavin, N.T. Sazonov, T.D. Kalenichenko, S.S. Kruglov, R.J. Leschukh, G.I. Gotsanyuk, A. Wierzbowski et al., D. Rehakova et al.); **belemnites** (G.J. Krymgolts, V.I. Havrylyshyn); **bivalves** and **brachiopods** (G.T. Pchelintsev; P.A. Gerasimov, R.S. Dmitrieva, B.S. Kokoshinska, O.I. Nikiferova, O.M. Schmidt, V.I. Hawrylyshyn, I.M. Yamnichenko, S.I. Pasternak, R.J. Leschukh, I.V. Shaynoga); **foraminifers** (O.A. Hoffman, L.G. Dain, V.G. Dulub, A.S. Terestchuk, M.T. Chip, V.P. Maslov, N.M. Zhabina); **tintinnids** (L.V. Linetska, N.M. Zhabina, D. Rehakova); **radiolarians** (P.Yu. Lozyniak); **dinocysts** (L.V. Linetska, D. Rehakova); **palynocomplexes** (M.I. Burova, M.E. Ohorodnik); **floristic remnants** (E.M. Migacheva, A.I. Turutanova-Ketova, F.A. Stanislavsky, V.A. Vakhrameev, Yu.V. Teslenko) [Славин, 1963; Калениченко и др., 1965; Геология..., 1966; Круглов и др., 1968; Калениченко, Круглов, 1969; Лінецька, 1972; Лещух, Гоцанюк, 2002, 2003; Гоцанюк, 2004; Rehakova et al., 2011, Wierzbowski et al., 2012; Стратиграфія..., 2014, etc.].

Stratigraphical units

Median and Outer Carpathians. The complex tectonic structure of Carpathians complicates the stratification of Jurassic deposits and creation of stratigraphic scheme. In the Median Carpathians the stratigraphical units were distinguished in Pieniny Zone and Belopotik Subnapper of Marmarosh Massif, and in the Outer Carpathians – in Kamyapotik and Porkulets Nappers (Fig. 2). Jurassic volcanic and sedimentary rocks are also present in the composition of the olistoliths in Lower Cretaceous olistostrome in Vezhany Napper.

Pieniny Zone. There were distinguished Perechyn, Priborzhava, Zhubrakiy, Svalyava suites and four strata – Variegated brecciated limestone, Siliceous limestone, Light-coloured limestone and Red-coloured limestone. There are the tectonic blocks or olistoliths of Jurassic rocks lying as detached bodies in Cretaceous thicknesses.

Perechyn suite (V.I. Slavin, V.Ya. Dobrynina, N.A. Yefimova, 1967) [Славин и др., 1967] – Hettangian–Lower Pliensbachian. Pieniny Zone. Tectonic blocks, sometimes olistoliths in Upper Cretaceous matrix. Two subsuites: lower (thickness




Fig. 2. The Generalized stratigraphic scheme of the Jurassic of the Western Ukraine

20-30 m) – mudstones and sandstones with interbedded limestones (Lower Hettangian–Sinemurian) and upper (30 m) – interbedding of marls, limestones and clayey shales with large nodules of cherts (Upper Sinemurian–Lower Pliensbachian). The lower and upper boundaries are unknown. Suite is dated by ammonites *Arietites bucklandi* Sow., *Paltechioceras nodotianum* (Orb.), *Vermiceras nodotianum* Orb., *Schlotheimia charmassei* Orb., *Echioceras pohioides* Fuc., *E. declivis* Truen., *E. raricostatum* Bayeux, belemnites *Salpingoteuthis tubularis* Young et Bird., *Passaloteuthis apicicurvata* Bl., *P. paxillosus* Schloth., *Nannobelus acutus* Miller, bivalves *Gryphaea arcuata* Linne, *G. orbiculata* Sow., *G. incurva* Sow., *Cardinia hybrida* Stutch., *C. listeri* Sow., brachiopods *Spiriferina alpina* Opp., etc.

Priborzhava suite (V.I. Slavin, V.Ya. Dobrynina, N.A. Yefimova, 1967) [Славин и др., 1967] – Upper Pliensbachian–Aalenian. Pieniny Zone. Tectonic blocks, somewhere olistoliths in Upper Cretaceous matrix. Dark gray and variegated mudstones and siltstones interbedded with sandy limestones with glauconite and marls. The base is not found; suite is overlapped unconformably by Zhubrakov suite. Thickness is to 60 m. Suite is dated by ammonites *Uptonia jamesoni* (Sow.), *Androgynoceras capricornus* Schlot., *Harpoceras falcifer* Sow., *Grammoceras thouarsense* d'Orb., *G. saemanni* Dumor., *Dactylioceras commune* Sow., *Leioceras opalinum* Reinecke, *L. acutum* Quenst., *Ludwigia murchisonae* Sow., *L. literata* Buck., *L. carinata* Buck., *Lytoceras ophioneum* Benes., *Lioceras costosum* Quenst. and belemnites *Passaloteuthis ima* Long., *P. bruguieri* Orb., *P. aplicicurvata* Bl., *Hastites chormonthensis* May., *Mesoteuthis triscissa* Gon., *M. tripartita* Schlot., *M. oxycona* Hell., *M. conoidea* Opp., *M. pyramidalis* Liet., *M. banzensis* Kolb, *M. stimula* Dun., *M. qenstedti* Opperl, *M. inornata* Philips, *Rhabdobelus exilis* Orb., etc.

Zhubrakov suite (V.I. Slavin, 1956) [Славин, 1956] – Bajosian–Callovian. Pieniny Zone. Tectonic blocks, sometimes olistoliths in Upper Cretaceous matrix. Gray, green, pink crystal or sometimes crinoid limestones often with cherts. Suite overlies unconformably Priborzhava suite, and is overlain by Svalyava suite with a break. Thickness is 40-60 m. Suite is dated by ammonites *Stephanoceras humphriesianum* Sow., *S. (Cadomites) deslongchampsii* Defrance, *Parkinsonia parkinsoni* Sow., *Oppelia* cf. *subradiata* Sow., *Leptosphinctes leptus* Buck., *Phylloceras asisbekovi* Kakh., *Eurystomiceras polyhelictum* Bockh, *Pseudophylloceras kudernatschi* Hauer, *Calliphylloceras heterophyl-*

loides Opp., *C. disputabile* Zitt., *Thysanolytoceras eudesianum* Orb., *Dinolytoceras crimea* Strem., *Holcophylloceras zignodianum* Orb., *Perisphinctes defrancei* Orb., *Peltoceras athleta* Phill. and belemnites *Holcobelus blainvillei* Voltz., *Cylindroteuthis puzosiana* Orbigny, *C. spicularis* Phillips.

Variegated brecciated limestone strata – Uppermost Bajosian – Oxfordian. Represented locally in Pieniny Zone. Red, dark-cherry, brown-greenish, yellowish brecciated limestones, limestone breccia and conglomerates. Sometimes there are layers of marls and silicites at the top. Strata lies transgressive on Lower and Middle Jurassic deposits and is overlain by strata of light-coloured limestones. Thickness up to 26 m. Strata is dated by ammonites *Bullatimorphites* ex gr. *bullatimorphus* (Buckm.), *B. ymir* (Opp.), *B. eszterensis* (Bockh), *B. cf. polypleurus* (Buckm.), *B. costatus* Arkel., *Procerites* ex gr. *imitator* (Buckm.), *Dimorphinites* cf. *dimorphus* (Orb.), *Parkinsonia (Durotrigensia) bomfordi* Arkell, *Nannolytoceras* cf. *tripartitum* (Rasp.), *Perisphinctes (Dichotomoceras) bifurcatus* (Quenst.), *Wagnericeras* cf. *kudernatschi* (Liss.), *Liosphinctes plicatilis* (Sow.), *Euaspidoceras paucituberculatum* (Arkell), *Gregoryceras*, *Perisphinctes*, *Lytoceras*, *Phylloceras*, *Holcophylloceras*, *Ptychophylloceras*, *Partschiceras*, *Spiroceras*, *Parachoffatia*, *Homoeplanulites*, *Epistrenoceras*, mollusks, dinocists *Colomisphaera fibrata* (Nagy), *Cadosina parvula* Nagy, foraminifers *Globuligerina*, etc. [Славин, 1963; Rehakova et al., 2011].

Siliceous limestone strata – Upper Oxfordian–Lower Kimmerigian. Represented locally in the Northeast part of Pieniny Zone. Red laminate siliceous limestones, marls, bioclastic limestones. Strata lies on variegated brecciated limestones and is overlain by Red-coloured limestone strata. Thickness is more than 2 m. Strata is dated by dinocists *Colomisphaera fibrata*, *Cadosina parvula* [Rehakova et al., 2011].

Light-coloured limestone strata – Kimmerigian–Lowermost Tithonian. Represented locally in Pieniny Zone. Whitish and light grey laminate massive micritic and biomicritic limestones. Sometimes at the top, there are eroded horizon with phosphorite nodules and belemnites. Strata lies on variegated brecciated limestones and is overlain by Svalyava suite. Thickness up to 16 m. Strata is dated by ammonites *Orthaspidoceras uhlandi*, *Calliphylloceras ptychoicum* Quenst., *Lytoceras quadrisulcatum* Opp., *Haploceras tithonium* Opp., *Neolissoceras gzasi* Orb., *Nebroditis*, *Phylloceras*, *Lytoceras*, *Taramelliceras* mollusks, etc. [Славин, 1963; Wierzbowski et al., 2012].

Red-coloured limestone strata – Upper Kimmergian–Lower Berriassian. Represented locally in the Northeast part of Pieniny Zone. There are red spotted limestones in lower part, and pink micritic limestones with numerous planktonic organisms above. Strata lies on siliceous limestones and is sharply overlain by basaltic rocks of Early Cretaceous. Thickness up to 17 m. Strata is dated by tintinnids *Praetintinnopsella andrusovi* Borza, *Calpionella alpina* Lor., *C. grandalpina* Nagy, *Crassicollaria massutiniana* (Colom), *C. parvula* Remane, *C. brevis* Remane, *Tintinnopsella carpathica* Murg. et Fil., *T. doliphormis* (Colom), *Remaniella ferasini* (Catalano), *R. catalanoi* Pop, *R. durandelgai* Pop, etc. [Rehakova et al., 2011].

Svalyava suite [Славин, 1963] – Upper Tithonian–Lower Barremian. Pieniny zone of Median Carpathians and Vyshkiv Subzone of Pripannonian Zone (Inner Carpathians). Olistoliths and tectonic blocks (xenoliths) in Upper Cretaceous matrix. Isolated blocks of different sizes (0,2 to 2 km), configuration and orientation (before inverted). Their constituent units of different age are oriented irregularly and are bounded by curved faults. Light-colored sometimes porcelaneous limestones, with layers and lenses of black cherts, black or green mudstones and marls. Lower and upper boundaries are unknown. Thickness in the Carpathians is more than 20 m. Suite is dated by ammonites *Beriasella* sp., *Crioceras duvali* Lev., *C. cf. nolani* Kil., *Phyllopacyceras infundibulum* Orb., *Spitidiscus cf. andrusovi* Karak., *Barremites* sp., *Lamellaptychus cf. didaji* Coq., tintinnids *Calpionella alpina* Lor., *C. undelloides* Colom, *Tintinnopsella carpathica* Murg. et Fil., etc.

Marmarosh Massif. There were distinguished Obnizh, Rudarnya, Baltagul and Dovgoruny suites in Belopotik Subnapper.

Obnizh suite (V.I. Slavin, 1963) [Славин, 1963] – Hettangian–Toarcian. Belopotik Subnapper of Marmarosh Massif. Black laminated mud limestones, mudstones, gray calcareous sandstones, massive limestone breccia, conglomerates and carbonaceous rocks. Suite lies transgressive on the Lower Triassic, Upper Carboniferous or Upper Proterozoic sediments; the upper contact is unknown. Thickness is to 100 m. Suite is dated by ammonites *Arietites (Coroniceras) ex gr. herbichi* Uhl., bivalves *Cardinia* sp.

Rudarnya suite [Славин, 1956] – Aalenian–Callovian. Belopotik Subnapper of Marmarosh Massif. Dark gray argillites with siderite nodules and interlayers of siltstones and sandstones. At the

bottom there are the basal conglomerates and sandstones with interlayers of calcareous mudstones, siltstones and lenses of sandy limestones (up to 40-45 m), or a pack of cinnamon-brown bauxites and argillites (1,5-2 m). Thickness up to 70-100 m. Suite lies unconformably on Triassic and Palaeozoic and is overlain unconformably by Upper Jurassic Baltagul suite. Age is determined according to ammonites *Leioceras opalinum* Rein., belemnites *Nannobelus brevis* Blainv., brachiopods *Leilleria monbizotensis* Roll., etc.

Baltagul suite (V.I. Slavin, 1963) [Славин, 1963] – Oxfordian–Lower Tithonian. Belopotik Subnapper of Marmarosh Massif. Variegated calcareous sandstones, siliceous mudstones, jaspers with radiolarians, crinoid limestones, conglomerates. At the bottom there are conglomerates (2-3 m), at the top the thin-bedded sandstones and mudstones. Suite lies on the Upper Triassic and Middle Jurassic rocks. The upper boundary is not determined. Thickness is to 100 m. Suite is dated by radiolarians *Saturnalis amissus* Squin, brachiopods *Rhynchonella cf. alemanica* Roll., *Zelleria montbizotensis* Roll., etc.

Dovgoruny suite (V.I. Slavin, 1956) [Славин, 1956] – Oxfordian – Tithonian. Belopotik Subnapper of Marmarosh Massif. At the bottom (up to 5 m) there are conglomerates, breccias, gravellites from the wreckage of metamorphic rocks (gneisses, muscovite and two-mica schists, etc.); higher – sandstones, limestones, mudstones. In upper part there are gray and green-gray laminated often marbled limestones, with lenses of gray-green quartz-carbonate-chlorite shales (1-1,5 m). Suite lies with a sharp angular unconformity on the Upper Proterozoic, Upper Carboniferous, Lower Triassic, Lower Jurassic formations. The upper limit is not determined. Thickness is 200-400 m. Age is determined according by ammonites *Perisphinctes stenocloides* Siem., *Progeronia breviceps* (Quenst.).

Kamyapotik Napper. There were distinguished Chivchini and Kamyapotik suites.

Chivchini suite (V.I. Slavin, 1956) [Славин, 1956] – Oxfordian–Tithonian. Kamyapotik Napper. In the lower part there are dark green, brick-brown porphyres, spilites, diabases and tuffs, above – breccia, conglomerates with lenses of biohermal and micritic platy limestones and jaspers. Thickness of flows is 15-300 m, tuffs – 12-180 m, limestones – 5-10 m, jaspers – 1-1,5 m. The lower boundary is not found. Suite is overlapped conformly by Kamyapotik suite. Thickness is more than 1000 m. Suite is dated by corals *Calamophylloopsis etalloni* (Koby), *Thecosmilium irregularis* Etall., *Th. koniakensis* Ogil.,

Montivaltia elongata E. et H., *Actinaraea granulata* (Munst.), etc. [Лозиняк, Марушкін, 1991].

Камуанопотик suite (J.O. Kulchyt'sky, O.V. Maksimov, 1961) [Кульчицький, Максимов, 1961] – Tithonian–Hauterivian. Камуанопотик Napper. Gray and dark gray layered or massive limestones (80% of the suites), mudstones, sandstones (generally calcareous with volcanic rocks in the clastic material) and gravelites. Suite lies conformly on Chivchiny suite and it is overlapped with erosion by the Lower Cretaceous sandstone strata [Лозиняк, Марушкін, 1991]. Thickness is to 200 m. Suite is dated by ammonites – *Costidiscus recticostatus* Orb., *Speetonicerias auerbachii* Eichwald, *Berriasella subchaperi* Retowski, bivalves *Camptonectes cottaldinus* Orb., *Syncyclonema germanica* Woll., brachiopods – *Suiaella weberi* Moisseev, radiolarians – *Cenospaera*, *Xiphosphaera*, *Cenodiscaella*, *Dicolocapsa*, *Dictyomitra*, tintinnids – *Calpionella alpina* Lor., *C. elliptica* Cad., nannoplankton – *Vipodorhabdus roeglii* (Thier.), *Cretarhabdus crenulatus* Braml. et Mart., *Watznaueria barnesae* (Black), *W. britannica* (Str.), *Polycostella beckmanni* Thier., etc.

Porkulets Napper. There is only distinguished the Trostjanets strata.

Trostjanets strata [Ломізе, 1975] – Tithonian – Lower Cretaceous. Porkulets Napper. Olistoliths and olistoplakes in Albian–Upper Cretaceous olistostrome [Славин, 1956] or tectonic blocks. Diabases and volcanic breccia with lenses of cream, pink and gray limestones. Streams of lava (5–32 m thickness) are divided by klastolavas. Underlying and overlying rocks are not reliably determined. Thickness is 150 m. Strata is dated by ammonites *Caliphilloceras ptichoicum* Quenst., *C. callipso* Orb. var. *zacarpatiensis* Slavin, *Lythoceras quadrisulcatum* Orb., *L. montanum* Opp., *Streblites loliensis* Slavin, tintinnids *Calpionella alpina* (Lor.), etc.

Basement of Transcarpathian Depression. In the Jurassic deposits of before-molassa basement of Transcarpathian Depression were distinguished the Jurassic carbonate-terrigenous strata, Svalyava, Sharock and Bakta suites.

Jurassic carbonate-terrigenous strata (M.G. Prykhodko, 2012) [Стратиграфія..., 2014]. Uzhgorod-Solotvino zone. Interbedding of gray mudstones, marls, sandstones, siltstones, and micritic and bioclastic limestones. This strata conformly or tectonically overlies the Triassic rocks; it is conformly overlapped by Dulov suite of Lower Cretaceous and is often significantly eroded. Thickness is to 900 m. Age as a whole Jurassic is determined by bivalves *Bositra (Posidonia) buchi* (Roemer),

B. dagestanica (Uhlig), tintinnids *Crassicollaria intermedia* (Delga), *C. brevis* Remane, *C. parvula* Remane, *Calpionella alpina* Lorenz, *C. alpina grandis* Doben, *Tintinnopsella carpathica* (Murg. et Fil.), *Calpionellites darderi* (Colom), *Lorenziella hungarica* Knauer et Nagy, and by the position in the section and the analogy with deposits of eastern Slovakia also.

Sharock suite (P.Yu. Lozynyak, M.J. Petrashkevich, 1993) [Лозиняк, Петрашкевич, 1993] – Lower–Middle Jurassic. Berehovo Subzone of Pripannonian Zone. Interbedding of gray limestones and marls, sandstones, mudstones and siltstones with lenses of diabases. Sharock suite lies conformly on the Triassic rocks and is overlain by Bakta suite. Thickness is to 150 m. Suite is dated by belemnites *Passoloteuthis*, *Holcobelus* and position in the section also.

Bakta suite (P.Yu. Lozynyak, M.J. Petrashkevich, 1993) [Лозиняк, Петрашкевич, 1993] – Middle–Upper Jurassic. Berehovo Subzone of Pripannonian Zone. Gray and variegated basic volcanics with interbedded gray sometimes variegated marls with numerous radiolarians, variegated massive limestones, jasper-like siliceous-carbonate rocks and dark gray mudstones. Suite lies conformly on sharock suite, and it is overlapped with a break (?) by the Cretaceous deposits. Thickness is to 200 m. Suite is dated by radiolarians *Cenodiscus rachovenssis* Loz., *Tricolocapsa plicarum* Yao, *Hemicrycaspa vialovi* Tichom., *H. a* (Zham.), *Dictyomitra venusta* Hinde etc., belemnites *Megateuthis*, as well as by the position in the section.

Svalyava suite (V.I. Slavin, 1963) [Славин, 1963] – Tithonian–Lower Barremian. Vyshkiv Subzone of Pripannonian Zone. Gray and light gray silicified micritic sometimes brecciated or clastic limestones with inclusions of black cherts and thin interlayers of dark gray siliceous mudstones. Suite is overlapped by Cretaceous Tissale suite; the lower boundary is not determined. Suite is dated by radiolarians *Xiphosphaera umbilicata* Rust, *Cenosphaera sphaeroconus* Rust, *Cenodiscaella nummulitica* Kh. Aliev, *Cornotella conica* Kh. Aliev, *Dictyomitra carpatica* Lozynyak and by the analogy with the deposits of Pieniny Zone.

Carpathian Foredeep and adjacent part of East-European Platform (Stryi Jurassic Deep). The Jurassic deposits form there a submeridional strip up to 100 km width. Thicknesses, as well as the completeness of sections, are increasing to the west from the few meters in Volyno-Podillya up to 2500 m near the Krakovets Fault. Jurassic deposits

transgressive overlies the eroded Paleozoic basement and are overlapped with erosion by the Cretaceous and Neogene formations. Lower and Middle Jurassic are extended in the northwestern part of Bilche-Volitsa zone of Carpathian Foredeep and locally on the platform and represented by mainly clastic sediments. The Upper Jurassic spread in a whole Stryi Jurassic Deep and together with Berriasian-Valanginian forms complex of carbonate rocks. In terrigenous deposits of Lower and Middle Jurassic were distinguished Komarno, Bortyatin, Podiltsy, Medenychi, Kochaniwka, Sokal and Yavoriv suites. Carbonate complex of Upper Jurassic–Lower Cretaceous consist of Boniv, Gorodok, Rudky, Morantsy, Karolino, Opary, Rava-Ruska, Podluby, Nizhnyv, Bukivna suites [Жабина, Анікеєва, 2007].

Komarno suite (V.G. Dulub et al., 2003) – Hetangian. Northwest of Bilche-Volitsa zone of Carpathian Foredeep. Interbedding of gray fine-grained quartz and quartzite-like sandstones, mudstones, siltstones. Suite is overlapped by Bortyatin suite; the lower boundary is not determined. Thickness is more than 560 m. Age is determined conditionally according by the presence of Mesozoic spores – *Densoisporites* sp., *Lycopodiumsporites* sp., *Granulatisporites minutus* Pot. et Kremp. as well as by the position in section.

Bortyatin suite (Yu.R. Karpenchuk, 1985; V.G. Dulub et al., 1986) – Sinemurian. Northwest of Bilche-Volitsa zone. Greenish-gray siltstones interbedded with gray fine-grained quartz sandstones and dark gray non-laminated silty mudstones and siltstones. It lies conformly on the Komarno suite and is overlain by Podiltsy suite. Thickness is to 295 m. Suite is dated by the palynocomplex *Limboisporites lundbladii* Nilsson, *Platyptera trilingua* (Horst) Schulz, *Stereisporites* sp., *Camptotriletes* sp., *Densoisporites* sp., *Dictyophyllidites harrisii* Couper, *Cyachidites junctus* (K.-M.) Alimov., as well as by the position in section.

Podiltsy suite (Yu.R. Karpenchuk, 1985; V.G. Dulub et al., 1986) – Pliensbachian. Northwest of Bilche-Volitsa zone. The rhythmic interbedding: at the lower part there are dark gray shales and siltstones, above gray and light gray cross-bedded siltstones and sandstones with interlayers of limestones in the upper part, and with streaks and nests of anhydrite at the top. It lies according on the Bortyatin suite and is unconformably(?) overlapped by Medenychi or Kochaniwka suites. Thickness is to 580 m. Suite is dated by the palynocomplex *Lycospora salebrosacea* (Mal.) Schulz, *L. gracilis* Sem., *Limboisporites lundbladii* Nilsson, *Platyptera trilingua*

(Horst) Schulz, *Foraminisporites jurassicus* Schulz, *Converrucosisporites luebbenensis* Schulz, *Uvaesporites verrucatus* Schulz., etc.

Medenychi suite (V.N. Utrobin, 1962) [Утробин, 1962] – Toarcian. Northwest of Bilche-Volitsa zone. Gray quartz and quartzite-like sandstones (somewhere cross-bedded, porous, with lenses and interlayers of coal) with streaks of gray non-carbonate siltstones, silty sometimes siliceous mudstones, creamy-gray limestones. Rocks contain coalified fitodetrit and somewhere imprints of plants. Suite lies with erosion on Podiltsy suite and is overlapped by Kochaniwka suite. Thickness is to 600 m. Suite is dated by microflora *Cycadopytes dilucidus* (Bolch.) Iljina, *C. orbicularis* (Sachon et Iljina), *Ginkgocycadophytus caperatus* Samoil., *Chasmatosporites apertus* (Rogalska) Nilsson, *Stereisporites congregatus* (Bolch.) Schulz, *S. bujargiensis* (Bolch.) Schulz, *S. incertus* (Bolch.) Sem., *Selaginella simplex* Krasn., *S.ensis* Chlon. et Krasn., *Osmundacidites jurassicus* (K.-M.) Kuzit., *O. versiformis* Sem. sp. nov., *O. cingulatus* Sem., *Marattisporites scabratus* Couper, *Tripartina variabila* Mal., *Klukisporites variegatus* Couper.

Kochaniwka suite (J.M. Sandler, 1962) [Сандлер, 1962] – Toarcian–Bathonian. Northwest of Bilche-Volitsa zone. Dark gray argillites with organogenic detritus, interbedded with thin-layered gray micaceous sandstones and siltstones, dark gray bioclastic limestones. It lies on Medenychi or Podiltsy suites, and is transgressive overlapped by Yavoriv suite. Thickness is to 582 m. Suite is dated by mollusks *Meleagrinnella ptchelincevae* Polub., *Paleoneilo galata* (Orb.), *Dacryomya zietenii* (Brauns.), *Nucula amygdaloides* Som., *Inoceramus ambiquus* Eichw., *Ozytoma startense* Polub., *Posidonia dagestanica* Uhlig., *Parkinsonia parkinsoni* Sowerby, *Partschiceras abichi* (Uhlig), *Pholadomya solitaria* Morris et Lycett., *Goniomya baysunensis* Borissjak, *Pleuromya decurtata* Phillips, *Cucullaea subdecussata* Gold., *Leda mucronata* Sow., *L. lacryma* Sow., *Phaenodesmia arzisiensis* Romanov, *Nucula subovalis* Goldf., *Parallelodon elongatum* Sow., *Oxytoma scaburgense* Rollier, *Pinna buchi* Koch et Dunker, *Lima (Plagiostoma) subrigidula* Schlippe, *Entolium demissum* Goldf., *Pleuromya balkhanensis* Pčel., *Reinholdella dreheri* (Bart.), *R. epistominoides* (Kapt.), *Camptotriletes cerebriformis* Naum., *C. triangulus* Larosch., foraminifers *Trochammina chodzica* Ant., *Quinqueloculina rawensis* (Pazdro), *Lenticulina erecta* Ant., *L. volubilis* Dain, palynocomplex *Duplexisporites anagrammensis* (K.-M.) Schug., *Marattisporites*

scabratus Couper, *Klukisporites variegatus* Couper., *Tripartina variabilis* Mal., *Trilites minutus* (Bolch.) Mal. emend. Sem., etc.

Sokal suite (V.I. Slavin, V.Ya. Dobrynina, 1958) [Славин, Добрынина, 1958] – Bajosian–Bathonian. Volyno-Podillya. Continental non-carbonate clays, siltstones, sandstones, gravelites, conglomerates. Suite consists of two subsuites: lower – gray lake-marsh sediments with imprints of flora and upper – variegated alluvial-proluvial-deluvial deposits. Suite lies on the Paleozoic and is overlapped by Rava-Ruska suite. Thickness is to 60 m. Age is determined according to floristic remnants *Hylomites zamites* Goepf., *Coniopteris lymenophylloides* Brongn., *Phoenicopsis speciosa* Heer., *Ptilophyllum cutchense* Old et Mor., *Pt. acutifolium* Teist., *Nilssonia orientalis* Heer., *Abietites densifolia* Thomas, *P. tyophyllum nordenskioldi* (Heer.) Nath., *P. lindstromii* Nath., *Leplophyllum subcirculata* Pryn., *Taeniopteris tenuinervis* Brans., *Eboracia labiflora* (Pill.) Tom., etc, and also by the sedimentary criterias and position in section.

Yavoriv suite (V.M. Utrobin, 1962) [Утробин, 1962] – Callovian. Northwest of Bilche-Volitsa zone and adjacent part of the platform. Gray ocherous non-carbonate sandstones and quartz siltstones, sometimes dolomitic or clayey, with glauconite. At the bottom there are conglomerates and gravellites and at the top – interlayers of gray sandy dolomites and breccia, dolomitic micritic or bioclastic limestones. Somewhere there are interlayers of gray, brown or mottled clays. The rocks contain coalified fitodetrit and ferruginous oolites. Suite lies transgressive on Kochaniwka suite and overlain conformly by Rudky suite. Thickness is to 100 m. Suite is dated by mollusks *Macrocephalites* sp., *Kepplerites* sp., *Kosmoceras* sp., *Camptonectes* cf. *lens* (Sow.), *Dinolytoceras* cf. *crimea* Streumoukhoff, *Melegrinella* sp., *Phylloceras* sp., *Posidonia* (*Bositra*) *buchi* Roemer, *Pholadomya* cf. *murchisoni* Sowerby, *Cucullaea subdecussata* Gold., *Leda mucronata* Sowerby, *Planularia folium* (Wisn.), *P. polyra* (Guemb.) and palynomorphs *Classopollis clasoides* Pflug.

Rudky suite (V.M. Utrobin, 1962) [Утробин, 1962] – Oxfordian. Northwest of Bilche-Volitsa zone and adjacent part of the platform. Biohermal limestones. At the bottom there are sponge limestones with chert nodules and interlayers of spongolites, above coral-algae limestones, and at the top oncolitic limestones and sometimes stromatolites. Between the bioherms there are bioclastic and biomicritic limestones interbedded with siltstones and

limestone breccia. Suite lies on Yavoriv suite and is overlapped by Opary, Pidluby or Rava-Ruska suites. Thicknesses of bioherms are up to 65-140 m; between the bioherms – 40-60 m. Suite is dated by ammonites *Subkossmatia* cf. *opis* (Sow.), brachiopods *Septaliphora badensis* (Opp.), *Cheirothyris aculeata* Leit., *Rhynchonella varians* Schloth, *R. badensis* Opp. and foraminifers *Globuligerina oxfordiana* Grig., *Trocholina transversarii* Paalz., *T. conica* (Schlumb.), *T. belorussica* Mitjan., *Bulbobaculites maynci* Mohl., *Alveosepta sequana* (Merian), *A. sequana* (Merian) *minor* Mohl., *Haplophragmium coprolithiformis* Mohl., *Marssonella doneziana* Dain, etc.

Boniv suite (V.G. Dulub, 1995) [Дулуб, 1995] – Oxfordian. Northwest of Bilche-Volitsa zone. Interbedding of gray, brown-gray micritic and pellete limestones and mudstones. It conformly lies on Yavoriv suite and is overlain by Morantsy suite. Thickness is 50-280 m. Suite is dated by foraminifers *Globuligerina oxfordiana*, *Alveosepta jaccardi* Schrodt, *Alveosepta sequana* var. *minor*, *Bulbobaculites maynci*, *Marssonella doneziana*, *Trocholina conica*, *Paalzwella turbinella* (Gumb.), *Discorbis speciosus* Dain, *H. coprolithiformis sequanum*, *Trocholina transversarii*, *Quinqueloculina semisphaeroidalis* Danitch. and tintinnids *Foliacella propartula* Makar., *Scalpratella angustioris* Makar., *Rosiella tintinnubulum* Makar., *Borzaiella terekensis* Makar., etc.

Gorodok suite (N.M. Zhabina, O.V. Anikeyeva, 2007) [Жабіна, Анікеєва, 2007] – Oxfordian. Northwest of Bilche-Volitsa zone and adjacent part of the platform. Gray, cream-gray bioclastic and oncolitic limestones with interlayers of dolomites, sandstones, siltstones, mudstones. It lies on Yavoriv suite or eroded Paleozoic basement and is overlapped by Rava-Ruska suite or Neogene. Thickness is 24-100 m. Suite is dated by foraminifers *Haplophragmium suprajurassicum* (Schwag.), *Bulbobaculites maynci*, *Alveosepta jaccardi*, *A. sequana*, *Marssonella doneziana*, *Discorbis speciosus*, *Paalzwella scalariformis* Dain, *Torinosuella peneropliformis* (Yabe et Hanzlik.), *Trocholina transversarii*, *Paalzwella turbinella*, *Spirothalmidium dilatatum* (Paalz.), etc.

Opary suite (I.B. Vishnyakov, 1978) [Вишняков, 1978] – Kimmeridgian–Lower Berriassian. Northwest of Bilche-Volitsa zone. Suite is composed of two subsuites: lower (up to 450 m) – gray micritic, bioclastic, brecciated limestones with a sponge-algal bioherms (Kimmeridgian); and upper (400-1100 m) – light-colored biohermal limestones (coral, algal, etc), bioclastic, oncolitic, brecciated

(Tithonian–Lower Berriasian). Suite lies on Rudky or Boniv suites and is overlapped with erosion by Neogene. Age is determined by foraminifers *Mesoendothyra izjumiana* Dain, *Everticyclammina virguliana* (Henson), *Charentia compressa* (Cushm. et Glaz.), *Orbignyoides podolicus* (Cushm. et Glaz.), *Alveosepta powersi* (Redm.), *A. personata* (Tobler), *Pseudocyclammina bukowiensis* Cushm. et Glaz., *P. sphaeroidalis* Hotting., *Rectocyclammina arrabidensis* Remalho, *Feurtillia frequens* Maync, *Anchispirocyclus lusitanica* (Egger), *Melathrokerion spirialis* Gorb., *Freixialina planispiralis* Remalho, *Quinqueloculina mitchurini* Dain, *Neotrocholina friburgensis* (Guill. et Reich.), *Palaeotextularia crimica* Gorb., *Belorussiella taurica* Gorb., etc.

Morantsy suite (V.G. Dulub, 1995) [Дулуб, 1995] – Kimmeridgian. Northwest of Bilche-Volitsa zone. Interbedding of dark gray mudstones, mud limestones (micritic, pelletous) and limestone breccia. Suite conformly lies on Boniv suite and is overlain by Karolino suite or upper subsuite of Opary suite. Thickness is to 245–300 m. Suite is dated by foraminifers *Mesoendothyra izjumiana*, *Pseudocyclammina powersi* Redm., *Conicospirillina basiliensis* Mohl., *Spirillina polygyrata* Gumb., *Alveosepta jaccardi*, *Haplophragmium coprolithiformis*, etc.

Pidluby suite (J.M. Sandler, 1962) [Сандлер, 1962] – Lower Kimmeridgian. Northwest of Bilche-Volitsa zone. Gray, cream, lime green pelletous, onkolitic limestones and dolomites with interlayers of gray mudstones at the bottom. It conformly lies on Rava-Ruska or Rudky suites and is overlapped by Nyzhniv suite. Thickness is to 250 m. Suite is dated by foraminifers *Choffatella tingitana* Hott., *Alveosepta personata*, *A. powersi*, *Torinosuella peneropliformis*, etc.

Rava-Ruska suite (V.I. Slavin, 1956) [Славин, 1956] – Lower Kimmeridgian. East periphery of Carpathian Foredeep and adjacent part of the platform. Suite consists of two subsuites. Lower subsuite (20–45 m) are represented by variegated mudstones, clays, siltstones, sandstones, gravellites, conglomerates, breccias, limestones, dolomites with interlayers and inclusions of gypsum and anhydrite; upper subsuite (200 m) – by gray dolomites, limestones, mudstones, anhydrites, gypsum, breccia. Suite lies on Gorodok or Rudky suites or eroded Paleozoic basement and is overlapped by Nyzhniv suite or the Cretaceous formations. Thickness is to 250 m. Suite is dated by foraminifers *Mesoendothyra izjumiana*, *Pseudocyclammina ukrainica* Dain, *P. parvula*, *Alveosepta powersi*, *A. personata*, *Choffatella tingi-*

tana, *Torinosuella peneropliformis*, *Discorbis sub-speciosus*, *Conicospirillina basiliensis*, *Labyrinthina mirabilis* Weynch., *Choffatella tingitana*, etc.

Nyzhniv suite (J.M. Sandler, 1962) [Сандлер, 1962] – Upper Kimmeridgian–Lower Tithonian. East periphery of Carpathian Foredeep and adjacent part of the platform. In lower part there are mottled conglomerate-breccia, above the light gray micritic and onkolitic limestones and dolomites interbedded with marls and dark gray mudstones. Suite lies on Pidluby or Rava-Ruska suites, or on the eroded Paleozoic basement. It is overlapped by Bukivna suite or with erosion by deposits of Albian, Cenomanian or Neogene. Thickness up to 200 m. Suite is dated by mollusks *Corbula inflexa* Roem., *Exogyra virgula* Defr., *Pteroceras oceani* Brogn., algae – *Petrascula bursiformis* Gium. and foraminifers *Mesoendothyra izjumiana*, *Pseudospirocyclus maynci* Hott., *P. mauretanicus* Hott., *Alveosepta personata*, *A. powersi*, *Haplophragmium coprolithiformis*, *Conicospirillina basiliensis*, *Anchispirocyclus lusitanica* (Egger), *Pseudolamarkina obliquicamerata* Dulub, etc.

Bukivna suite (V.I. Slavin, 1956) [Славин, 1956] – Upper Tithonian – Lower Berriasian. East periphery of Carpathian Foredeep and adjacent part of the platform. Light-colored bioclastic and oncolitic limestones, sometimes interbedded with limestone breccia. Suite lies on Nyzhniv suite and is overlapped by the Upper Berriasian or Cenomanian. Thickness is to 80 m. Suite is dated by mollusks *Corbula inflexa* Roem., *Purpurina subnodosa* Roem., *Cerithium septemplicatum* Roem., *Pleuromya jursasica* Brg., *Anisocardia pulchella* Lor., *A. parvula* Roem., *A. legayi* Lor., *Lucina substriata* Roem., *Mytilus longaevis* Ctj., *Avicula gessneri* Th., *Hinnites velatus* Gdf., *Exogyra virgula* Defr., *Nautilus geinitzi* Opp., *Trochus betancourti* Lor., *Turbo durki* Lor., *Bulla cylindrella* Buv., *Sphaenia saemanni* Lor., *Anisocardia intermedia* Lor., *Lucina vernerii* Etal., *Astarte saemanni* Lor., *Opis portlandicus* Lor., *Amonia suprajurensis* Roem., *Nerinea constricta* Roem., *Nautilus geinitzi* Opp., brachiopods *Terebratula subsella* Leym., *Waldheimia pentagonalis* Br., serpulids *Serpula conformis* Gdlf., *S. subflaccida* Rtal., algae *Petrascula bursiformis*, foraminifers *Anchispirocyclus lusitanica*, *Feurtillia frequens*, *Freixialina planispiralis*, *Melathrokerion spirialis*, *Belorussiella taurica* Gorb., *T. burlini* Gorb., *T. molesta* Gorb., etc.

Karolina suite (V.G. Dulub, 1995) [Дулуб, 1995] – Tithonian – Lower Valanginian. Northwest of Bilche-Volitsa zone. Two subsuites: Lower (Tithonian–Lower Berriasian) is represented by limestones, limestone breccia, mudstones, sandstones

with glauconite (up to 430 m), Upper (Upper Berri-Asian–Lower Valanginian) is represented by interbedding of limestones and mudstones (up to 200 m). Suite lies on Morantsy suite, and is overlapped with erosion by Neogene formations. Suite is dated by tintinnids *Longicollaria dobeni* (Borza), *Crassicollaria brevis* Remane, *C. intermedia* Durand Delga, *C. massutiniana* (Colom), *C. parvula* Remane, *Calpionella alpina*, *C. grandalpina* Nagy, *C. elliptalpina* Nagy, *C. elliptica* Cadish, *Remaniella ferasini* Catalano, *R. cadishiana* (Colom), *Tintinnopsella longa* (Colom), *T. remanei* Borza, *T. doliphormis* (Colom), *Calpionellopsis oblonga* (Cadish), *C. simplex* (Colom), *Calpionellites coronata* Trejo and foraminifers – *Pseudocyclamina bukowiensis*, *Anchispirocyclina lusitanica*, *Freixialina planispiralis*, *Protopenneroplis striata* Weynsh., *Mesoendothyra izjumiana*, *Feurtillia frequens*, *Melathrokerion spiralis*, *Belorussiella taurica*, *Palaeotextularia crimica*, *Trocholina burlini*, *T. molesta*, *Neotrocholina friburgensis*, *N. vasserodi* Guill., *N. valdensis* Reich., *N. infragranulata* Noth, *Marsonella oxycona* (Reuss), *M. pseudocostata*, *Quinqueloculina triola* Macieva et Temirbek., *Moesiloculina danubiana* (Neagu), *Scytiloculina confusa* Neagu, *Spiroloculina speciosa* Macieva et Temirbek., etc.

Kovel Ledge of the East-European platform basement. In the Northwestern Ukraine, in the area of the Shatsk Lakes the Middle–Upper Jurassic deposits were opened by wells. There were distinguished Svityaz and Shatsk suites [Гаврилишин, 1993].

Svityaz suite (V.I. Havrylyshyn, 1993) [Гаврилишин, 1993] – Bathonian–Callovian. Northwestern part of Volyn and Southwestern Bielorussian. At the bottom there are gray lake-marsh silty and sandy muds with coalified fitodetrit and fragments of stems, above – gray clayey quartz sands with glauconite and small shelly detritus. Suite lies on the Proterozoic and is overlain by Shatsk suite. Thickness is 10-40 m and more. Suite is dated by palynomorphs *Classopollis classoides* Pflug., *Scidopityspollenites macroverrucosus* (Their.), etc. (Дулуб и др., 2003).

References / Список літератури

1. Бирюлев Л.В., Измалкова С.П., Никулина Т.В., Полонский Б.Т. Методические указания по применению стратиграфических схем и условных обозначений для геологических карт Украинских Карпат и Волыно-Подоллии масштаба 1:50 000. Киев, 1974. 64 с.

Biryulev L.V., Izmalkova S.P., Nikulina T.V., Polonsky B.T., 1974. Methodical instructions for the use of stratigraphic charts and symbols for geological maps

Shatsk suite (V.I. Havrylyshyn, 1993) [Гаврилишин, 1993] – Callovian–Oxfordian. Southwestern part of Bielorussian and northwestern part of Volyn. Gray calcareous sandstones with organogenic detritus transformed into bioclastic sandy limestones with oolites. It lies on the Svityaz suite, and with erosion are overlapped by the Cretaceous. Thickness is 15-20 m. Suite is dated by bivalves – *Trigonia recticostata* Lycet, *T. bronni* Agassis, *T. densicostata* Roeder, gastropods *Pleurotomaria thoetensis* Heb. et Desl., etc.

Conclusions. In this paper the Generalized Correlative Scheme of Stratigraphy of Jurassic of the Western Ukraine is represented. The current stratigraphic scheme [Стратиграфія ..., 2014] was supplemented and detailed according to modern data, analysis and re-interpretation of many-years investigations Jurassic deposits by different specialists. The characteristics of stratigraphic units: distribution, lithological and paleontological composition, position in section, thicknesses and criteria for dating are given.

The most complete and precise sequence of the Jurassic stages is determined on the territory of Carpathian Foredeep and adjacent part of the East European platform. In other regions these deposits are present fragmentarily. The stratigraphic units are mainly dated by the complexes of macro- and microfossils. In Precarpathians the Lower Jurassic are dated by palynocomplexes, Middle Jurassic – by mollusks, foraminifers and palynocomplexes, Upper Jurassic – by foraminifers, mollusks and tintinnids. In Carpathians the Jurassic rocks are predominantly tectonic blocks and olistoliths occurring in the thicknesses of Cretaceous age. Here the primary sequence of strata was determined. In Carpathians the Lower and Middle Jurassic are dated by ammonites and belemnites and Upper Jurassic – by ammonites, tintinnids and dinocysts. Now the stage division of Jurassic in the basement of Transcarpathian Deep is not determined. There the distinguished strata are dated by age of mollusks, radiolarians and tintinnids. Correlation of the Jurassic stratigraphic units of the Western Ukraine is done.

of the Ukrainian Carpathians and Volyn-Podolia of scale 1:50 000. Kiev, 64 p. (in Russian).

2. Вишняков И.Б. Особенности геологического строения и развития Волыно-Подоллии и Предкарпаття: автореф. дис. ... канд. геол.-минерал. наук. Львов, 1978. 25 с.

Vishnyakov I.B., 1978. The features of geological structure and development of Volyno-Podolia and Precarpathians: The candidat thesis. Lvov, 25 p. (in Russian).

3. **Гаврилишин В.І.** Стратиграфія юрських відкладів Волині. *Геологія і геохімія горючих копалин*. 1993. № 1 (82). С. 8–12.

Havrylshyn V.I., 1993. Stratigraphy of Jurassic deposits in Volyn. *Geologiya i geokhimiya goryuchyh kopalyн*, № 1 (82), p. 8–12 (in Ukrainian).

4. **Геологія СССР**. Т. 48. Карпаты. Ч. 1. Геологическое описание: Сидоренко А.В. (гл. ред.). Москва: Недра, 1966. 540 с.

Geology of USSR. Vol. 48 Carpathians. Pt 1. Geological description: Sidorenko A.V. (Ed.), 1966. Moscow: Nedra, 540 p. (in Russian).

5. **Гоцанюк Г.І.** Моллюски і стратиграфія юрських відкладів Пенінської зони Українських Карпат: автореф. канд. дис. Київ, 2004. 22 с.

Gotsanyuk G.I., 2004. Mollusks and stratigraphy of the Jurassic deposits of the Pieniny zone of Ukrainian Carpathians: The candidat thesis. Kyiv, 22 p. (in Ukrainian).

6. **Дулуб В.Г.** Стратифікація депресійних утворень верхньої юри Більче–Волицької зони Передкарпатського прогину. *Нафта і газ України: наук.–практ. конф.*, 17–19 трав. 1994 р. Львів: УНГА, 1995. Т. 1. С. 118.

Dulub V.G., 1995. The stratification of depressive formations of Upper Jurassic in Bilche-Volitsa zone of Carpathian Foredeep. *Oil and gas of Ukraine: science-practical conf.*, 17–19 May 1994. Lviv: UNGA, vol. 1, p. 118 (in Ukrainian).

7. **Жабіна Н.М., Анікеєва О.В.** Оновлена стратиграфічна схема верхньої юри–неокому Українського Передкарпаття. *Зб. наук. пр. УкрДГРІ*. 2007. № 3. С. 46–56.

Zhabina N.M., Anikeyeva O.V., 2007. The new stratigraphic scheme of Upper Jurassic-Neocomian of Ukrainian Precarpathians. *Zbirnyk naukovykh prats Ukr-DGRI*, № 3, p. 46–56 (in Ukrainian).

8. **Жабіна Н.Н., Шлапинский В.Е., Приходько М.Г., Анікеєва Е.В., Мачальский Д.В.** Стратиграфія юрських отложений Запада Украины. *Buletinul Institutului de Geologie și Seismologie al AȘM*. 2015. № 1-2. С. 65-76.

Zhabina N.M., Shlapinsky V.Ye., Prykhodko M.G., Anikeyeva O.V., Machalsky D.V., 2015. Stratigraphy of Jurassic deposits of the West of Ukraine. *Buletinul Institutului de Geologie si Seismologie al ASM*, № 1-2, p. 65-76 (in Russian).

9. **Калениченко Т.Д., Круглов С.С., Мигачева Е.М.** До стратиграфії відкладів середньої юри Радянського Закарпаття. *Доп. АН УРСР*. 1965. № 9. С. 1193–1196.

Kalenichenko T.D., Kruglov S.S., Migacheva E.M., 1965. To the stratigraphy of Middle Jurassic deposits of the Soviet Transcarpathians. *Dopovidi AN URSSR*, № 9, p. 1193–1196 (in Ukrainian).

10. **Калениченко Т.Д., Круглов С.С.** Аммониты нижней юры Пенинской зоны (Закарпатье). *Палеонтол. сб.* 1969. № 6. Вып. 2. С. 32–38.

Kalenichenko T.D., Kruglov S.S., 1969. The Ammonites of Lower Jurassic of Pieniny Zone (Transcarpathians). *Paleontologichsky sbornik*, № 6, iss. 2, p. 32–38 (in Russian).

11. **Круглов С.С., Гаврилишин В.И., Крымгольц Г.Я.** Новые данные по стратиграфии юрских отложений Украины (с. Приборжавское в зоне Пенинских утесов Закарпаття). *Геол. журн.* 1968. № 5 (122). С. 87–91.

Kruglov S.S., Havrylshyn V.I., Krimgolts G.J., 1968. The new data on the stratigraphy of the Jurassic deposits of Ukraine (s. Priborgzhavskoye in the Zone of Pieniny Cliffs of Transcarpathians). *Geol. zh.*, № 5 (122), p. 87–91 (in Russian).

12. **Кульчицкий Я.О., Максимов О.В.** Стратиграфія крейдових відкладів хребта Чорногори в Українських Карпатах. *Доп. АН УРСР*. 1961. № 8. С. 1066–1068.

Kulchitsky Ya.O., Maksimov O.V., 1961. Stratigraphy of Cretaceous deposits of Chornogory Ridge in Ukrainian Carpathians. *Dopovidi AN URSSR*, № 8, p. 1066–1068 (in Ukrainian).

13. **Лещух Р.Й., Гоцанюк Г.І.** Палеонтологічне обґрунтування стратифікації юрських відкладів Пенінської зони Українських Карпат. *Геол. журн.* 2002. № 1 (299). С. 93–99.

Leschukh R.J., Gotsanyuk G.I., 2002. Paleontological substantiation of the stratification of the Jurassic deposits of the Pieniny zone of the Ukrainian Carpathians. *Geol. zh.*, № 1, (299). p. 93–99 (in Ukrainian).

14. **Лещух Р.Й., Гоцанюк Г.І.** Нова біостратиграфічна схема юрських відкладів Пенінської зони Українських Карпат. *Вісн. Львів. ун-ту. Сер. геол.* 2003. Вип. 17. С. 3–8.

Leschukh R.J., Gotsanyuk G.I., 2003. New biostratigraphic scheme of the Jurassic deposits of the Peninsula zone of the Ukrainian Carpathians. *Visnyk Lvivskogo Universitetu. Ser. Geol.*, vol. 17, p. 3–8 (in Ukrainian).

15. **Лінецька Л.В.** Tintinnoidea, Saccosoma та Globochaete карбонатного мезозою гори Великий Камінець (Бескидова зона Радянських Карпат). *Доп. АН УРСР. Сер. Б.* 1972. № 1. С. 17–20.

Linetska L.V., 1972. Tintinnoidea, Saccosoma and Globochaete of carbonate Mesozoic of the Velyky Kamynets Mountain (Beskidy Zone of the Soviet Carpathians). *Dopovidi AN URSSR. Ser. B*, № 1, p. 17–20 (in Ukrainian).

16. **Лозиняк П.Ю., Марушкін О.І.** Про вік чивчинської світи Українських Карпат. *Палеонтол. сб.* 1991. № 28. С. 71–73.

Lozinyak P.Yu., Marushkin O.I., 1991. About the age of Chivchiny suite of Ukrainian Carpathians. *Paleont. sb.*, № 28, p. 71–73 (in Ukrainian).

17. **Лозиняк П.Ю., Петрашкевич М.Й.** Нові дані про юрські відклади фундаменту Закарпатського прогину. *Палеонтол. зб.* 1993. № 29. С. 61–65.

Lozinyak P.Yu., Petrashkevich M.J., 1993. New data on the Jurassic basement of the Transcarpathian deep. *Paleont. zb.*, № 29, p. 61–65 (in Ukrainian).

18. **Ломизе М.Г.** Вулканизм флишевой геосинклинали Советских Карпат. *Вест. МГУ. Сер. геол.* 1975. № 2.

Lomize M.G., 1975. Volcanism of Flysch Geosyncline of the Soviet Carpathians. *Vestnik MGU. Ser. geol.*, № 2 (in Ukrainian).

19. **Объяснительная** записка к региональной стратиграфической схеме юрских отложений Предкарпатского прогиба и Волыно-Подольской окраины Восточно-Европейской платформы: Дулуб В.Г., Бурова М.И., Буров В.С., Вишняков И.Б. Ленинград: Мингео УССР, 1986. 58 с.

Explanatory note to regional stratigraphic scheme of Upper Jurassic deposits of Carpathian Foredeep and Volyno-Podolian edge of East European Platform: Dulub V.G., Burova M.I., Burov V.S., Vishnyakov I.B., 1986. Leningrad: Mingeo USSR, 58 p. (in Russian).

20. **Пояснювальна** записка до стратиграфічної схеми юрських відкладів Передкарпаття (Стрийський юрський басейн): Дулуб В.Г., Жабіна Н.М., Огородник М.Є., Смірнов С.Є. Львів: ЛВ УкрДГРІ, 2003. 32 с.

Explanatory note to regional stratigraphic scheme of Jurassic deposits of Pre-Carpathians (Stryi Jurassic Basin): Dulub V.G., Zhabina N.M., Ohorodnik M.E., Smirnov S.E., 2003. Lviv: LV UkrDGRI, 32 p. (in Ukrainian).

21. **Сандлер Я.М.** До характеристики юрських відкладів південно-західної частини Руської платформи і Передкарпатського прогину. *Геол. журн.* 1962. Т. 22, вип. 6. (87) С. 79–83.

Sandler J.M., 1962. By characteristics of the Jurassic sediments of the Southwestern part of Russian Platform and Carpathian Foredeep. *Geol. zh*, vol 22, N 6 (87), p. 79–83 (in Ukrainian).

22. **Славин В.И.** Схема стратиграфии мезозойских отложений Западных областей УССР. *Тр. Всесоюз. совещ. по разработке унифицированной схемы стратиграфии мезозойских отложений Русской платформы.* Ленинград: Гостоптехиздат, 1956. С. 87–100. (Тр. ВНИГРИ. Вып. 20).

Slavin V.I., 1956. Scheme of stratigraphy of Mesozoic deposits in the Western regions of the USSR. *Proc. all-Union Conf. to develop a Unified scheme of stratigraphy of Mesozoic deposits of the Russian Platform.* Leningrad: Gostoptekhizdat, p. 87–100. (Tr. VNIIGRI; Iss. 20) (in Russian).

23. **Славин В.И.** Триасовые и юрские отложения Восточных Карпат и Паннонского срединного массива. Москва: Госгеолиздат, 1963. 172 с.

Slavin V.I., 1963. Triassic and Jurassic deposits of the Eastern Carpathians and Pannonian Median Massif. Moscow: Gosgeolizdat, 172 p. (in Russian).

24. **Славин В.И., Добрынина В.Я.** Стратиграфия юрских отложений Львовской мульды и Предкарпатского прогиба. *Бюл. МОИП.* 1958. Т. 33, вып. 2. С. 43–54.

Slavin V.I., Dobrynina V.Y., 1958. The stratigraphy of the Jurassic deposits of the Lviv trough and Carpathian Foredeep. *Bul. MOIP*, vol. 33, № 2, p. 43–54 (in Russian).

25. **Славин В.И., Добрынина В.Я., Ефимова Н.А.** Новые данные о юрских отложениях Советских Карпат. *Карпато-Балкан. геол. ассоциация. VIII Конгр. Белград: доклады.* Киев, 1967. Т. 1. Ч. 2. С. 345–349.

Slavin V.I., Dobrynina V.Y., Efimova N.A., 1967. *New data on the Jurassic deposits of Soviet Carpathians. Carpathian-Balkan Geological Association. 8th Congr. Belgrade: Reports.* Kiev, vol. 1, pt. 2, p. 345–349 (in Russian).

26. **Стратиграфія** верхнього протерозою та фанерозою України: у 2 т. Т. 1: Стратиграфія верхнього протерозою, палеозою та мезозою України: Гожик П.Ф. (відп. ред.). Київ: Логос, 2014. 636 с.

Stratigraphy of Upper Proterozoic and Phanerozoic of Ukraine: in 2 volumes. Vol. 1. Stratigraphy of Upper Proterozoic, Paleozoic and Mesozoic of Ukraine: Gozhyk P.F. (Ed.), 2014. Kyiv: Logos, 636 p. (in Ukrainian).

27. **Стратиграфические** схемы фанерозоя и докембрия Украины / УМСК Украины. Гос. комитет Украины по геологии и использованию недр. Киев, 1993. 60 с.

Stratigraphic schemes of the Phanerozoic and Precambrian of Ukraine / UMSK of Ukraine. Gos. Committee of Ukraine on Geology and Use of Mineral Resources. Kyiv, 1993. 60 p. (in Russian).

28. **Утробин В.Н.** Основные черты стратиграфии юрских отложений Предкарпатского прогиба и юго-западной окраины Русской платформы. *Докл. АН СССР.* 1962. Т. 147, № 4. С. 908–911.

Utrobin V.N., 1962. The main features of the stratigraphy of the Jurassic deposits of Carpathian Foredeep and Southwestern edge of the Russian Platform. *Doklady AN SSSR*, vol. 147, № 4, p. 908–911 (in Russian).

29. **Rehakova D., Matyja B., Wierzbowski A., Schlogl J., Krobicki M., Barski M.** Stratigraphy and microfacies of the Jurassic and lowermost Cretaceous of the Veliky Kamenets section (Pieniny Klippen Belt, Carpathians, Western Ukraine). *Volumina Jurassica.* 2011. Vol. 9. P. 61–104.

30. **Wierzbowski A., Krobicki M., Matyja B.**, 2011. The stratigraphy and paleogeographic position of the Jurassic succession of the Priborzhavske-Perechin Zone in the Pieniny Klippen Belt of the Transcarpathian Ukraine. *Volumina Jurassica*, vol. 10, p. 25–60 (in English).

Стаття надійшла

13.11.2017