

UDC 547.74:547.232

A. H. Hotynchan¹, O. M. Kovtun², Yu. P. Kovtun¹

¹Institute of Organic Chemistry of the National Academy of Sciences of Ukraine,
5 Akademik Kukhar str., 02094 Kyiv, Ukraine

²Dragomanov Ukrainian State University, 9 Pyrohova str., 01601 Kyiv, Ukraine

The Synthesis of Pyrroles from Nitroolefins

Abstract

The synthesis of pyrroles occupies a key place in synthetic organic chemistry due to the numerous biological properties of pyrrole derivatives, in particular antimicrobial, antibacterial, antifungal, antimalarial, anticancer activities, etc. Therefore, pyrroles serve as building blocks in the creation of potential pharmaceuticals and also serve as the basis for the synthesis of boradipyrromethene dyes. One of the most well-known approaches to the synthesis of pyrroles is the reaction between nitroolefins, 1,3-dicarbonyl compounds, and amines, also known as the Grob-Camenisch reaction. This review is devoted to the historical chronology from the discovery of this transformation dating back to 1950s to the present, and covers the development of various modifications of the above reaction in the synthesis of pyrroles.

Keywords: pyrrole; Grob–Camenisch reaction; nitroolefins

А. Г. Готинчан¹, О. М. Ковтун², Ю. П. Ковтун¹

¹Інститут органічної хімії Національної академії наук України,
вул. Академіка Кухаря, 5, м. Київ, 02094, Україна

²Український державний університет імені Михайла Драгоманова,
вул. Пирогова, 9, м. Київ, 01601, Україна

Синтез піролів з нітроолефінів

Анотація

Синтез піролів займає ключове місце в синтетичній органічній хімії завдяки численним біологічним властивостям похідних піролу, зокрема антимікробній, антибактеріальній, протигрибковій, антималярійній, протираковій тощо. Саме тому піроли слугують будівельними блоками у створенні потенційних фармацевтичних препаратів, а також є основою для синтезу барвників бордипірометенового ряду. Одним із найвідоміших підходів до синтезу піролів є реакція між нітроолефінами, 1,3-дикарбонільними сполуками та амінами, відома як реакція Гроба-Каменіша. Цей огляд висвітлює історичну хронологію від відкриття цього перетворення, датованого 1950-ми роками, до сьогодення та охоплює розробку різноманітних модифікацій вищезгаданої реакції в синтезі піролів.

Ключові слова: пірол; реакція Гроба-Каменіша; нітроолефіни

Citation: Hotynchan, A. H.; Kovtun, O. M.; Kovtun, Yu. P. The synthesis of pyrroles from nitroolefins. *Journal of Organic and Pharmaceutical Chemistry* **2025**, 23 (2), 48–58.

<https://doi.org/10.24959/ophcj.25.323775>

Received: 26 February 2025; **Revised:** 4 June 2025; **Accepted:** 7 June 2025

Copyright © 2025, A. H. Hotynchan, O. M. Kovtun, Yu. P. Kovtun. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0>).

Funding: The authors received no specific funding for this work.

Conflict of interests: The authors have no conflict of interests to declare.

■ Introduction

Heterocyclic compounds are very important in the field of organic chemistry as they are widely spread in nature and commonly used as building blocks in pharmacology. Among heterocycles, pyrrole is of great interest due to its presence in a variety of medicines, such as atorvastatin [1],

tolmetin [2], ketorolac [3] etc., and its application in the dye synthesis, for example, BODIPY [4]. Therefore, the pyrrole core synthesis methods play a key role in contemporary organic chemistry.

There are some well-known synthetic routes for the pyrrole synthesis. The first one is the Hantzsch reaction, including the transformation between β -ketoesters, ammonia source, and

haloketones [5]. Alternatively, the Knorr synthesis (the reaction between α -amino-ketones and 1,3-dicarbonyl compounds) [6] or the Paal-Knorr synthesis (the reaction between 1,4-diketones and different ammonia sources) [7] can be employed. Due to increased interest, other methods of the pyrrole synthesis were developed. One of them is based on the *aza*-Wittig reaction, involving the interaction between 1-aza-1,3-bis(triphenylphosphoranylidene)propane and 1,2-diketo-compounds [8], and another one includes the reaction between 3-formylchromones and amines under TMSCl-mediated conditions [9].

Nitroolefins are an efficient initial material for the synthesis of organic compounds. Their synthetic attractiveness refers to the simplicity of their preparation *via* the condensation between nitroalkanes and aldehydes, known as the Henry reaction [10]. In β -nitrostyrene, the conjugation of the double bond with the nitro group enables it to participate in the Michael addition [11], a reaction that plays a crucial role in the synthesis of heterocycles. For example, nitroolefins are widely used in the synthesis of aziridines, pyrrolidines, oxazoles, indoles, etc. [12].

Taking the aforementioned information into account, the aim of this review is to discuss the historical development and progress in the pyrrole synthesis methods from nitroolefin derivatives.

Discovery of the Grob-Camenisch reaction

The first literary mention of the pyrroles synthesis from nitro compounds refers to the work of Grob and Kamenisch, dating back to 1953 [13].


Using 1-nitro-2-methylaminopropane and acetoacetic ester, the authors synthesized ethyl 1,2,4-trimethyl-2-pyrrolicarboxylate with a yield of 31%. They suggested the following mechanism (**Scheme 1**). In the solution, 1-nitro-2-methylaminopropane (**1**) exists in equilibrium with methylamine (**2**) and 1-nitropropene (**3**). After the addition of acetoacetic ester, the latter undergoes the condensation to form the corresponding enamine **4**, which further reacts with 1-nitropropene, resulting in Michael's product **5**. This product **5** cyclizes into dihydropyrrole **6**, eliminating a molecule of nitrous acid, which, in turn, oxidizes intermediate **6**, forming pyrrole **7**.

The further development of the reaction involving nitroolefins, 1,3-dicarbonyl compounds, and amines focused on identifying alternative synthetic approaches to increase pyrrole yields and expand the range of substrates that could be used. Currently, several variants of this reaction are known, including two-, three-, and four-component Grob-Camenisch-type reactions.


A two-component Grob-Camenisch-type reaction

The two-component Grob-Camenisch-type synthesis of pyrroles involves the reaction of β -enaminoesters or ketones with nitrostyrenes. First reported in 1981, this method utilized enamines **8**, derived from acetoacetic ester and (*E*)-(2-nitroprop-1-en-1-yl)benzene (**9**) [14]. Refluxing the reaction mixture in ethanol for four to six hours allows the pyrroles **10** to be obtained with yields ranging from 16% to 80% (**Scheme 2**).

The authors also found that using *N*-unsubstituted enamine **11** in the reaction with nitro



Scheme 1. The synthesis of pyrrole by Grob-Camenisch



Scheme 2. The synthesis of pyrroles from methyl- and cyclopropyl-derived enaminoesters

compound **9** by refluxing in ethanol for 14 hours could produce pyrrole **12** with a yield of 51% (**Scheme 3**).


Enaminoketones can also be used in a Grob-Camenisch-type reaction. However, harsher conditions are required, delivering the products with lower yields compared to standard substrates. Specifically, the reaction of enamine **13** with nitrostyrene **10** in a melt at 150°C produces pyrrole **14** with a yield of only 26% (**Scheme 4**) [14].

Further studies of the pyrrole synthesis from nitrostyrenes and β -enaminoesters or β -enaminoketones showed that enaminones **15** reacted with nitrostyrene or *p*-tolyl nitrostyrene (**16**) under solvent-free conditions, yielding pyrroles in 70–90% yield (**Scheme 5**) [15]. The mechanism


proposed involves the Michael-type addition of β -enaminones to nitroolefins, forming intermediate **17**, which cyclizes to pyrroline **18** with the elimination of nitrous acid. The subsequent oxidation results in the production of the final pyrrole **19**.

The scope and diversity of pyrroles synthesized from β -enaminones or esters and nitroolefins can be enhanced by using solvents or catalysts. For instance, the reaction of *N*-phenyl-substituted enamine **20** with substituted nitropropene **21** in methanol at 120°C yields penta-substituted pyrroles **22** (**Scheme 6**) [16].


The authors observed that catalytic amounts of bases (e.g., sodium acetate or piperidine) and polar aprotic solvents (e.g., DMSO or acetonitrile)




Scheme 3. The synthesis of pyrrole from *N*-unsubstituted enaminoester



Scheme 4. The synthesis of pyrrole from enaminone based on dimedone



Scheme 5. The reaction mechanism of the pyrrole formation in the reaction between enaminones and nitrostyrenes



Scheme 6. The synthesis of *N*-substituted pyrroles **22**

significantly decreased pyrrole yields. Electron-donating substituents on the aryl groups of nitrostyrenes and β -enaminoesters substantially improve yields compared to electron-withdrawing ones. It is noteworthy that the use of [2-nitroprop-1-en-1-yl]furan made it possible to synthesize pyrrole with a furyl group in position 3, yielding 79% of the product. Additionally, the application of nitrostyrene as a reagent facilitates the production of α -CH pyrrole with good yields.

Catalytic reactions provide an effective preparative method for synthesizing pyrroles from β -enaminoesters and nitroolefins. For instance, using iodine as a catalyst allows the formation of *N*-substituted pyrroles from nitrostyrene and β -enaminoesters derived from the acetoacetic ester. It is noteworthy that nitrostyrenes bearing electron-donating substituents result in pyrroles with significantly higher efficiency, whereas *N*-alkyl-substituted 3-aminobut-2-enoates lead to slightly lower yields due to competing side reactions [17].

The Ph_3PAuCl catalyst combined with AgOTf significantly enhances the pyrrole synthesis *via* a two-component reaction. Reactions of *N*-substituted enamines **23** obtained from acetoacetic ester or acetylacetone with substituted nitrostyrenes **24** lead to pyrroles **25** with yields exceeding 80% (**Scheme 7**) [18]. It is noteworthy


that the nature of substituents in aryl groups has minimal impact on pyrrole yields, and *N*-alkyl-substituted enamines react without side reactions, also giving pyrroles in high yields.

The PEG-400 catalyst demonstrated a high effectiveness for synthesizing isoxazole derivatives bearing a pyrrole moiety in position 4 [19]. The reaction of ethyl 3-((3-methyl-5-((*E*)-styryl)-isoxazol-4-yl)amino)but-2-enoate derivatives (**26**) with nitrostyrenes **27** gave the corresponding pyrroles **28** with yields from 70 to 90% (**Scheme 8**).


A solid-phase method for the pyrrole synthesis was also developed [20]. Initially, Rink Amide resin (**29**) is acetoacetylated with diketene to form amide **30**, which is subsequently converted to polymer-bound enaminone **31** upon the treatment with primary amines. Further reaction of polymer-bound **31** with nitroolefins in a DMF/EtOH solvent mixture at 60°C yielded pyrroles **32**, which upon the treatment with trifluoroacetic acid produced final amides **33**. In this method, the pyrrole yield exceeds 80% when using both aliphatic and aromatic amines, as well as aliphatic nitroolefins (**Scheme 9**).

A three-component Grob-Camenisch-type reaction


The three-component Grob-Camenisch synthesis of pyrroles involves the reaction of amines, 1,3-dicarbonyl compounds, and nitrostyrenes,



Scheme 7. The reaction mechanism of the pyrrole formation in the reaction between enamines and nitrostyrenes catalyzed with Ph_3PAuCl



Scheme 8. The synthesis of pyrroles 28



Scheme 9. The solid-phase synthesis of pyrroles from nitroolefins

which make it possible to prepare pyrroles with catalytic activity. For instance, research [21] demonstrates that lactic acid serves as an effective medium for synthesizing tetrasubstituted pyrroles from acetylacetone and substituted anilines, achieving yields of approximately 70–90%. The study indicates that neither electron-withdrawing nor electron-donating substituents on the aniline significantly influence pyrrole yields.

Iron-based catalysts, such as FeCl_3 , are widely employed in the catalytic synthesis of pyrroles from nitroolefins and 1,3-dicarbonyl compounds. For instance, FeCl_3 facilitates a three-component reaction, yielding pyrroles at 70–80% using 1,3-dicarbonyl compounds like acetoacetic ester, acetylacetone, or ethylbenzyl acetate [22]. As a Lewis acid, FeCl_3 promotes the formation of β -enaminoester **34**, which undergoes the Michael-type addition to nitrostyrene, followed by the cyclization of the resulting adduct **35** into pyrrole *via* the nitro group conversion to its aci-nitro form (Scheme 10).


Interestingly, iron(III) chloride makes it possible to synthesize pyrroles from nitrostyrene bearing ethynyl substituents in the *ortho*-position,

obtaining the target products with a yield of about 50% [23]. Additionally, the catalyst facilitates the pyrrole synthesis from peptides with a free amino group [24].


An alternative to FeCl_3 is the use of Fe_3O_4 nanoparticles, in particular $\text{Fe}_3\text{O}_4@\text{SiO}_2\text{-CPTMS-guanidine-SO}_3\text{H}$, which contain a terminal sulfo group [24]. This catalyst allows the synthesis of tetrasubstituted pyrroles from acetoacetic ester, acetylacetone, and *para*-chloro- or *para*-bromoanilines with yields of approximately 90% (Scheme 11). The sulfo group is supposed to catalyze the reaction similarly to FeCl_3 .

Cerium(III) chloride serves as an alternative catalyst to FeCl_3 , effective in the microwave-assisted synthesis using nitromethane as a solvent [26]. This method allows the synthesis of tetrasubstituted pyrroles from acetoacetic ester or acetylacetone, nitrostyrenes, and anilines, with yields of about 80%. However, when *para*-chloronitrostyrene is used, the yield decreases to about 50%.

An alternative to CeCl_3 is cerium(IV) ammonium nitrate (CAN), which catalyzes the reaction of nitrostyrene **38**, acetoacetic ester (**36**),



Scheme 10. The reaction mechanism of the three-component synthesis of pyrroles from nitroolefins catalyzed with FeCl_3



Scheme 11. The reaction mechanism of the three-component synthesis of pyrroles from nitroolefins catalyzed with $\text{Fe}_3\text{O}_4@\text{SiO}_2\text{-CPTMS-guanidine-SO}_3\text{H}$


and benzylamine (**37**) at room temperature in methanol [27]. This method produces pyrroles with yields ranging from 50 to 75% (**Scheme 12**).

In addition to other metal-containing catalysts, Ph_3PAuCl combined with AgOTf [28] and zirconyl dichloride complexes [29] makes it possible to use acetoacetic ester and various β -diketoamides as 1,3-dicarbonyl compounds, yielding pyrroles at approximately 70%.

Furthermore, diacetoxyiodobenzene serves as an effective non-metal catalyst for the pyrrole syn-

thesis, producing pyrroles from acetoacetic ester and acetylacetone with yields of about 70%, unaffected by substituents on nitrostyrene or aniline [30, 31].

The combination of oxone and iodobenzene makes it possible to synthesize pyrroles from nitrostyrenes, acetylacetone, or acetoacetic ester, and anilines, yielding 80–90% of the desired product. Notably, neither electron-donating nor electron-withdrawing substituents on nitrostyrene or aniline impact the pyrrole yield [31].



Scheme 12. The reaction mechanism of the three-component synthesis of pyrroles from nitroolefin **38**

Ionic liquids, such as *N*-methyl-2-pyrrolidonium methyl sulfonate, serve as effective catalysts for the three-component synthesis of pyrrole involving nitroolefins, amines, and 1,3-dicarbonyl compounds. Notably, the yields of pyrroles from substituted nitrostyrenes and anilines, ranging from 70 to 90%, show no correlation with the substituent effects. However, using aliphatic nitroolefins and amines decreases yields to 30% [32].

A four-component Grob-Camenisch-type reaction


Another convenient option for the synthesis of pyrroles is a four-component reaction involving aldehydes, nitroalkanes, 1,3-dicarbonyl compounds, and amines. Currently, only catalytic variants of this transformation have been developed using FeCl_3 [32], palladium [34], and tungsten-

based complexes [35], CuO nanoparticles [36], ionic liquids [37, 38], iodine [39], clay [40], $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ [41], as well as organic acids – lactic and gluconic [42] (**Scheme 13**).

The reaction mechanism includes the catalyzed formation of a β -enaminone from a nitroolefin, followed by the Michael addition to form an intermediate. This intermediate undergoes elimination of a hyponitrous acid molecule, resulting in the formation of pyrrole.

The synthesis of *N*-unsubstituted α -CH pyrroles

The synthesis of *N*-H α -CH pyrroles is of significant interest in the Grob cyclization, as these compounds serve as precursors for luminescent BODIPY borofluoride complexes and porphyrins. The earliest reported attempt to synthesize such



Scheme 13. The reaction mechanism of the four-component synthesis of pyrroles from nitroolefins catalyzed by Lewis acids


pyrroles was by Grob [13] where 1-nitropropan-2-amine **39** and acetoacetic ester were used. However, instead of the expected N-H pyrrole, an *N*-substituted derivative **44** was obtained. This outcome was attributed to intermediate **41** possessing two nucleophilic centers: a carbon atom and a nitrogen one. Hence, two Michael additions occur, leading to intermediate **42**, which undergoes further cyclization, thus yielding *N*-isopropyl derivative **44** instead of the expected N-H one (Scheme 14).

In order to synthesize the corresponding N-H α -CH pyrrole, Grob first isolated intermediate **45**, which failed to transform into pyrrole **46** under different conditions (Scheme 15). The author attributed this to the conjugation of the nitrogen atom lone pair in intermediate **45** with the ethoxycarbonyl group reducing its nucleophilicity. However, interestingly, upon the treatment of **45** with methylamine, pyrrole **48** could be isolated, assuming that the transamination


occurred with the formation of **47**, which then cyclized to **48**. From this observation, it could be suspected that the presence of even weak electron-donating groups favors the pyrrole synthesis by the Grob cyclization.

In the subsequent study, Gómez-Sánchez [43] investigated the synthesis of N-H α -CH pyrroles using nitro compounds, with a focus on the formation of anomalous Michael products from nitrostyrene and acetylacetone. By employing non-catalytic amounts of sodium methoxide with acetylacetone and nitrostyrene **49**, the authors obtained anomalous Michael products **50** (Scheme 16). The latter product, upon the treatment with saturated methanolic ammonia solution at 0°C, gives rise to pyrrole **51** (Scheme 16).



Based on this observation, the authors developed a one-pot procedure of the pyrrole synthesis, including the treatment of nitrostyrene **49** with acetylacetone, acetoacetic ester, or methyl acetoacetate in methanol with a non-catalytic




Scheme 14. The synthesis of pyrrole by Grob-Camenisch



Scheme 15. Transformations of intermediate **45**

Scheme 16. The synthesis of anomalous Michael's product **50** and pyrrole **51**

Scheme 17. The one-pot pyrrole synthesis by Gómez-Sánchez



Scheme 18. The synthesis of other pyrrole derivatives by the Gómez-Sánchez-like procedure

amount of sodium methoxide at 0°C for 1 hour with further addition of an ammonia source, such as aniline, benzylamine or saturated methanolic ammonia solution. This synthetic procedure allowed obtaining pyrroles **52** with yields of up to 80% (Scheme 17).

Two additional literature references describe similar methods for synthesizing N-H α -CH pyrroles, involving the initial formation of the Michael addition product between nitrostyrenes **53** and acetoacetic ester, followed by the treatment with ammonia (Scheme 18). Using this approach, pyrroles **54** were obtained, though in low yields, from 25 to 30% [44, 45].

Conclusions

The Grob-Camenisch synthesis of pyrroles is highly valued for its straightforward methods and procedures, as evidenced by numerous publications. The main advantage of this reaction is the possibility of obtaining tetra- and three-substituted pyrrole derivatives. This reaction also allows the synthesis of pyrroles in multi-gram amounts. It is noteworthy that for this method, it is possible to use catalysts of different types (Lewis and Brønsted acids, nanoparticles, etc.), which broadens the synthetic possibilities of the pyrrole synthesis.

References

- Novozhilov, Y. V.; Dorogov, M. V.; Blumina, M. V.; Smirnov, A. V.; Krasavin, M. An Improved Kilogram-Scale Preparation of Atorvastatin Calcium. *Chem. Cent. J.* **2015**, *9* (1). <https://doi.org/10.1186/s13065-015-0082-7>.
- Artico, M.; Corelli, F.; Massa, S.; Stefancich, G. Non-steroidal antiinflammatory agents. A novel synthesis of 1-methyl-5-p-tolylpyrrole-2-acetic acid (tolmetin). *J. Het. Chem.* **1982**, *19* (6), 1493–1495. <https://doi.org/10.1002/jhet.5570190647>.
- Müller, P.; Polleux, Ph. Synthesis of a Ketorolac Model via Aromatic Carbenoid Insertion. *Helv. Chim. Acta.* **1998**, *81* (2), 317–323. <https://doi.org/10.1002/hlca.19980810212>.
- Ulrich, G.; Zissel, R.; Harriman, A. The Chemistry of Fluorescent Bodipy Dyes: Versatility Unsurpassed. *Angew. Chem. Int. Ed.* **2008**, *47* (7), 1184–1201. <https://doi.org/10.1002/anie.200702070>.
- Hantzsch, A. Neue Bildungsweise von Pyrrolderivaten. *Ber. Dtsch. Chem. Ges.* **1890**, *23* (1), 1474–1476. <https://doi.org/10.1002/cber.189002301243>.
- Knorr, L. Synthese von Pyrrolderivaten. *Ber. Dtsch. Chem. Ges.* **1884**, *17* (2), 1635–1642. <https://doi.org/10.1002/cber.18840170220>.
- Paal, C. Synthese von Thiophen- und Pyrrolderivaten. *Ber. Dtsch. Chem. Ges.* **1885**, *18* (1), 367–371. <https://doi.org/10.1002/cber.18850180175>.
- Katritzky, A. R.; Jiang, J.; Steel, P. J. 1-Aza-1,3-bis(triphenylphosphoranylidene)propane: A Novel :CHCH₂N: Synthone. *J. Org. Chem.* **1994**, *59*, 4551. <https://doi.org/10.1021/jo00095a034>.

9. Plaskon, A. S.; Ryabukhin, S. V.; Volochnyuk, D. M.; Shivanyuk, A. N.; Tolmachev, A. A. The Synthesis of 5-Hetaryl-3-(2-Hydroxybenzoyl) Pyrroles. *Tetrahedron* **2008**, *64* (25), 5933–5943. <https://doi.org/10.1016/j.tet.2008.04.041>.
10. González-Olvera, R.; Vergara-Arenas, B. I.; Negrón-Silva, G. E.; Angeles-Beltrán, D.; Lomas-Romero, L.; Gutiérrez-Carrillo, A.; Lara, V. H.; Morales-Serna, J. A. Synthesis of β -Nitrostyrenes in the Presence of Sulfated Zirconia and Secondary Amines. *RSC Adv.* **2015**, *5* (120), 99188–99192. <https://doi.org/10.1039/c5ra17168g>.
11. Stowe, G. N.; Janda, K. D. A Diels-Alder Reaction Conducted within the Parameters of Aqueous Organocatalysis: Still Just Smoke and Mirrors. *Tetrahedron Lett.* **2011**, *52* (17), 2085–2087. <https://doi.org/10.1016/j.tetlet.2010.10.134>.
12. Halimehjani, A. Z.; Namboothiri, I. N. N.; Hooshmand, S. E. Part I: Nitroalkenes in the Synthesis of Heterocyclic Compounds. *RSC Adv.* **2014**, *4* (89) 48022–48084. <https://doi.org/10.1039/c4ra08828j>.
13. Grob, C. A.; Camenish, K. Eine neue Pyrrolring-Synthese. *Helv. Chim. Acta.* **1953**, *36* (1), 49–58. <https://doi.org/10.1002/hlca.19530360109>.
14. Meyer, H. Dihydropyridine, VI. Diimidomalonsäurederivate in der Hantzsch-Pyridin-Synthese. *Liebigs Ann. Chem.* **1981**, *1981* (9), 1523–1533. <https://doi.org/10.1002/jlac.198119810902>.
15. Yavari, I.; Ghazvini, M.; Aminkhani, A. Solvent-Free Synthesis of 1,2,3,4-Tetrasubstituted Pyrroles from Enaminones and β -Nitrostyrenes. *J. Chem. Res.* **2011**, *35* (10), 558–560. <https://doi.org/10.3184/174751911X13164434098696>.
16. Guan, Z. H.; Li, L.; Ren, Z. H.; Li, J.; Zhao, M. N. A Facile and Efficient Synthesis of Multisubstituted Pyrroles from Enaminoesters and Nitroolefins. *Green Chem.* **2011**, *13* (7), 1664–1668. <https://doi.org/10.1039/c1gc15278e>.
17. Xu, H.; Li, Y.; Xing, M.; Jia, J.; Han, L.; Ye, Q.; Gao, J. Synthesis of Pyrroles from β -Enamines and Nitroolefins Catalyzed by I_2 under High-Speed Vibration Milling (HSVM). *Chem. Lett.* **2015**, *44* (4), 574–576. <https://doi.org/10.1246/cl.141102>.
18. Abdulkader, A.; Xue, Q.; Lin, A.; Zhang, M.; Cheng, Y.; Zhu, C. Gold-Catalyzed Cascade C-C and C-N Bond Formation: Synthesis of Polysubstituted Indolequinones and Pyrroles. *Tetrahedron Lett.* **2013**, *54* (44), 5898–5900. <https://doi.org/10.1016/j.tetlet.2013.08.100>.
19. Ponduri, R.; Kumar, P.; Vadali, L. R. PEG-400 Promoted a Simple, Efficient, and Recyclable Catalyst for the One-Pot Eco-Friendly Synthesis of Functionalized Isoxazole Substituted Pyrroles in Aqueous Medium. *Synth. Commun.* **2018**, *48* (24), 3113–3122. <https://doi.org/10.1080/00397911.2018.1535078>.
20. Trautwein, A. W.; Jung, G. Solid-phase synthesis of pyrroles from enaminones and nitroalkenes. *Tetrahedron Lett.* **1998**, *39*, 8263–8266. [https://doi.org/10.1016/S0040-4039\(98\)01887-5](https://doi.org/10.1016/S0040-4039(98)01887-5).
21. Akbaslar, D.; Giray, E. S.; Algul, O. Revisit to the Synthesis of 1,2,3,4-Tetrasubstituted Pyrrole Derivatives in Lactic Acid Media as a Green Solvent and Catalyst. *Mol. Divers.* **2021**, *25* (4), 2321–2338. <https://doi.org/10.1007/s11030-020-10122-1>.
22. Sarkar, S.; Bera, K.; Maiti, S.; Biswas, S.; Jana, U. Three-Component Coupling Synthesis of Diversely Substituted N-Aryl Pyrroles Catalyzed by Iron(III) Chloride. *Synth. Commun.* **2013**, *43* (11), 1563–1570. <https://doi.org/10.1080/00397911.2011.650273>.
23. Sarkar, S.; Bera, K.; Jalal, S.; Jana, U. Synthesis of Structurally Diverse Polyfunctional Pyrrolo[1,2-a]Quinolines by Sequential Iron-Catalyzed Three-Component Coupling and Gold-Catalyzed Hydroarylation Reactions. *Eur. J. Org. Chem.* **2013**, *27*, 6055–6061. <https://doi.org/10.1002/ejoc.201300659>.
24. Jad, Y. E.; Gudimella, S. K.; Govender, T.; De La Torre, B. G.; Albericio, F. Solid-Phase Synthesis of Pyrrole Derivatives through a Multicomponent Reaction Involving Lys-Containing Peptides. *ACS Comb. Sci.* **2018**, *20* (4), 187–191. <https://doi.org/10.1021/acscombsci.8b00006>.
25. Rostami, H.; Shiri, L. $Fe_3O_4@SiO_2$ —CPTMS—Guanidine— SO_3H -Catalyzed One-Pot Multicomponent Synthesis of Polysubstituted Pyrrole Derivatives under Solvent-Free Conditions. *Russian J. Org. Chem.* **2019**, *55* (8), 1204–1211. <https://doi.org/10.1134/S1070428019080207>.
26. Silveira, C. C.; Mendes, S. R.; Martins, G. M.; Schlösser, S. C.; Kaufman, T. S. Modular $CeCl_3 \cdot 7H_2O$ -Catalyzed Multi-Component Synthesis of 1,2,3,4-Tetrasubstituted Pyrroles under Microwave Irradiation and Their Further Trichloroisocyanuric Acid-Mediated Conversion into 5-Sulphenylpyrrole Derivatives. *Tetrahedron* **2013**, *69* (43), 9076–9085. <https://doi.org/10.1016/j.tet.2013.08.035>.
27. Magar, D. R.; Ke, Y. J.; Chen, K. Three-Component Synthesis of Functionalized N-Protected Tetrasubstituted Pyrroles by an Addition-Elimination-Aromatization Process. *Asian J. Org. Chem.* **2013**, *2* (4), 330–335. <https://doi.org/10.1002/ajoc.201200193>.
28. Li, L.; Chen, Q.; Xiong, X.; Zhang, C.; Qian, J.; Shi, J.; An, Q.; Zhang, M. Synthesis of Polysubstituted Pyrroles via a Gold(I)-Catalyzed Tandem Three-Component Reaction at Room Temperature. *Chinese Chem. Lett.* **2018**, *29* (12), 1893–1896. <https://doi.org/10.1016/j.ccllet.2018.09.004>.
29. Goyal, S.; Patel, J. K.; Gangar, M.; Kumar, K.; Nair, V. A. Zirconocene Dichloride Catalyzed One-Pot Synthesis of Pyrroles through Nitroalkene-Enamine Assembly. *RSC Adv.* **2015**, *5* (5), 3187–3195. <https://doi.org/10.1039/c4ra09873k>.
30. Jadhav, N. C.; Pahalkar, A. R.; Desai, N. V.; Telvekar, V. N. Design, Synthesis and Molecular Docking Study of Novel Pyrrole-Based α -Amylase and α -Glucosidase Inhibitors. *Med. Chem. Res.* **2017**, *26* (10), 2675–2691. <https://doi.org/10.1007/s00044-017-1965-z>.
31. Jadhav, N. C.; Jagadhane, P. B.; Patile, H. V.; Telvekar, V. N. Three-Component Direct Synthesis of Substituted Pyrroles from Easily Accessible Chemical Moieties Using Hypervalent Iodine Reagent. *Tetrahedron Lett.* **2013**, *54* (23), 3019–3021. <https://doi.org/10.1016/j.tetlet.2013.04.014>.
32. Pachecne, L. A.; Pereira, V. F.; Martins, G. M.; Martendal, E.; Xavier, F. R.; Mendes, S. R. One-Pot Multicomponent Synthesis of 1,2,3,4-Tetra-substituted Pyrroles Catalyzed by $[NMPH]CH_3SO_3$. *Tetrahedron Lett.* **2019**, *60* (38), 151043. <https://doi.org/10.1016/j.tetlet.2019.151043>.
33. Maiti, S.; Biswas, S.; Jana, U. Iron(III)-Catalyzed Four-Component Coupling Reaction of 1,3-Dicarbonyl Compounds, Amines, Aldehydes, and Nitroalkanes: A Simple and Direct Synthesis of Functionalized Pyrroles. *J. Org. Chem.* **2010**, *75* (5), 1674–1683. <https://doi.org/10.1021/jo902661y>.
34. Reddy, G. R.; Reddy, T. R.; Joseph, S. C.; Reddy, K. S.; Reddy, L. S.; Kumar, P. M.; Krishna, G. R.; Reddy, C. M.; Rambabu, D.; Kapavarapu, R.; Lakshmi, C.; Meda, T.; Priya, K. K.; Parsa, K. V. L.; Pal, M. Pd-Mediated New Synthesis of Pyrroles: Their Evaluation as Potential Inhibitors of Phosphodiesterase 4. *Chem. Commun.* **2011**, *47* (27), 7779–7781. <https://doi.org/10.1039/c1cc12321a>.
35. Atar, A. B.; Jeong, Y. T. Heterogenized Tungsten Complex: An Efficient and High Yielding Catalyst for the Synthesis of Structurally Diverse Tetra Substituted Pyrrole Derivatives via Four-Component Assembly. *Tetrahedron Lett.* **2013**, *54* (41), 5624–5628. <https://doi.org/10.1016/j.tetlet.2013.08.016>.
36. Saeidian, H.; Abdoli, M.; Salimi, R. One-Pot Synthesis of Highly Substituted Pyrroles Using Nano Copper Oxide as an Effective Heterogeneous Nanocatalyst. *C. R. Chim.* **2013**, *16* (11), 1063–1070. <https://doi.org/10.1016/j.crci.2013.02.008>.
37. Meshkatsadat, M. H.; Mahmoudi, A.; Lotfi, S.; Pouramiri, B.; Foroumadi, A. Green and Four-Component Cyclocondensation Synthesis and in Silico Docking of New Polyfunctionalized Pyrrole Derivatives as the Potential Anticholinesterase Agents. *Mol. Div.* **2022**, *26* (6), 3021–3035. <https://doi.org/10.1007/s11030-021-10362-9>.
38. Meshram, H. M.; Madhu Babu, B.; Santosh Kumar, G.; Thakur, P. B.; Bangade, V. M. Catalyst-Free Four-Component Protocol for the Synthesis of Substituted Pyrroles under Reusable Reaction Media. *Tetrahedron Lett.* **2013**, *54* (19), 2296–2302. <https://doi.org/10.1016/j.tetlet.2013.01.098>.

39. Reddy, G. R.; Reddy, T. R.; Joseph, S. C.; Reddy, K. S.; Pal, M. Iodine Catalyzed Four-Component Reaction: A Straightforward One-Pot Synthesis of Functionalized Pyrroles under Metal-Free Conditions. *RSC Adv.* **2012**, 2 (8), 3387–3395. <https://doi.org/10.1039/c2ra00982j>.
40. Bharate, J. B.; Sharma, R.; Aravinda, S.; Gupta, V. K.; Singh, B.; Bharate, S. B.; Vishwakarma, R. A. Montmorillonite Clay Catalyzed Synthesis of Functionalized Pyrroles through Domino Four-Component Coupling of Amines, Aldehydes, 1,3-Dicarbonyl Compounds and Nitroalkanes. *RSC Adv.* **2013**, 3 (44), 21736–21742. <https://doi.org/10.1039/c3ra43324b>.
41. Khan, A. T.; Lal, M.; Ray Bagdi, P.; Sidick Basha, R.; Saravanan, P.; Patra, S. Synthesis of Tetra-Substituted Pyrroles, a Potential Phosphodiesterase 4B Inhibitor, through Nickel(II) Chloride Hexahydrate Catalyzed One-Pot Four-Component Reaction. *Tetrahedron Lett.* **2012**, 53 (32), 4145–4150. <https://doi.org/10.1016/j.tetlet.2012.05.133>.
42. Li, B. Le; Li, P. H.; Fang, X. N.; Li, C. X.; Sun, J. L.; Mo, L. P.; Zhang, Z. H. One-Pot Four-Component Synthesis of Highly Substituted Pyrroles in Gluconic Acid Aqueous Solution. *Tetrahedron* **2013**, 69 (34), 7011–7018. <https://doi.org/10.1016/j.tet.2013.06.049>.
43. Gómez-Sánchez, A.; Stiefel, B. M.; Fernández-Fernández, R.; Bellanato, J. Unusual Michael Reaction of Acyclic 1,5-Dicarbonyl Compounds with Nitro-Olefins. A Novel Pyrrole Synthesis. *J. Chem. Soc., Perkin Trans.*, **1982**, 1, 441–447. <https://doi.org/10.1039/P19820000441>.
44. Yamamoto, S.; Matsunaga, N.; Hitaka, T.; Yamada, M.; Hara, T.; Miyazaki, J.; Santou, T.; Kusaka, M.; Yamaoka, M.; Kanzaki, N.; Furuya, S.; Tasaka, A.; Hamamura, K.; Ito, M. Design, Synthesis, and Biological Evaluation of 4-Phenylpyrrole Derivatives as Novel Androgen Receptor Antagonists. *Bioorg. Med. Chem.* **2012**, 20 (1), 422–434. <https://doi.org/10.1016/j.bmc.2011.10.067>.
45. Okamoto, M.; Kojima, H.; Saito, N.; Okabe, T.; Masuda, Y.; Furuya, T.; Nagano, T. Virtual Screening and Further Development of Novel ALK Inhibitors. *Bioorg. Med. Chem.* **2011**, 19 (10), 3086–3095. <https://doi.org/10.1016/j.bmc.2011.04.008>.

Information about the authors:

Andrii H. Hotynchan (*corresponding author*), Ph.D. student of the Department of Colour and Structure of Organic Compounds, Institute of Organic Chemistry of the National Academy of Sciences of Ukraine. <https://orcid.org/0000-0002-1569-8590>; e-mail for correspondence: ahotynchan@gmail.com.

Olena M. Kovtun, Associate Professor of the Chemistry Department, Natural Faculty, Dragomanov Ukrainian State University.

Yuriy P. Kovtun, Dr. Sci. in Chemistry, Professor, Leading Researcher of the Department of Colour and Structure of Organic Compounds, Institute of Organic Chemistry of the National Academy of Sciences of Ukraine. <https://orcid.org/0000-0002-2673-5342>.